

ANNEX F

DNA repair and mutagenesis

CONTENTS

	<i>Page</i>
INTRODUCTION	2
I. DNA DAMAGE AND REPAIR	2
A. THE ROLE OF DNA REPAIR GENES IN CELL FUNCTION	2
B. TYPES OF DAMAGE AND PATHWAYS OF REPAIR	3
C. SUMMARY	8
II. REPAIR PROCESSES AND RADIOSENSITIVITY	9
A. RADIOSENSITIVITY IN MAMMALIAN CELLS AND HUMANS	9
1. The identification of radiosensitive cell lines and disorders	9
2. Mechanisms of enhanced sensitivity in human disorders	11
3. Analysis of genes determining radiosensitivity	12
4. Summary	17
B. RELATIONSHIP BETWEEN REPAIR AND OTHER CELL REGULATORY PROCESSES	18
1. Radiosensitivity and defective recombination in the immune system: non-homologous end joining of DNA double-strand breaks	18
2. Radiosensitivity and the cell cycle	21
3. Apoptosis: an alternative to repair?	23
4. Summary	25
III. HUMAN RADIATION RESPONSES	26
A. CONTRIBUTION OF MUTANT GENES TO HUMAN RADIOSENSITIVITY	26
B. INFLUENCE OF REPAIR ON RADIATION RESPONSES	29
C. SUMMARY	32
IV. MECHANISMS OF RADIATION MUTAGENESIS	33
A. MUTATION AS A REPAIR-RELATED RESPONSE	33
B. THE SPECTRUM OF RADIATION-INDUCED MUTATIONS	35
C. MOLECULAR ANALYSIS OF RADIATION-INDUCED MUTATIONS	36
D. EFFECT OF RADIATION QUALITY	39
E. NOVEL MECHANISMS OF GENETIC CHANGE	41
F. MUTATION FREQUENCIES AND CONSEQUENCES	44
G. SUMMARY	47
CONCLUSIONS	47
<i>References</i>	49

INTRODUCTION

1. Risk estimates for the induction of human disease are obtained primarily from epidemiological studies. These studies can clearly distinguish radiation effects only at relatively high doses and dose rates. To gain information at low doses and low dose rates, which are more relevant to typical human radiation exposures, it is necessary to extrapolate the results of these studies. To be valid, this extrapolation requires a detailed understanding of the mechanisms by which radiation induces cancer and genetic disorders.

2. Several lines of evidence show that sites of radiation-induced cell lethality, mutation, and malignant change are situated within the nucleus and that DNA is the primary target. When DNA is damaged by radiation, enzymes within the cell nucleus attempt to repair that damage. The efficiency of the enzymatic repair processes determines the outcome: most commonly, the structure of DNA is repaired correctly and cellular functions return to normal. If the repair is unsuccessful, incomplete, or imprecise, the cell may die or may suffer alteration and loss of genetic information (seen as mutation and chromosomal aberration). These information changes determine heritable genetic defects and are thought to be important in the development of radiation-induced cancer. The more complete the knowledge of the ways in which human cells respond to damage and of the mechanisms underlying the formation of mutations and chromosomal aberrations, the more accurate will be the predictions of the oncogenic and hereditary effects of ionizing radiation.

3. DNA repair is itself controlled by a specific set of genes encoding the enzymes that catalyse cellular response to DNA damage. Loss of repair function, or alteration of the control of repair processes, can have very serious consequences for cells and individuals. It is anticipated that DNA repair plays a critical role in protecting normal individuals from radiation effects, including cancer. Clinical experience has revealed individuals who are both hypersensitive to radiation and cancer-prone; some of these individuals have recently been shown to have defects in genes involved in the response to DNA damage.

4. In recent years there have been significant advances in the molecular analysis of repair processes and the understanding of the mechanisms that induce genetic changes. Additionally, new methods have been developed to simplify the identification of the genes involved. As the details of damage-repair processes become clearer, it is seen that these processes have considerable overlap with other cellular control functions, such as those regulating the cell cycle and immune defences. In this Annex the Committee continues to review such developments in molecular radiobiology, as it began to do in Annex E, "Mechanisms of radiation carcinogenesis" of the UNSCEAR 1993 Report [U3], in order to improve the understanding of how radiation effects are manifested in cells and organisms.

I. DNA DAMAGE AND REPAIR

A. THE ROLE OF DNA REPAIR GENES IN CELL FUNCTION

5. The information needed to control cellular functions such as growth, division, and differentiation is carried by the genes. Genes, which are specific sequences of DNA, act mainly through the production of complementary messages (mRNA) that are translated into proteins. Proteins can have a structural role but commonly work as enzymes, each of which catalyses a particular metabolic reaction. Thus specific genes contain the code for (encode) specific cellular functions. The production of proteins can be timed so that they work at specific points in the development of a cell or organism, but protein function can also be controlled by post-translational modifications. These modifications are carried out by other proteins, so that a complex set of interactions is necessary to fine-tune cellular functions. Proteins involved in important aspects of cell metabolism (e.g. DNA replication) may also work in multi-protein complexes [A1]. There is some evidence also that some of the complexes are assembled into larger structures situated in defined regions of the nucleus (e.g. the nuclear matrix) [H2].

6. Loss or alteration of information in a specific gene may mean that none of that gene product (protein) is formed, or that the protein is less active, or that it is formed in an uncontrolled fashion (e.g. at the wrong time or in the wrong amount). While some minor genetic alterations may not affect protein activity or interactions, others may significantly disrupt cellular function. Since certain proteins work in a number of different processes or complexes, the loss or impairment of one type of protein can affect several different functions of the cell and organism (pleiotropic effect).

7. A very large number of genes, 60,000-70,000 [F14], are required to control the normal functions of mammalian cells and organisms. However, the genes form only a small part of the genome (the complete DNA sequence of an organism), the remainder of which largely consists of many copies of repetitive DNA sequence. The genes are linked in linear arrays interspersed by non-coding sequences, to form chromosomes located in the cell nucleus. Most genes are present in only two copies, each on a separate homologous chromosome, one inherited from the mother and one from the father. To monitor damage and to maintain the genes without significant alteration is a major concern for the cell. Repair

processes are common to all organisms from bacteria to humans and have evolved to correct errors made in replicating the genes and to restore damaged DNA. This fact has in recent years provided a useful tool for molecular geneticists in the analysis of repair processes; well characterized micro-organisms can serve as model systems to understand the structure and function of repair genes. The information gained in this way can sometimes also be used directly to isolate human genes of related function [L1]. While the structure and function of repair genes appears to be highly conserved from lower to higher organisms, the regulation of their activities may differ in different organisms.

8. The consequences of loss of repair capacity are seen in a number of human syndromes and mutant cell lines. These show hypersensitivity to environmental agents, and the human syndromes often have multiple symptoms, including cancer-proneness, neurological disorder, and immune dysfunction. Good progress has been made over the last few years in mapping and cloning the genes involved.

9. Ionizing radiation damages DNA and causes mutation and chromosomal changes in cells and organisms. Damage by radiation or radiomimetic agents also leads to cell transformation (a stage in cancer development) and cell death. In the light of current research, it is seen that the final response to radiation damage is determined not only by cellular repair processes but also by related cellular functions that optimize the opportunity for recovery from damage. For example, radiation damage may cause an arrest in the cell cycle; this is thought to be a damage-limitation step, allowing time for repair and reducing the consequences of a given dose [L22]. There is now some understanding of the way in which radiation alters cell cycle timing (Section II.B.2), although the roles of a number of enzymatic activities that are induced or repressed shortly after irradiation remain to be clarified (Section III.B).

10. The severity of DNA damage, or the context in which damage occurs (e.g. during DNA replication), will often dictate a repair strategy that places survival first and incurs genetic change. DNA replication may bypass sites of single-stranded DNA damage, inserting an incorrect base opposite the altered or lost base. Additionally, in attempting to repair damage to DNA, enzymes may not be able to restore the structure with fidelity. Thus, mutation and chromosomal rearrangement are not passive responses to damage; rather, they are a consequence of the interaction of cellular processes with damage. The types of genetic change that occur will depend on the types of initial DNA damage, from their potential to miscode at replication and from the probability that specific repair enzymes will act on given types of damage.

B. TYPES OF DAMAGE AND PATHWAYS OF REPAIR

11. DNA is a double-helical macromolecule consisting of four units: the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T) and cytosine (C). The bases

are arranged in two linear arrays (or strands) held together by hydrogen bonds centrally and linked externally by covalent bonds to sugar-phosphate residues (the DNA "backbone"). The adenine base pairs naturally with thymine (A:T base pair), while guanine pairs with cytosine (G:C base pair), so that one DNA strand has the complementary sequence of the other. The sequence of the bases defines the genetic code; each gene has a unique sequence, although certain common sequences exist in control and structural DNA elements. Damage to DNA may affect any one of its components, but it is the loss or alteration of base sequence that has genetic consequences.

12. Ionizing radiation deposits energy in tracks of ionizations from moving charged particles within cells, and radiations of different quality may be arbitrarily divided into sparsely ionizing, or low linear energy transfer (low-LET), and densely ionizing (high-LET). Each track of low-LET radiations, such as x rays or gamma rays, consists of only a relatively small number of ionizations across an average-sized cell nucleus (e.g. a gamma-ray electron track crossing an 8 μm diameter nucleus gives an average of about 70 ionizations, equivalent to about 1 mGy absorbed dose, although individual tracks vary widely about this value because of track stochastics and varying path lengths through the nucleus). Each track of a high-LET radiation may consist of many thousands of ionizations and give a relatively high dose to the cell; for example, a 4 MeV alpha-particle track has, on average, about 23,000 ionizations (370 mGy) in an average-sized cell nucleus [G27, U3]. However, within the nucleus even low-LET radiations will give some small regions of relatively dense ionization over the dimensions of DNA structures, for example, where a low-energy secondary electron comes to rest within a cell.

13. Radiation tracks may deposit energy directly in DNA (direct effect) or may ionize other molecules closely associated with DNA, especially water, to form free radicals that can damage DNA (indirect effect). Within a cell the indirect effect occurs over very short distances, of the order of a few nanometres, because the diffusion distance of radicals is limited by their reactivity. Although it is difficult to measure accurately the different contributions made by the direct and indirect effects to DNA damage caused by low-LET radiation, evidence from radical scavengers introduced into cells suggests that about 35% is exclusively direct and 65% has an indirect (scavengeable) component [R21]. It has been argued that both direct and indirect effects cause similar early damage to DNA; this is because the ion radicals produced by direct ionization of DNA may react further to produce DNA radicals similar to those produced by water-radical attack on DNA [W43].

14. Ionization will frequently disrupt chemical bonding in cellular molecules such as DNA, but where the majority of ionizations occur as single isolated events (low-LET radiations), these disruptions will be readily repaired by cellular enzymes. However, the average density of ionization by high-LET radiations is such that several ionizations are likely to occur as the particle traverses a DNA double helix.

Therefore, much of the damage from high-LET radiations, as well as a minority of the DNA damage from low-LET radiations, will derive from localized clusters of ionizations that can severely disrupt the DNA structure [G27, W44]. While the extent of local clustering of ionizations in DNA from single tracks of low- and high-LET radiations will overlap, high-LET radiation tracks are more efficient at inducing larger clusters, and hence more complex damage. Also, high-LET radiations will induce some very large clusters of ionizations that do not occur with low-LET radiations; the resulting damage may be irreparable but may also have unique cellular consequences (see paras. 192, 199, and 201) [G28]. Additionally, when a cell is damaged by high-LET radiation, each track will give large numbers of ionizations, so that the cell will receive a relatively high dose and there will be a greater probability of correlated damage within a single DNA molecule (or chromosome) or in separate chromosomes. As a consequence, the irradiation of a population of cells or a tissue with a "low dose" of high-LET radiation results in a few cells being hit with a relatively high dose (one track) rather than in each cell receiving a small dose. In contrast, low-LET radiation is more uniformly distributed over the cell population; at doses of low-LET radiation in excess of about 1 mGy (for an average-size cell nucleus of 8 μm diameter), each cell nucleus is likely to be traversed by more than one sparsely-ionizing track.

15. The interaction of ionizing radiation with DNA produces numerous types of damage; the chemical products of many of these have been identified and classified according to their structure [H4, S3]. These products differ according to which chemical bond is attacked, which base is modified, and the extent of the damage within a given segment of DNA. Table 1 lists some of the main damage products that can be measured following low-LET irradiation of DNA, with a rough estimate of their abundance. Attempts have also been made to predict the frequencies of different damage types from a knowledge of radiation track structure, with certain assumptions about the minimum energy deposition (number of ionizations) required. Interactions can be classified

according to the probability they will cause a single-strand DNA alteration (e.g. a break in the backbone or base alteration) or alterations in both strands in close proximity in one DNA molecule (e.g. a double-strand break), or a more complex type of DNA damage (e.g. a double-strand break with adjacent damage). Good agreement has been obtained between these predictions and direct measurements of single-strand breaks, but there is less good agreement for other categories of damage [C47]. While complex forms of damage are difficult to quantify with current experimental techniques, the use of enzymes that cut DNA at sites of base damage suggests that irradiation of DNA in solution gives complex damage sites consisting mainly of closely-spaced base damage (measured as oxidised bases or abasic sites); double-strand breaks were associated with only 20% of the complex damage sites [S87]. It is expected that the occurrence of more complex types of damage will increase with increasing LET, and that this category of damage will be less repairable than the simpler forms of damage. Theoretical simulations have predicted that about 30% of DNA double-strand breaks from low-LET radiation are complex by virtue of additional breaks [N19] and that this proportion rises to more than 70%, and the degree of complexity increases, for high-LET particles [G29].

16. Some of the DNA damage caused by ionizing radiation is chemically similar to damage that occurs naturally in the cell: this "spontaneous" damage arises from the thermal instability of DNA as well as endogenous oxidative and enzymatic processes [L2, M40]. Several metabolic pathways generate oxidative radicals within the cell, and these radicals can attack DNA to give both DNA base damage and breakage, mostly as isolated events [B46]. The more complex types of damage caused by ionizing radiation may not occur spontaneously, since localized concentrations of endogenous radicals are less likely to be generated in the immediate vicinity of DNA. This theme is taken up in Annex G, "*Biological effects at low radiation doses*", which considers the cellular responses to low doses of radiation.

Table 1
Estimated yields of DNA damage in mammalian cells caused by low-LET radiation exposure
[L60, P31, W39]

<i>Type of damage</i>	<i>Yield (number of defects per cell Gy⁻¹)</i>
Single-strand breaks	1 000
Base damage ^a	500
Double-strand breaks	40
DNA-protein cross-links	150

^a Base excision enzyme-sensitive sites [P31] or antibody detection of thymine glycol [L60].

17. Measurement of the endogenous levels of DNA base damage has been difficult because of the artefactual production of damage during the preparation of the DNA for analysis (e.g. by gas chromatography/mass spectrometry) [C55]. This difficulty explains the presence in the literature of considerably inflated (by factors of at least 100) values for background

levels of base damage. Interestingly, the recognition of damage by base excision repair enzymes (paragraph 22) has provided a less discordant method of measurement, although the specificity of the enzymes for different types of base damage is not precisely known. These enzymes cut the DNA at the site of base damage, to give a single-strand break that

can be measured accurately by a number of techniques. Using this method, measurement of an important form of oxidative damage, 7,8-dihydro-8-oxoguanine (generally known as 8-oxoguanine)), has given steady-state levels of 500-2000 per cell, depending on cell type [P30]. Using similar measurement methods, the level of 8-oxoguanine induced in cellular DNA by gamma rays is about 250 per cell per Gy [P31]. A newly developed ultrasensitive assay for another type of base damage in human cellular DNA, thymine glycol, couples antibody detection with capillary electrophoresis. This method showed a linear response for yield of thymine glycol with gamma-ray dose down to 0.05 Gy, giving a level of about 500 thymine glycols per cell per Gy against a background of 6 thymine glycols per cell [L60]. The difficulties experienced in measuring base damage accurately in cellular DNA and the relatively low levels now found for the commoner types of damage have also called into question the extent to which some previously identified forms of base damage occur in cells following irradiation.

18. The measurement of endogenous levels of other types of DNA damage, such as double-strand breaks, has involved similar technical difficulties. Many of the methods used to measure double-strand breaks in mammalian cells introduce this form of damage either inadvertently or deliberately as part of the methodology. This is because the mammalian genome is so large that it had to be reduced in size by random breakage first before useful measurements could be made. This problem has been overcome in part by the introduction of methods based on the gentle release of DNA from cells by their lysis in a gel matrix [C64, O3], but there is commonly still a background level of DNA breakage amounting to a few per cent of the total DNA. However, as documented in Section II.B, it is unlikely that mammalian cells have a high steady-state level of DNA double-strand breakage, since these breaks act as a signal for damage-recognition processes that can block the cell cycle or induce programmed cell death. It is possible that even one unrepaired double-strand break can trigger this cellular response (paragraph 101). It has also been found that one unrepaired double-strand break can cause lethality in irradiated yeast cells (paragraph 108). Thus, tolerance of this form of damage in cells is likely to be very low.

19. While the precise nature of the damage will influence repairability, it is possible to consider a few general categories of damage in order to describe their consequences. A simplified classification can be based on the ability of enzymes to use the complementary base structure of DNA to facilitate repair of the damage site. Thus, damage to single strands (base modifications, single-strand breaks) can be removed or modified, followed by resynthesis using the undamaged strand as a template. Where the damage affects both strands of a DNA molecule in close proximity (double-strand breaks, cross-links), it is more difficult to repair and requires different enzymatic pathways for its resolution. To resolve successfully more complex types of damage may require enzymes from more than one repair pathway. To illustrate the knowledge of the different repair pathways available to the cell, the following account (to paragraph 34) includes a discussion of

the repair of damage caused by various DNA-damaging agents as well as ionizing radiation.

20. DNA repair enzymes can be characterized as cellular proteins acting directly on damaged DNA in an attempt to restore the correct DNA sequence and structure. These relatively specialized enzymes appear to undertake the initial stages of recognition and repair of specific forms of DNA damage. For example, DNA glycosylases catalyze the cleavage of base-sugar bonds in DNA, acting only on altered or damaged bases [W1]. Further, there are several different types of glycosylase that recognize chemically different forms of base damage. However, enzymes that carry out normal DNA metabolism are also part of the repair process for many different forms of damage. In the latter category there are, for example, enzymes involved in the synthesis of DNA strands (DNA polymerases) and enzymes involved in the joining of the DNA backbone (DNA ligases). Several different types of DNA polymerases and ligases have been identified; it is thought that they have different roles in normal DNA metabolism and that only some are active in DNA repair [L3, P1].

21. The simplest repair processes directly reverse the damage; for example, many organisms, but not mammals, possess an enzyme that directly photoreactivates the UV-induced dimerization of pyrimidine bases [S1]. Similarly, the enzyme O⁶methylguanine-methyltransferase directly removes methyl groups induced in DNA by alkylating carcinogens [P13]. However, most damage types require the concerted action of a number of enzymes, forming a repair pathway. Several apparently discrete repair pathways have been identified, as described below and illustrated in Figure I.

22. Damage to individual bases in DNA may be corrected simply by removing the base, cleaning up the site, and resynthesis. In this process, termed the base-excision repair pathway, a DNA glycosylase removes the damaged base, a DNA endonuclease cuts the DNA backbone, the sugar-phosphate remnants are removed by a phosphodiesterase, and a polymerase fills in the gap using the opposite base as a template (Figure Ia) [L2, S49]. Even where a single base is damaged, therefore, several different enzymes are required to give correct repair. The latter part of this process may also be used to repair single-strand breaks in DNA. Radiation-induced DNA breaks are generally not rejoined by a simple ligation step, because sugar damage and, often, base loss occur at the site of a break. Base-excision repair is generally localized to the single DNA base and is very rapid [D16, S58]; however, in mammalian cells a minority of repair patches of up to 6 bases have been found, indicating a second "long-patch" pathway (see paragraph 75).

23. Many DNA glycosylases are specific for the removal of one type of altered base from DNA; for example, uracil-DNA glycosylase removes only uracil and some oxidation products of uracil [F15]. However, there is some overlap in the specificity of some base-excision repair enzymes. An




Figure 1.
Mechanisms of DNA repair (simplified).

Figure 1a: Base damage is excised by a specific glycosylase: the DNA backbone is cut and the gap filled by a polymerase. The resulting gap is refilled.

Figure 1b: Bulky base damage is removed along with an oligonucleotide (of about 30 bases in human cells). Resynthesis takes place using the opposite strand as a template.

Figure 1c: A double-strand break is rejoined end-to-end.

Figure 1d: A double-strand break is repaired with the help of a homologous undamaged molecule (shown in red). Strand invasion allows resynthesis on complementary sequence, followed by a resolution of the strands and rejoining.

example of this is seen in the response to ionizing radiation of bacterial cells defective for one or more nucleases involved in base-excision repair: while cells defective for either nuclease alone were not radiation sensitive, the loss of both nucleases made the cells extremely radiation sensitive [C60, Z10]. Some glycosylases have a very broad specificity for different types of base structure, such as 3-methyladenine-DNA glycosylase II, which acts on a variety of chemically-modified purines and pyrimidines, 5-methyloxidized thymines, and hypoxanthine. It has recently been found that 3-methyladenine-DNA glycosylase II will also remove natural bases from DNA, especially purines, at significant rates. This finding has led to

the novel suggestion that the rate of excision of broad-specificity glycosylases is a function of the chemical bond stability in DNA. On this basis, damaged bases are excised more readily than natural bases because their chemical bonds are less stable [B57].

24. A specialized form of base-excision repair involves the removal of mismatched DNA bases that occur as errors of DNA replication or from the miscoding properties of damaged bases. For example, 8-oxoguanine is a common product of oxidative damage to guanine bases (paragraph 17); this product is highly mutagenic because of its ability to miscode

(polymerases incorporate adenine instead of cytosine opposite 8-oxoguanine, thus changing the DNA sequence). Cells have evolved three different methods to deal with the formation of 8-oxoguanine: one glycosylase can correct the mismatch by removing 8-oxoguanine from DNA, while a different glycosylase can remove the mismatched adenine after DNA replication [A27, T48]. Excision and resynthesis of the missing base occur as in the general base-excision repair process. Additionally, another enzyme can remove the precursor of 8-oxoguanine (8-hydroxy-GTP) from the nucleotide pool before it is incorporated into DNA [M51]. A number of other mismatches, arising most commonly in DNA replication and recombination, are corrected by a separate "long patch" pathway, known as the MutHLS pathway, that is similarly important in protecting cells from high frequencies of mutation (see paragraph 165) [K38].

25. Where a damaged base is close to another damaged base or to a single-strand break, which is the most simple form of clustered damage, repair may be compromised. Examination of model DNA substrates with base damage on opposite strands, separated by different numbers of bases, has shown that glycosylases do not repair both sites of damage when they are very close to one another (1-3 bases) [C56, C57, H49]. Further, attempted repair of both sites of base damage can result in a DNA double-strand break, because the DNA backbone is cut as part of the repair process.

26. In contrast to base-excision repair, nucleotide-excision repair removes a whole section of single-stranded DNA containing a site of damage, generally a bulky DNA adduct causing distortion of the double helix. These enzymes have to perform such functions as recognition of damage, cutting of the strand at a specified distance either side of the damage, and unwinding and removal of the strand (Figure 1b). As might be expected, at least 11 enzymes have already been identified as components of nucleotide-excision repair [W52], not including polymerase and ligase. The enzymes of nucleotide-excision repair are highly conserved from microbes to humans, and this feature has been used to assist in the isolation of the genes encoding these functions. Several of these genes have been found to be mutated in humans, giving rise to a series of disorders, including xeroderma pigmentosum and Cockayne's syndrome [H3]. Individuals inheriting mutated nucleotide-excision repair genes are generally sensitive to sunlight and chemical agents causing bulky damage in DNA, but a few individuals show cross-sensitivity to ionizing radiation [A2, R15]. Additionally, a small fraction of damage induced by gamma rays is not repaired in cells derived from individuals with xeroderma pigmentosum, which suggests that ionizing radiation induces some bulky damage (e.g. purine dimers) that cannot be removed by the base-excision repair pathway [S56]. Alternatively, a fraction of non-bulky base damage (8-oxoguanine, thymine glycol) may be removed by the nucleotide-excision repair system, particularly in long-lived cells such as neurons that sustain a great deal of endogenous oxidative damage [R22]. These possibilities may explain why severe cases of xeroderma pigmentosum also suffer progressive neurological degeneration.

27. A surprising recent discovery was that some nucleotide-excision repair enzymes are also involved in the normal process of gene expression (transcription). Thus, when genes are actively expressing, they require some of the same functions needed for repair, such as unwinding the DNA helix, and it seems that the same proteins are used. This finding explains the previously puzzling observation of a link between human disorders with sensitivity to sunlight and those with complex defects in gene expression (such as trichothiodystrophy [L70]).

28. Another important discovery about nucleotide-excision repair is that it operates at different rates in different parts of the genome [H1]. Thus, actively expressing genes are repaired much faster than the remainder of the genome. Much of the detail of this process has now been elucidated: it is thought that when damage occurs in a gene that is actively expressing, the proteins involved in this process (the RNA polymerase II transcription complex) stop working, and the stalled complex acts as a signal to the repair proteins to go to the damage site. This two-tier repair system has been found in organisms from bacteria to humans, and the protein mediating the signal to bring repair to the damage site has been identified in bacteria [S2]. The presence in the cell of a fast transcription-coupled repair process has also been found to have genetic consequences: only one of the DNA strands of the duplex is transcribed, and only this strand is repaired rapidly. It has been found in normal cells that most of the mutations induced by DNA damage are in the non-transcribed strand, presumably because lack of fast repair allows the damage to interact with other processes, causing mutations [M2]. In contrast, repair-defective cell lines show a completely altered mutation spectrum, with most mutations recovered in the transcribed strand. Much of this detail has been established using UV-light damage, but the repair of other types of DNA damage, including certain forms of base damage induced by ionizing radiation such as thymine glycols, is influenced by transcription-coupled processes [H1, C41]. Again, certain human sun-sensitive disorders have been found to lack either the fast repair path (Cockayne's syndrome) or the slower overall path of nucleotide-excision repair (xeroderma pigmentosum group C). Loss of the fast or slower pathways may also affect the clinical outcome of sun-sensitive disorders: Cockayne's syndrome does not result in cancer-proneness, while xeroderma pigmentosum patients are highly prone to skin cancers [M2].

29. More severe forms of damage require yet more resources for their correct repair. This is especially true of damage affecting both DNA strands simultaneously, since there is no undamaged strand to act as a template for repair. Severe damage may occur directly by a damaging agent causing complex DNA changes or may arise during the replication of unrepaired single-stranded DNA damage. It is likely that such severe damage will be repaired by recombination enzymes, which rejoin or replace damaged sequences through a variety of mechanisms. In general, there are two main types of recombination repair processes: homologous recombination and illegitimate recombination, although site-specific recombination processes also occur. The principles of recombination repair have been well

established in micro-organisms, and it has recently been found that similar processes occur in human cells.

30. Homologous recombination takes advantage of the sequence identity between certain regions of DNA to repair damage; such regions exist, for example, in the maternal and paternal copies of chromosomes and in the duplicated chromosome (sister chromatids) following DNA replication. The DNA sequence, from which information is derived to repair the damaged copy, must be identical over a considerable length (≥ 200 base pairs). It is known in the budding yeast (*Saccharomyces cerevisiae*), for example, that homologous recombination is the main method for DNA double-strand break repair. Several of the *rad52* group of yeast mutants were isolated on the basis of their extreme sensitivity to ionizing radiation and have been shown to be defective in both DNA double-strand break repair and homologous recombination [G1]. In recombination, the broken 3' end of a DNA strand invades an unbroken double-stranded homologue, and resynthesis on this template re-forms the damaged strand (Figure Id). Separation of the joint product of this reaction requires the activity of enzymes cutting and rejoining the newly-synthesized DNA strands. Depending on which strands are cut and rejoined, this reaction may also result in crossing over (genetic exchange) of DNA strands.

31. Illegitimate recombination (including DNA end-joining processes) is a common mechanism for rejoining broken DNA sequences in mammalian cells (Figure Ic). When foreign DNA molecules are integrated into the genome [R1] or when the genomic breakpoints of deletions and rearrangements are analysed (Section IV.C), it is found that these genomic sites show little sequence homology. There appears to be more than one repair pathway involved, and terms such as non-homologous end joining and direct-repeat end joining are used to describe different pathways in this Annex (see Section II.B.1). It can be argued that illegitimate recombination is a mechanism for rapidly rejoining broken DNA ends without the need for the complex machinery of homologous recombination [R1]. It is also likely that because of the large amount of repetitive DNA sequence in mammalian cells, if the processes of homologous recombination were generally available in cells, there would be an intolerable level of reshuffling of the genome. While homologous recombination is thought to be a mechanism for repairing DNA with little error, illegitimate recombination is likely to cause alteration and/or loss of DNA sequence.

32. It is likely that both homologous and illegitimate recombination processes are able to repair severe damage in the genome. However, a surprising recent discovery in mammalian cells is that some of the enzymes involved in repairing radiation-induced breakage of DNA also take part in a site-specific recombination process, V(D)J immune-system recombination. This process assembles functional immune genes from separate genomic regions, through somatic gene rearrangement, and is dealt with in more detail in Section II.B.1.

33. There is also evidence that cells have specific surveillance mechanisms for DNA damage and that these mechanisms interface with other aspects of cellular metabolism such as cell-cycle progression [M3]. Thus it is envisaged that when the genome, and perhaps other parts of the cell, sustains damage, a response mechanism is set up to maximize the chance of repairing the damage (or in some cases to commit the cell to a programmed death). The details of these mechanisms, as well as how the overall response is coordinated, are not yet clear.

34. More than 50 genes are already known to affect the repair of DNA damage in lower eukaryotic organisms such as yeasts [F1], but this figure includes genes involved in processes such as cell-cycle checkpoints (Section II.B.2). Additionally, new genes are being found continually, both in searches for homologues of existing repair genes and in genome mapping projects. In view of the numbers already discovered, the multiplicity of types of damage requiring repair, and the recent discoveries of complexity in repair pathways, it would not be surprising if the overall number in humans is a few hundred genes. Therefore, a significant fraction of the genome (paragraph 7) is devoted to maintaining the integrity of DNA. Since the damage to DNA from ionizing radiation is also very diverse, many of these genes will play a role in its repair.

C. SUMMARY

35. Ionizing radiation interacts with DNA to give many different types of damage. Radiation track structure considerations indicate that the complexity of the damage increases with linear energy transfer, and that this complexity may distinguish radiation damage from alterations occurring spontaneously and by other agents. Attempts to measure endogenous levels of damage have suffered from high levels of artefacts, and despite improved methods there are still large margins of error in these estimates. At present, therefore, it is difficult to compare radiation-induced levels of damage with those occurring spontaneously, especially when damage complexity is taken into account. The importance of the relationship between spontaneous and induced levels of damage in the determination of low dose responses is taken up in Annex G, "*Biological effects at low radiation doses*".

36. A large number of genes have evolved in all organisms to repair DNA damage; the repair gene products operate in a co-ordinated fashion to form repair pathways that control restitution of specific types of damage. Repair pathways are further co-ordinated with other metabolic processes, such as cell cycle control, to optimize the prospects of successful repair.

37. It is likely that the simpler forms of DNA damage (single sites of base damage, single-strand breaks) arising endogenously and from exposure to ionizing radiation will be repaired rapidly and efficiently by base-excision repair processes, so these types of damage are not normally a

serious challenge to biological organisms. However, because of the relatively large amount of base damage and single-strand breaks induced (Table 1), if base-excision repair systems are compromised, the consequences would be very serious for the cell and the individual. DNA damage such as double-strand breaks represents a more difficult problem for cellular repair processes, but more

than one recombination repair pathway has evolved to cope with this damage. Damage caused by large clusters of ionizations in or near DNA, giving more complex DNA alterations, may represent a special case for which separate repair pathways have to come together to effect repair, or where there is a consequent loss or alteration of the DNA sequence as a result of incorrect or inadequate repair.

II. REPAIR PROCESSES AND RADIOSENSITIVITY

A. RADIOSENSITIVITY IN MAMMALIAN CELLS AND HUMANS

1. The identification of radiosensitive cell lines and disorders

38. Individuals vary in their sensitivity to ionizing radiation. Highly radiosensitive individuals have been detected when they present for cancer therapy; these are seen as rare patients suffering severe normal tissue damage after standard therapy treatments. It has been possible to group some radiosensitive patients into defined disorders, such as ataxia-telangiectasia and the Nijmegen breakage syndrome, but others appear to be asymptomatic (that is, with none of the symptoms of known sensitivity disorders, but also discovered following treatment for cancer). Additionally there are individuals who show less extreme radiosensitivity, some of whom may be variants of known disorders such as ataxia-telangiectasia.

39. Ataxia-telangiectasia is the best described of radiosensitive disorders. It has a complex phenotype; cerebellar ataxia, neuromuscular degeneration, dilated ocular blood vessels (telangiectasia), immunodeficiency, chromosomal instability, and a substantially increased incidence of some cancers and neoplasms are common to ataxia-telangiectasia patients [B10]. Ataxia-telangiectasia is inherited primarily as an autosomal recessive trait, although it has been suggested that both radiosensitivity and cancer-proneness behave with some dominance. The disease is progressive, with most affected individuals surviving only to adolescence or early adulthood. Lymphocytic leukaemia and non-Hodgkin's lymphoma appear to be the commonest forms of cancer, but solid tumours in various organs are also associated with ataxia-telangiectasia [H9]. Estimates of the frequency of the disorder vary but suggest an average of about 1 per 100,000 [P8, S12, W10].

40. The radiosensitive phenotype of ataxia-telangiectasia is also readily demonstrated in cells cultured from patients, using cell survival and chromosome damage assays. For example in a survey comparing the survival of cells cultured from 42 normal individuals with those from 10 ataxia-telangiectasia individuals following x-irradiation, the ataxia-telangiectasia cells were, on average, 2.7 times more sensitive than the normal cells (see Figure II) [C8]. Compared to normal cells, an elevated frequency of chromosomal aberrations is found both spontaneously and after irradiation of ataxia-telangiectasia

cells. Also, while irradiation of normal cells in the pre-synthesis (G_0) phase of the cell cycle yields only chromosome-type aberrations, both chromatid- and chromosome-type aberrations are found in ataxia-telangiectasia [T1]. A striking feature of ataxia-telangiectasia cells is their resistance to radiation-induced DNA synthesis delay: normal cells show a rapid inhibition of DNA synthesis after irradiation, while ataxia-telangiectasia cells have a delayed and/or much reduced inhibition [P4].




Figure II. Survival of human fibroblast cells after x-irradiation as measured by their colony-forming ability [C8]. The range of D_0 (the dose required to kill 63% of the cells) is 0.3–0.6 Gy in 10 patients with ataxia-telangiectasia and 1–1.6 Gy in 42 normal individuals.

41. Nijmegen breakage syndrome is a clinically separate radiosensitive disorder characterized by variable immune deficiencies, microcephaly, developmental delay, chromosomal instability, and cancer susceptibility [B4, S7, V7, W4]. Lymphoreticular cancers again seem to characterize this disorder [S8]. Nijmegen breakage syndrome patients show no ataxia or telangiectasia, but their cellular phenotype is very similar to that of ataxia-telangiectasia [A17, J1, N6, T2]. Other patients with similarities to ataxia-telangiectasia and Nijmegen breakage syndrome have been found and in some cases classified separately by genetic analysis ([C12, W5]; see also below). One case has been reported of combined ataxia-telangiectasia and Nijmegen breakage syndrome and called A-T_{FRESNO} [C18]. Also, there are a number of reports of families or individuals who show symptoms that partially overlap

ataxia-telangiectasia and Nijmegen breakage syndrome; an example would be a family with individuals showing either ataxia-telangiectasia or a disorder involving ataxia, microphthalmia, and congenital cataract [Z3].

42. To add to this complexity, a number of individuals with a variant form of ataxia-telangiectasia have been described; these patients may show a slower onset of symptoms and have intermediate levels of cellular radiosensitivity [C4, C7, F2, J2, T3, Y1, Z1].

43. In addition to individuals having multiple symptoms associated with radiosensitivity, some otherwise normal persons have been found to be highly radiosensitive. Woods et al. [W9] described an apparently normal 13-year-old girl who developed multiple complications following radiotherapy for Hodgkin's disease. Cells derived from a skin biopsy showed a highly sensitive radiation survival response, similar to that for ataxia-telangiectasia cells. Plowman et al. [P7] reported similar findings for a 14-year-old boy with acute lymphoblastic leukaemia; again, no ataxia-telangiectasia or Nijmegen breakage syndrome-like symptoms were present, but both whole-body and cellular radiosensitivity were as extreme as for ataxia-telangiectasia. This individual has now provided the first example of radiosensitivity in humans where a defined cellular repair defect (in the ligase IV enzyme) can be readily demonstrated; repair of radiation-induced DNA double-strand breaks and interphase chromosome damage are similarly impaired (see also paragraph 66) [B15].

44. There is some evidence for ionizing radiation sensitivity in several other cancer-prone disorders, although the published data do not always agree on the degree of sensitivity (see also Section III.A). Bloom's syndrome is a rare autosomal recessive disorder showing severe growth retardation, variable immune deficiencies, and abnormal spermatogenesis [G20]. The age of onset of cancer is considerably earlier than normal; about one third of surviving cancer patients with Bloom's syndrome develop multiple primary tumours with no consistent pattern of cancer type or location. Individuals with Bloom's syndrome develop a distinct facial rash from sunlight sensitivity, and their cells are not only hypersensitive to several different DNA-damaging agents but also show DNA replication abnormalities [L47]. Genetic instability is seen in high levels of spontaneously-occurring chromosomal aberrations and sister chromatid exchanges; chromosomal sensitivity to x rays has been found especially in cells in the G₂ phase of growth [A16, K30]. One case of Bloom's syndrome developed oesophageal stricture following standard radiotherapy treatment for lung cancer. This is very rarely seen in such treatment and is suggestive of hypersensitivity to radiation [K31].

45. Fanconi's anaemia is a cancer-prone disorder, most commonly presenting with acute myeloid leukaemia (15,000-fold increased risk), although solid tumours are also found. Bone marrow failure is a common diagnostic feature of this disorder, although symptoms may include congenital malformations, abnormal skin pigmentation, and skeletal and renal abnormalities [J15]. Fanconi's anaemia cells show high levels of chromosomal aberrations and are hypersensitive to

DNA cross-linking agents (e.g. mitomycin C, diepoxybutane). Additionally, a high proportion of deletions has been reported in certain genes of Fanconi's anaemia cell lines, giving a higher frequency of mutation in assays measuring loss of heterozygosity (see paragraph 172) [S68]. There has been some dispute over the extent to which Fanconi's anaemia cells are sensitive to ionizing radiation; a lack of genetic classification in these experiments may account for some of the variability found (see paragraph 68). However, when they compared published data on the sensitivity of human fibroblasts, Deschavanne et al. [D7] concluded that Fanconi's anaemia was one of the few disorders for which sensitivity to radiation could be distinguished from that of normal cells.

46. The genes controlling radiosensitivity in humans will not be identified simply by analysing radiosensitive disorders. This is because mutations in many genes are deleterious to the extent that the development of a viable organism is inhibited. This point was illustrated by the creation of a "knockout" mouse for the UV-damage repair gene *ERCC1* (knockout meaning that both copies of the gene are inactivated). No human variant for the *ERCC1* gene has been found, and the knockout mouse dies before weaning, apparently as the result of a massive load of (unrepaired) damage [M7]. Therefore, to examine the full range of genes involved, it has been necessary to derive radiosensitive mutant lines from cells in culture. To this end, more than 50 mutant cell lines sensitive to various genotoxic agents have been identified; many of these show some degree of x-ray sensitivity and are being used to dissect repair pathways in mammalian cells [C16, H11]. These cell lines are especially useful for gene cloning, since this has often proved difficult to achieve using cells derived from human patients. It is possible at the present stage of knowledge to group these mutant cell lines into several categories based on their responses, and recently several of the genes involved have been mapped or cloned (Table 2). As radiosensitive cell lines have been developed in laboratories around the world, almost all have been found to represent defects in different genes. For this reason, and because ionizing radiation produces a diversity of DNA damage, it is anticipated that a large number of the human genes involved in determining radiation resistance remain undiscovered.

47. The discovery and analysis of the genetic basis of radiation sensitivity is also being pursued through other strategies, including the biochemical analysis of repair reactions and the purification of repair proteins, as well as the identification of human repair genes by homology to their counterparts in lower organisms (Section II.A.3). In many instances, the genes discovered by these routes are found to give rise to a high level of sensitivity when mutated but to affect only a small fraction of the human population or to be inconsistent with life. However, some genes affecting radiation sensitivity will probably have more subtle effects, either because the particular gene mutation only partially reduces gene product activity or because the gene is not vital for cellular response to radiation. Studies exploring the latter types of response, which may affect a much larger fraction of the human population, are described in Section III.A.

Table 2
Classification of radiosensitive disorders and cell lines

<i>Type of defect</i>	<i>Disorder / defective cell line(s)</i>	<i>Human gene designation^a</i>	<i>Human gene location</i>	<i>Animal model phenotype^b</i>
Probable DNA break repair defect and loss of cell-cycle control following damage	Ataxia telangiectasia Nijmegen breakage syndrome <i>irs 2/V-series</i>	<i>ATM</i> <i>NBS1</i> <i>XRCC8</i>	11q23 8q21 ?	AT-like ^c - -
DNA double-strand break repair defective and V(D)J recombination defective	<i>xrs</i> XR-1/M10 V3/ <i>scid/SX9</i> 180BR	<i>XRCC5</i> <i>XRCC4</i> <i>XRCC7</i> <i>LIG4</i>	2q35 5q13 8p11-q11 13q33-q34	Immune deficiency ^d Embryonic lethal Immune deficiency ^d Embryonic lethal
Sensitivity to many different agents; some have DNA single-strand break repair defect and/or replication defect	46BR Bloom's syndrome EM9 <i>irs1</i> <i>irs1SF</i> UV40	<i>LIG1</i> <i>BLM</i> <i>XRCC1</i> <i>XRCC2</i> <i>XRCC3</i> <i>XRCC9(FANCG)</i>	19q13 15q26 19q13 7q36 14q32 9p13	Viable (acute anaemia) Embryonic lethal Embryonic lethal Embryonic lethal - -
Radiosensitivity inferred from structural homology of genes to those known to be involved in response to radiation damage in lower organisms	- - - - -	<i>ATR</i> <i>hRAD50</i> <i>hRAD51</i> <i>hRAD52</i> <i>hRAD54</i> <i>hMRE11</i>	3q22-23 5q23-31 15q 12p13.3 1p32 11q21	Embryonic lethal Embryonic lethal Embryonic lethal Viable Viable Embryonic lethal

a XRCC = x-ray cross complementing gene.

b Knockout mice except for *XRCC7*; (-) indicates no model yet available

c Symptoms similar to ataxia telangiectasia (see paragraph 62).

d Severe combined immune deficiency.

2. Mechanisms of enhanced sensitivity in human disorders

48. In addition to being radiosensitive, ataxia-telangiectasia cell lines show enhanced sensitivity to agents that have in common an ability to damage DNA molecules by producing highly reactive chemical radicals, causing both base damage and sugar damage, leading to breakage of DNA strands. Ataxia-telangiectasia cells have been found to be hypersensitive to a variety of chemicals that cause such DNA damage through radical action (bleomycin, neocarzinostatin, hydrogen peroxide, streptonigrin, phorbol ester [M5]). Also, inhibitors of DNA topoisomerases that can trap these enzymes during DNA-strand passage, leaving open breaks, are more effective at inducing chromosomal damage and cell killing in ataxia-telangiectasia cells than in normal cells [C1, H7, S11]. More recently, ataxia-telangiectasia cell lines have been found to be hypersensitive to restriction endonucleases; these enzymes produce only double-strand breaks in DNA by direct enzymatic cutting [C17, L56]. There is also evidence of modest chromosomal hypersensitivity in some ataxia-telangiectasia lines to agents such as UV light, especially when irradiated in extended G₁ phase, possibly because of the excessive production of breaks when DNA synthesis is attempted [E1, K4].

49. Experiments varying the time component either during or following irradiation have revealed the general nature of the defect in ataxia-telangiectasia cells. It was shown that normal cells held after irradiation in a non-growing state had

some recovery (or sparing) from lethal effects, while ataxia-telangiectasia cell lines showed little or no sparing [C9, U15, W2]. More strikingly, irradiation at low dose rates, where the same dose was given over a period of days instead of minutes (factor of 500 difference in dose rates) showed a very large sparing effect on normal cells and little or no effect on ataxia-telangiectasia cells [C10]. These observations are consistent with an inability of ataxia-telangiectasia cells to recover from radiation damage, and they also show that the defect cannot be abrogated simply by allowing more time for damage restitution.

50. Lack of a sparing effect appears to be typical of a particular class of radiosensitive cell lines. Thus lines that are known to have a defect in the repair of DNA double-strand breaks, for example the *xrs* series and XR-1, also lack recovery under irradiation conditions in which their normal counterparts show a large sparing effect [S14, T7]. Similarly, yeast radiosensitive lines that are unable to rejoin double-strand breaks, because of a defect in recombination (*rad50*, *rad51*, *rad52*), also lack sparing [R2, R3]. A substantial body of data in yeast supports the contention that the double-strand break is the DNA damage most likely to be lethal to cells, and that its repair is responsible for the recovery seen under sparing conditions [F3].

51. In contrast to these cellular studies implicating strand breakage as the type of DNA damage involved in the ataxia-telangiectasia defect, it has been difficult to prove that ataxia-telangiectasia cells have a break-repair defect at the molecular

level [M5]. Recently, irradiation at 37°C with low-dose-rate gamma rays has shown a small increase in DNA double-strand breaks following repair ("residual damage") in ataxia-telangiectasia cells relative to normal cells [B9, F10, F16]. However, cytogenetic studies have provided more satisfactory evidence for a break-repair defect: a significantly elevated fraction of unrestituted chromosomal breaks remain in ataxia-telangiectasia cells after irradiation [C11, T1]. Support for these findings has been obtained from the measurement of both DNA double-strand breaks and chromosomal breaks (using prematurely condensed chromosomes to allow rapid analysis) after gamma irradiation of normal and ataxia-telangiectasia lymphoblastoid cells at different phases of the cell cycle [P5]. A consistent decrease in the rapid component of repair was found in ataxia-telangiectasia relative to normal cells; this decrease was small and usually statistically non-significant for DNA double-strand breaks, but larger and significant for chromosomal breaks. Differences in amounts of residual chromosomal damage between normal and ataxia-telangiectasia cells give a close, but not exact, approximation to their relative survival levels after irradiation [C14, P5].

52. Hamster cell lines showing strong similarities to ataxia-telangiectasia have been isolated and their responses characterized (the *irs2* line [J5] and the V-series [Z5]). These lines are hypersensitive to agents known to cause DNA breakage, have radioresistant DNA synthesis, and have no measurable biochemical defect in DNA break repair [T8, Z6]. It has also been shown that the radiation sparing effect is absent in *irs2*, while it is present in other lines that have similar radiosensitivity but do not show ataxia-telangiectasia-like characteristics (such as *irs1* and *irs3*) [T30].

53. Overall, these studies strongly support the view that the increased sensitivity of ataxia-telangiectasia and related cell lines to agents such as ionizing radiation derives from an inability to recover from DNA breakage, leading to a higher level of residual chromosomal damage. However, the molecular mechanisms leading to radiosensitivity in this disorder are still not fully understood, despite considerable recent progress in defining the function of the ataxia-telangiectasia gene product (see Section II.A.3).

54. The functional defects in other cancer-prone disorders have also not been well characterized. Primary cells from Fanconi's anaemia patients have spontaneous delay and arrest in the G₂ phase of growth, as well as an increased frequency of chromosomal aberrations [J15] and recombination [T39]. G₂ delay and aberration frequency increases are corrected by lowering the oxygen tension during growth, leading to the suggestion that reduced detoxification of oxygen radicals may be responsible for the phenotype [C62, J17]. However, immortalized Fanconi's anaemia fibroblasts have lost this oxygen effect, showing that this factor is not a basic (or underlying) defect [S48]. In Bloom's syndrome, the high frequency of sister chromatid exchanges and specific types of chromosome aberrations suggested a defect in DNA repair and/or DNA replication.

3. Analysis of genes determining radiosensitivity

55. The classification of radiosensitive disorders and cell lines into genetic groups, followed by mapping and cloning of the affected genes, has dramatically increased knowledge of the molecular mechanisms of recovery from DNA damage. Once the affected gene has been cloned, its sequence may reveal the nature of the gene product (protein), because of similarities to known genes. Gene sequence data from at-risk groups will also allow deleterious mutations to be identified, and permit analysis of the role of these genes in disorders such as cancer. Manipulation and expression of the gene under defined experimental conditions allow specific functions to be studied in cells and in animals. Animal models of the human disorder can be created by replacing the normal pair of genes with defective copies (a knockout animal; see paragraph 46) and assessing the resulting phenotype.

56. Genetic classification of the ataxia-telangiectasia disorder initially indicated that several different genetic groups might exist [C3, J2], but mapping and cloning of a gene (*ATM*) found to be mutated in patients has cast doubt on this designation. The genetic mapping data, based on ataxia-telangiectasia family studies, mostly placed the affected gene into one chromosomal region, 11q23.1 [G2, Z2]. Positional cloning procedures in this region led to the identification of the *ATM* gene, which has homology to a gene family encoding PI-3 kinases [B31, S24, S25]. The PI-3 kinase family contains a number of large proteins involved in cell-cycle checkpoints, the regulation of chromosome-end length, and DNA break repair, including site-specific recombination (see Section II.B.1). It is therefore likely that *ATM* and other members of this family are involved in the detection of certain types of DNA alterations and may coordinate response by signalling these changes to other regulatory molecules in the cell [J8, K3, S25, T29].

57. Analysis of mutations in the *ATM* gene of ataxia-telangiectasia patients showed that the majority are compound heterozygotes (i.e. the mutations in the two gene copies derive from independent events) and that these commonly lead to an inactive, truncated protein [G19, M30]. However, individuals from 10 families in the United Kingdom with less severe symptoms (paragraph 42) all have the same mutation in one copy of *ATM* (a 137-bp insertion caused by a point mutation in a splice site) but differ in the mutation in the other gene copy. The less severe phenotype appears to arise from a low level of production of normal protein from the insertion-containing gene copy. Two more families with this less severe phenotype have mutations leading to the production of an altered but full-length protein, again suggesting that the severity of the symptoms in ataxia-telangiectasia is linked to the genotype of the individual [L41, M30]. However, it is possible that individuals with less severe symptoms have an increased risk of developing specific types of cancer [S78] (see also paragraph 139).

58. A further two families presenting with many of the symptoms of ataxia-telangiectasia (paragraph 39) but

without dilated ocular blood vessels failed to show mutations in the *ATM* gene. The cellular characteristics of family members, such as radiation sensitivity and DNA-synthesis delay (paragraph 40), were mostly intermediate between those of classical ataxia-telangiectasia patients and normal individuals. Examination of other genes implicated in the repair of DNA double-strand breaks revealed that the human homologue of the yeast *MRE11* gene (paragraph 74) was mutated in these families [S82].

59. A rare form of leukaemia, sporadic T-cell prolymphocytic leukaemia (T-PLL), shows a high frequency of *ATM* mutations; although some of the changes identified in the *ATM* gene were rearrangements, most mutations in T-PLL were single DNA base-pair changes in the kinase region of the *ATM* gene and did not lead to protein truncation [V5, Y8]. This finding has prompted the suggestion that the *ATM* gene acts as a tumour suppressor in cells that may develop T-PLL. There is no evidence for an involvement of the *ATM* gene in T-cell lymphoblastic leukaemias [L58], but in B-cell chronic lymphocytic leukaemia (B-CLL), 34%-40% of tumour samples showed about 50% reduction in levels of ATM protein [S70, S76]. *ATM* mutations were detected in the tumours of 6 out of 32 patients (18%); also 2 of these 6 patients had mutations in both tumour and normal cell DNA, indicating that they were carriers of an *ATM* gene defect (i.e. they inherited a mutation in one copy of the *ATM* gene; see paragraph 137 *et seq.*) [S76]. Similar results were obtained in a separate study [B62], and although they are based at present on small samples, the data suggest that the frequency of *ATM* mutations in the normal cells of patients developing B-CLL may be much higher than in the general population [B62, S76]. Patients with ATM deficiency also had significantly shorter survival times [S70].

60. Study of the ATM protein has shown that it has a nuclear location and is expressed in many different human tissues. Additionally, ATM protein does not increase in amount in response to cell irradiation, consistent with the idea that it is part of a DNA-damage-detection system rather than being regulated in response to DNA damage [L41]. However, the ATM protein does associate with DNA, and this interaction increases when the DNA is irradiated [S77]. The *ATM* gene has a complex structure with multiple transcription start sites, leading to messenger RNAs of different length and predicted secondary structure. This multiplicity of *ATM* transcripts may allow cells to modulate ATM protein levels in response to alterations in environmental signals or cellular metabolism [S40].

61. Initially it was also suggested that the Nijmegen breakage syndrome involves more than one genetic group [J1], but recent evidence shows that only one gene is involved [M41] and that this gene maps to chromosome 8q21 [M41, S59]. These data support clinical findings (paragraph 41) showing that the Nijmegen breakage syndrome is a separate radiosensitive disorder, distinct from ataxia-telangiectasia. The cloning of the gene (*NBS1*) mutated in this syndrome has confirmed this (paragraph 74), with the majority of patients carrying small deletions in this gene [M45, V6].

62. Knockout mice that lack a functional homologue of the ataxia-telangiectasia gene (*Atm*) have recently been bred; they show many of the symptoms of the human disorder, but also give further insights into the action of this gene [B33, E9, X3]. For example, the *Atm* knockout mice are viable but growth-retarded and infertile. They are also very sensitive to acute gamma radiation; at a whole-body dose of 4 Gy, about two thirds of the *Atm* knockout mice died after 5-7 days, while normal and heterozygous mice remained without morbidity after two months [B33]. Primary cells derived from the *Atm* knockout mice also show many of the characteristic features of cells from ataxia-telangiectasia patients [X4]. The cells grow poorly, are hypersensitive to gamma radiation, and fail to undergo arrest of the cell cycle following irradiation (see also paragraph 102). The *Atm* gene product was found to locate to homologous chromosomes as they associate (synapse) at meiosis in germ cells [K27]; loss of fertility in the knockout mice results from failure of meiosis due to abnormal synapsis and subsequent chromosome fragmentation [X3]. Immune defects occur in these mice, and the majority develop thymic lymphomas and die before four months. Based on the understanding to date of the defective response to DNA breakage in ataxia-telangiectasia, the meiotic failure and immune defects in these mice could both relate to an inability to respond to "programmed" DNA double-strand breaks (site-specific breaks that occur in the course of normal cellular processes). Such breaks are thought to be essential in meiosis, to drive the process of meiotic recombination, and they are required for V(D)J recombination in immune system development (see paragraph 82).

63. Unlike in the human disorder, no evidence was found of gross cerebellar degeneration in *Atm* knockout mice aged 1-4 months, but it is possible that the animals are dying too early for this symptom to be revealed [B33, X4]. However, behavioural tests indicated some impairment of cerebellar function [B33], and detailed studies of the brain in these animals showed that *Atm* deficiency can severely affect dopaminergic neurons in the central nervous system [E14]. There is an almost complete absence of radiation-induced apoptotic cell death (Section II.B.3) in the developing central nervous system of *Atm*-deficient mice, while the thymus shows normal levels of apoptosis after irradiation [H48]. Additionally, elevated levels of oxidative damage were recorded in mouse tissues affected by the loss of ATM, especially the cerebellum [B63].

64. In yeast cells, the closest sequence homologue to the *ATM* gene product is the Tel1 protein; mutations in the *TEL1* gene are associated with shortened chromosome ends (telomeres) and genetic instability (three- to fourfold increased levels of mitotic recombination and chromosome loss) [G22]. It has been known for many years that chromosomes in ataxia-telangiectasia cells show a relatively high incidence of telomere fusions, and recent studies have shown that preleukaemic cells from ataxia-telangiectasia patients have an increased rate of telomere loss [M29]. This loss may contribute to chromosomal instability [R40]. However, as yet the relationship between the telomere fusions and tumorigenesis in ataxia-telangiectasia is not clear.

Interestingly, loss of the *TEL1* gene does not lead to x- or gamma-ray sensitivity [G22, M31], but the combined loss of *TEL1* and another gene in the same family (*ESR1/MEC1*; see next paragraph) gives extreme sensitivity to gamma rays, suggesting that these two genes may functionally overlap in protecting the cell against radiation damage [M31]. In fission yeast, following prolonged growth, loss of these two genes led to circular chromosomes lacking telomeric sequences [N26].

65. A further member of the human PI-3 kinase family, related to *ATM*, has been found through homology searches [B32, C38]. The gene (named *ATR* or *FRP1*) encodes a protein that is most closely similar to the fission yeast *rad3* gene product, which is itself structurally and functionally related to the budding yeast *ESR1/MEC1* and the fruit fly *mei-41* gene products. Mutations in these genes render the yeast or flies sensitive to killing by both ionizing and UV radiation, and are known to play important roles in mitotic and meiotic cell-cycle controls. Expression of an inactive form of *ATR* at a high level in human fibroblasts increased radiation sensitivity by a factor of 2-3 and abrogated radiation-induced G₂ arrest (see paragraph 108). Additionally, cells with defective *ATR* develop abnormal nuclear morphologies, which may indicate further cell-cycle perturbations [C48, W50]. The *ATR* gene product is expressed at a high level in germinal tissue and localizes along unsynapsed meiotic chromosomes [K27], thereby playing a role that is complementary to that of the *ATM* gene product (paragraph 62). Thus in addition to their roles in mitotic cells, *ATM* and *ATR* gene products may be involved in the co-ordination of meiotic chromosome synapsis, perhaps by signalling breaks and monitoring repair synthesis to guard against genetic instability.

66. Some other genes affecting radiosensitivity in humans have also been mapped and cloned. Following the cloning of the human DNA ligase I gene, it was discovered that this was defective in a unique individual with growth retardation, immunological abnormalities, and cellular sensitivity to a variety of DNA-damaging agents, including radiation [B5]. The genes for other human DNA ligases have also been cloned using homology searches or protein purification [C31, W19], and subsequently another unique radiosensitive individual with impairment of double-strand break rejoining (described in paragraph 43) was found to be defective in DNA ligase IV [R38]. It has been established in yeast and mammalian cells that DNA ligase IV is specifically involved in the repair of DNA breaks by non-homologous end joining (Section II.B.1) [S64, T40, W46]. DNA ligase III has been implicated in the base-excision repair pathway (see paragraph 78).

67. Bloom's syndrome patients of different ethnic origins (Ashkenasi Jewish, French-Canadian, Mennonite, and Japanese) were found by cell fusion analysis to fall into one genetic group [W36]. The gene defective in Bloom's syndrome (named *BLM*) has been positionally cloned and shown to be similar to the bacterial RecQ helicase, a type of enzyme that opens up the DNA helix and is associated with genetic recombination in bacteria [E6, K39].

Knocking out the gene homologous to *BLM* in mice results in embryonic growth retardation and lethality, apparently because of a wave of increased apoptotic death [C61]. The yeast homologue of the *BLM* gene product interacts with DNA topoisomerase enzymes, known to be required for the resolution of interlocking DNA molecules following replication. In the absence of the *BLM* gene product, it is suggested that replicated DNA (in mitosis or meiosis) is entangled and may give rise to sister-chromatid exchange and chromosomal non-disjunction [W35]. Support for this idea comes from studies with the fission yeast, where the recQ helicase is required for the recovery from cell-cycle arrest during DNA replication following DNA damage; absence of the recQ helicase leads to an increase in the rate of genetic recombination [S60].

68. Cells derived from Fanconi's anaemia patients have been classified into eight genetic groups [J16, J19, S46], consistent with the heterogeneity of symptoms found (paragraph 45). To date, three of these genes have been mapped, *FANCA* to chromosome 16q24 [G21, P20], *FANCC* to 9q22 [S46], and *FANCD* to 3p22-26 [W37]. The *FANCA* and *FANCC* genes have now been cloned [F9, L43, S47]; these gene sequences predict proteins that are structurally different from each other and from other known proteins. Mutations of the *FANCA* gene were analysed in 97 patients from different ethnic groups with Fanconi's anaemia; the majority of mutations detected were either DNA base-pair alterations or small deletions and insertions scattered throughout the gene, with a smaller number of large deletions [L49]. The *FANCC* gene product is a cytoplasmic protein [Y7], but it interacts with the *FANCA* gene product, and the complex translocates to the nucleus [K40]. While this nuclear localization is consistent with a possible role in DNA repair, the precise function of these gene products remains unclear. *FANCC* also has binding sites for the p53 tumour suppressor protein (paragraph 100), and binding of p53 to the gene can regulate its expression [L46].

69. One of the more recently identified genetic groups of Fanconi's anaemia, FA-G, has been found to be caused by mutation of a previously-identified gene, *XRCC9* [D18]. This gene was identified by the complementation of a rodent cell line sensitive to a variety of DNA-damaging agents including a twofold enhanced sensitivity to x rays, and is potentially involved in post-replication repair of DNA [L59] (Table 2).

70. Human genes implicated in the repair of radiation-induced DNA damage have also been cloned by their structural homology to genes involved in specific repair pathways in lower organisms. For example, it is known that the main pathway for the repair of DNA double-strand breaks in lower organisms involves homologous recombination (paragraph 30); in budding yeast, the genes responsible are *RAD50*, *RAD51*, *RAD52*, *RAD53*, *RAD54*, *RAD55*, *RAD57*, *MRE11*, and *XRS2* (often called the *RAD52* group of genes). Mutations in these genes render the yeast cells defective in mitotic and/or meiotic

recombination and sensitive to ionizing radiation. At least some of the products of these genes act together in a multi-protein complex to effect recombination [H27, J14, U18]. In yeast, functional relationships have also been found between proteins of the PI-3 kinase family and some of the Rad52 group proteins. For example, the activity of the *RAD53* gene product, another protein kinase, is regulated by both *ESR1/MEC1* and *TEL1* gene products, suggesting that the Rad53 protein acts as a signal transducer in the DNA damage response pathway downstream of these ATM-like proteins [S43]. Sequence similarities have led to the cloning of human homologues of some of the yeast *RAD52* gene group: *hRAD50* [D16], *hRAD51* [S10, Y6], *hRAD52* [M10, S44], *hRAD54* [K28], and *hMRE11* [P22]. Preliminary evidence has been obtained from their abilities to partially correct the defects in radiosensitive yeast cell lines that some of these genes, or their mouse homologues, are involved in the repair of radiation damage [K28, M37]. Further, knocking out the *RAD54* gene in both the fruit fly [K41] and mouse [E12] confers radiation sensitivity and a defect in homologous recombination. There is also evidence for regulated expression of these recombination proteins through the cell cycle, with levels of RAD51 and RAD52 increasing during the S phase and peaking in G₂ in human and rodent cells [C5].

71. The human *RAD51* gene product has been shown to bind DNA and thereby underwind the double helix, an early step in the recombination process [B34]. The *RAD51* gene in yeast is not essential for cell survival, but recently the homologous gene has been knocked out in mice and surprisingly was found to give embryonic lethality [L42, T34]. Knockout embryos arrested early in development but progressed further if there was also a mutation in the gene encoding the tumour suppressor p53 (*TP53*), possibly because of a reduction in programmed cell death (see paragraph 112). This may suggest that *RAD51* in mammals has functions in addition to recombination and repair, and in a separate study the human RAD51 protein has been shown to directly interact with p53 [S45]. Cells from the mouse *Rad51* knockout were hypersensitive to gamma rays, although this response has not been quantified, and show chromosome loss in mitotic cells [L42]. In mitotic cells, the mouse RAD51 protein is found to concentrate in multiple foci within the nucleus at the DNA-synthesis (S) phase of the cell cycle [T37]. In meiotic cells, RAD51 is found on synapsing chromosomes and disappears shortly after this stage [P23], but this pattern of localization of RAD51 is disrupted in *Atm*-deficient mice [B47].

72. The pattern of cellular localization found for the RAD51 protein has also been found for the product of a gene commonly mutated in familial breast and ovarian cancer patients, *BRCA1*; subsequently, RAD51 and BRCA1 proteins were shown to interact directly [S51]. *Brcal*-deficient mouse cells show a modest increase in sensitivity to ionizing radiation and have a reduced capacity to repair base damage in transcribing DNA (transcription-coupled repair, paragraph 28) [G34]. In human *BRCA1*-deficient cells, increased sensitivity to cell killing by gamma rays and reduced repair of DNA double-

strand breaks can be partially corrected by introduction of the normal gene, while mutant *BRCA1* genes fail to restore these defects [S86]. A product of a second gene associated with familial breast cancer susceptibility, *BRCA2*, has similarly been shown to interact with RAD51 [M42, S61], as well as with p53 [M46]. *BRCA1* and *BRCA2* are highly expressed in rapidly proliferating cells, with expression highest at the start of S phase of the cell cycle [R23]. Analysis of mutations in the *BRCA* genes has shown that families with a high proportion of breast cancer tend to have mutations in different parts of the gene from families having a predisposition to ovarian cancer [G37, G38]. Despite their importance in cancer predisposition, the molecular function of the *BRCA* genes is unknown, and the finding of interaction with RAD51 provided the first clue that their role may be in DNA repair. As noted in paragraph 62, the ATM protein also has a specific pattern of localization to meiotic chromosomes, and loss of ATM can disrupt the localization pattern of RAD51. Thus, through their connections with *RAD51*, these data provide for the first time a mechanistic link between four different genes involved in cancer susceptibility (*ATM*, *TP53*, *BRCA1*, and *BRCA2*).

73. Mice have been bred with a knockout of the *Brcal* gene or the *Brcal2* gene; like *Rad51* knockout mice, they were both found to arrest early in embryonic development [H40, L50, S61]. However, viable mice have been bred with a mutation in *Brcal* or *Brcal2* that does not completely inactivate the gene. A conditional deletion of the *Brcal* gene in mice, confining the defect to the mammary glands, gave a low frequency of mammary tumour formation with long latency (see also paragraph 106) [X7]. Embryonic fibroblasts derived from *Brcal*-deficient mice proliferate poorly and show genetic instability [X8] and a reduction in the frequency of homologous recombination [M50]. About one third of the *Brcal2*-deficient mice survived to adulthood but showed small size, poor tissue differentiation, absence of germ cells, and development of thymic lymphomas [C59]. Cells derived from viable *Brcal2*-defective mice proliferate poorly, apparently as a result of spontaneously high levels of p53 and p21, causing cell-cycle arrest (see paragraph 100) [C59, P32]. These cells accumulate high levels of chromosomal aberrations and show enhanced sensitivity to DNA-damaging agents, including x rays [M48, P32]. The evidence linking the *BRCA* genes to DNA repair processes suggests that they may not function as tumour suppressors but are involved in the maintenance of genome integrity.

74. The yeast *RAD50*, *MRE11*, and *XRS2* gene products assemble into a multi-protein complex that is implicated in the processing of broken DNA, as well as in a number of other functions, including meiotic recombination and telomere maintenance [H46]. This multiplicity of roles for the complex may arise from its nuclease activities on both single- and double-stranded DNA ends that trim up the DNA in preparation for end-joining and homologous recombination processes. Human gene products with similar functions have been identified (paragraph 70); significantly, the gene mutated in the Nijmegen breakage syndrome (*NBS1*) has been found

to be the functional homologue of *XRS2* (paragraphs 41 and 61) [C54, M45, V6], and individuals with a variant form of ataxia-telangiectasia have mutations in the human homologue of *MRE11* (paragraphs 42 and 58) [S82]. The *NBS1* gene product contains motifs that are commonly present in proteins involved in cell-cycle regulation and DNA-damage response, suggesting that like ATM, this gene has a role in the signalling mechanism following damage to DNA by ionizing radiation. Study of the purified protein shows that DNA unwinding and nuclease activities of RAD50/MRE11 are promoted by the presence of NBS1 [P34]. The human RAD50 and MRE11 proteins co-localize in cell nuclei following ionizing radiation damage (but not after UV irradiation); their sites of localization were distinct from those of the RAD51 protein, consistent with their different roles in DNA repair [D17]. Co-localization of MRE11/RAD50 proteins was much reduced in ataxia-telangiectasia cells, suggesting that *ATM* gene signalling is important for the assembly of the break-rejoin complex [M28]. Additionally, the BRCA1 protein has been shown to colocalize and interact with RAD50, and radiation-induced sites of localization including MRE11, RAD50 and NBS1 proteins were much reduced in breast-cancer cells lacking BRCA1 [Z12].

75. Human genes encoding enzymes responsible for the repair of the numerous forms of base damage caused by DNA-damaging agents, including ionizing radiation, are also being discovered [L69, S49]. A number of key enzymes (DNA glycosylases, endonucleases) in the base-excision repair pathway have been isolated by biochemical purification, followed by protein sequencing and gene identification, or through their homologies to known enzymes in lower organisms. For example, the hOGG1 glycosylase [R41] removes 8-oxoguanine opposite a cytosine (paragraph 24), while the hNTH1 glycosylase [A28] removes oxidized pyrimidines such as thymine glycol (paragraph 17). The pathway of base-excision repair that results in the incorporation of a single nucleotide ("short-patch repair") has been reconstituted under cell-free conditions, using purified enzymes [K24]. Additionally, a second "long-patch repair" pathway has been identified, in which between two and six nucleotides are replaced following repair of a reduced or oxidized baseless site; this pathway also requires the structure-specific nuclease DNase IV (also known as FEN1) to remove the displaced nucleotides during repair [K42]. The reconstitution of base-excision repair of oxidized pyrimidines in DNA has revealed that hNTH1 glycosylase is strongly stimulated by one of the proteins involved in nucleotide-excision repair, XPG (xeroderma pigmentosum group G protein). The XPG protein binds to non-paired regions of DNA, and acts as a structure-specific nuclease where the unpaired region is greater than five base pairs, as will happen at sites of bulky damage. However, at sites of oxidative base damage it seems that the unpaired region is less than five base pairs, and instead of cutting DNA the XPG protein promotes the activity of the hNTH1 in removing the base damage [K46]. These findings may explain the extreme symptoms of some patients lacking XPG, including growth failure and neurological

dysfunction, and the early death of mice carrying a knockout of the *Xpg* gene [H51] (while mice that are totally defective for nucleotide-excision repair such as those defective in *Xpa* are viable [N27]).

76. The main endonuclease involved in the repair of baseless sites (Figure 1a) in human cells, known variously as HAP1, APE, APEX, or Ref-1, was cloned in several laboratories following biochemical purification and was found to have a surprising additional function when compared with the bacterial enzyme. It stimulates the DNA-binding activity of transcription factors involved in signal transduction such as Fos, Jun, NFκB, and p53 by reduction/oxidation mechanisms [X5]. Signalling and repair responses to oxidative damage may therefore be coordinated through this one enzyme; this may be especially important during tissue proliferation, since *Ref-1* knockout mice die during embryonic development [X6]. Knockout mice have been produced for several other genes in the base-excision pathways, and a number of these have also proved to be embryonic-lethal, showing their importance for the normal functioning of cells and tissues [W45]. However, a knockout mouse for the broad-specificity glycosylase 3-methyladenine-DNA glycosylase II was recently shown to be viable [E13], possibly because of its mode of action (paragraph 23) and the likelihood that other more specialized glycosylases can substitute for it in removing damaged bases.

77. The abundant enzyme poly(ADP-ribose) polymerase, also known as PARP, is rapidly recruited to sites of DNA breakage following irradiation, where PARP transiently synthesizes long, branched chains of poly(ADP-ribose) on itself and other cellular proteins. These chains are degraded by another enzyme, poly(ADP-ribose) glycohydrolase, with a half-life of only a few minutes. The role of PARP in response to DNA damage, which has long been in dispute [C49], has been clarified from recent experiments in which the *Parp* gene was knocked out in mice [M9]. These mice are viable and fertile, although adult size is smaller than average and litter sizes are smaller than for normal mice. Following whole-body irradiation (8 Gy gamma rays), the *Parp* knockout mice died much more rapidly than normal mice from acute radiation toxicity to the small intestine; the survival half-time of these irradiated PARP-deficient mice was comparable to that of irradiated *Atm* knockout mice (paragraph 62). PARP-deficient cells or cells in which a mutated PARP is expressed [S63], show increases in chromosomal aberrations, sister-chromatid exchanges, and apoptosis following DNA damage. The average length of telomeres is significantly shorter in PARP-deficient mouse cells, leading to an increased frequency of chromosomal fusions and other aberrations [D21]. Recent biochemical data suggest that PARP has an important function as a molecular sensor of DNA breaks, especially single-strand breaks, and its absence reduces the efficiency of base-excision repair [O4]. In addition, it has been suggested that the synthesis of poly(ADP-ribose) chains causes negative-charge repulsion of damaged DNA strands, preventing accidental recombination between homologous sequences [S62].

78. To date, studies with radiosensitive mammalian lines have identified more than eight genetic groups [J3, T9, Z13]. The genes responsible have provisionally been named the *XRCC* (for *X-Ray Cross-Complementing*) group, despite the fact that some of the cell lines are not primarily sensitive to x rays. The human gene corresponding to the first group, *XRCC1*, has been cloned and encodes a protein interacting with DNA ligase III [C2]. Further, the *XRCC1* protein has been shown to associate with DNA polymerase β [C39, K29], which fills in the gaps created during the repair of damaged bases. It is suggested that *XRCC1* may act as a scaffold protein in the final steps of base-excision repair (Figure Ia), supporting the activity of DNA polymerase β and DNA ligase III [K29]. The *XRCC2* gene [C66, L65, T35] and the *XRCC3* gene [L65] have been cloned recently, and both genes have structural homology to the yeast and human *RAD51* genes [T47]; it is likely, therefore that they are involved in repair of damage by homologous recombination. Other human genes recently cloned using mammalian cell lines (*XRCC4*, *XRCC5*, *XRCC7*) are involved in the repair of radiation-induced DNA double-strand breaks as well as in immune gene recombination, as detailed below (Section II.B.1). Current mapping data place all of the remaining genes on different chromosomes, and none maps to the location of characterized radiosensitive human syndromes (Table 2).

4. Summary

79. Several radiosensitive human disorders and examples of individual radiosensitivity have been identified in recent years. The sensitivity in these individuals is characterized by a greatly increased susceptibility to cancer, although this cancer is not necessarily radiation-induced. Many radiosensitive

lines of cultured cells have also been established, and these have been useful for identifying the genes involved and for functional studies. In general, it has been found that enhanced sensitivity arises from an inability to recover from DNA damage, because of a reduction in damage detection and/or repair processes. Reduction of repair capacity commonly leads to a lack of low-dose-rate sparing and a higher level of genetic changes.

80. Considerable progress has recently been made in defining repair gene functions in human and other mammalian cells, and a summary of well-defined repair pathways for damage by ionizing radiation is shown in Figure III. A number of important genes have been cloned, including the gene *ATM*, which determines sensitivity in the ataxia-telangiectasia disorder. Studies with the *ATM* gene product and the production of mice defective for the gene suggest that it participates in the detection of DNA double-strand breaks and passes this information on to other important molecules regulating cellular response processes. Other recently discovered genes act directly in the repair of radiation damage; for example, the *RAD51* gene is vital for repair by homologous recombination. The *RAD51* gene product has also been found to interact with products of the breast-cancer susceptibility genes, *BRCA1* and *BRCA2*, suggesting an unsuspected role for these genes in damage recovery. In accordance with this finding, mice defective for the *BRCA* genes have symptoms very similar to *Rad51*-defective mice, and the *Brca2*-defective mice are radiation-sensitive. Additionally, some repair genes when knocked out in animals give embryonic lethality, showing that these genes have important roles in basic cellular processes influencing tissue development.




Figure III. Important genes in DNA repair pathways in human cells following damage by ionizing radiation.

Some of the important genes identified in each pathway are shown (the *h* prefix indicates a human homologue of a yeast gene); names of proteins are shown in brackets if different from the gene names. The *BRCA* genes and the *PARP* gene (square brackets) have speculative roles in recombination repair and base-excision repair, respectively. Pathways that can lead to mutation are indicated.

81. The conservation of repair genes from lower to higher organisms and progress in understanding the human genome suggest that further rapid progress will be made in the discovery and analysis of genes influencing radiation sensitivity. These genes will be important tools in understanding the extent of variation in radiation sensitivity in the human population and should help to identify individuals especially at risk.

B. RELATIONSHIP BETWEEN REPAIR AND OTHER CELL REGULATORY PROCESSES

1. Radiosensitivity and defective recombination in the immune system: non-homologous end joining of DNA double-strand breaks

82. It has been noted that radiosensitive disorders are often associated with some degree of immune deficiency. The recent discovery that the *scid* (severe combined immune-deficient) mouse strain is radiosensitive has led to a reexamination of this relationship and to rapid progress in understanding at least one mechanism of DNA-break rejoining. In the immune system, functional immunoglobulin and T-cell receptor genes are assembled from separate genetic regions during lymphoid differentiation [A5]. The separate regions (termed "V" for variable; "D" for diversity, and "J" for joining) involved in this type of site-specific recombination are flanked by recognition (signal) sequences at which double-strand DNA breakage occurs prior to the rejoining of the regions. Mice homozygous for the *scid* mutation lack functional T-cells and B-cells because of a defect in V(D)J recombination [H8, L9], and about 15% of the mice develop lymphomas [C50]. It was shown that fibroblasts derived from homozygous *scid* mice are three to four times as sensitive as normal mice to x rays or gamma rays and have a reduced ability to repair DNA double-strand breaks [B8, H10]. Mice heterozygous for the *scid* mutation also show some increased sensitivity to gamma rays, compared to normal mice, and cultured bone marrow cells from *scid* heterozygotes are marginally more radiosensitive than normal cells [K45]. Detailed studies of repair capacity in *scid* cells show that the rate of repair of double-strand breaks is reduced by a factor of about 5, but if sufficient time is allowed (24 h) the final levels of repair are similar to those of normal cells [N20]. Subsequently, similar tests on several radiosensitive mammalian lines derived from cultured cells showed that those with large defects in double-strand break repair are also defective in V(D)J recombination, while radiosensitive lines, including ataxia-telangiectasia cells, with near-normal double-strand break repair are not V(D)J-recombination-deficient (Table 2) [H12, K25, P6, T4, T11].

83. At least four different gene products are common to V(D)J recombination and repair of radiation-induced DNA breaks; three of these products are defined by their respective rodent sensitive lines: *xrs*, *scid/V3*, and *XR-1/M10* (Table 2). Recently there has been a breakthrough in identifying the genes and gene products complementing

the defects in these lines, to define a repair pathway termed non-homologous end joining (paragraph 31). The *xrs*-complementing gene (also known as *XRCC5*) codes for a subunit of the Ku antigen (p86), a DNA end-binding protein discovered in normal human cells that reacted with sera from patients with certain autoimmune diseases [T5]. The *scid/V3* protein (*XRCC7* gene product) has been found to be the catalytic subunit of a large DNA-dependent protein kinase, also known as p460, belonging to the PI-3 kinase family and related to the ataxia-telangiectasia gene product (see paragraph 56) [H17]. The Ku antigen consists of two subunits, p70 and p86, which interact to form a dimer; this dimer binds to broken ends of DNA, recruiting the p460 subunit and conferring kinase activity on the complex [G7]. The DNA-dependent protein kinase has been shown to phosphorylate a number of substrates *in vitro*, including the p53 tumour suppressor and RNA polymerase II [A4], but it is not clear whether these constitute important targets *in vivo*. The most recently cloned gene of this series, the human *XRCC4* gene complementing the *XR-1* line, encodes a protein unrelated to any other yet described [L30]. Functional studies of the *XRCC4* protein show that it is a nuclear phosphoprotein that is a substrate *in vitro* for the DNA-dependent protein kinase and that it associates with the recently discovered DNA ligase IV [C51, G30]. This discovery suggests that *XRCC4* acts as a go-between in the assembly of a DNA-break repair complex in which the final step is mediated DNA ligase IV (paragraph 66). Experiments using cell extracts show that the rejoining of breaks by mammalian DNA ligases is stimulated by purified Ku86 protein, especially when the break ends cohere poorly [R24]. This observation suggests that Ku86 may function to stimulate ligation, perhaps through its ability to bridge the gap between broken ends. However, the precise mode of action of these proteins in promoting the repair of DNA double-strand breaks (and presumably in initiating a coordinated response through kinase action on relevant proteins) is presently under intensive study.

84. The different subunits of the DNA-dependent protein kinase are not induced following x irradiation [J22]. However, the gene encoding Ku86 in primates (but not in rodents) can be expressed in a form that has been found to respond to radiation damage. In this form the gene gives rise to a related protein, termed KARP-1, which includes a p53 binding site [M43], and the RNA transcript and protein were increased by a factor of up to 6 at 90 min following x-ray irradiation. Interestingly, this induction did not occur in cells defective in either p53 or ATM, suggesting that at least some of the non-homologous end joining is activated through interactions with p53 and ATM signalling processes [M44].

85. Sequencing the entire gene encoding the p460 kinase from *scid* cells showed that a single DNA base alteration leads to a premature stop codon in the highly conserved terminal region of the gene [A20], as other data had predicted [B37, D11]. This mutation gives a protein truncated by 83 amino acids in *scid* cells, leading to a partial abolition of kinase activity. Mutations in this gene have subsequently been

found in other radiation-sensitive lines of cultured cells; for example, the SX9 line (Table 2) has a mutation giving rise to a single amino-acid substitution at a position before that of the *scid* mutation, and leads to a more severe phenotype than the *scid* mutation [F25]. The partial loss of activity in *scid* [A19, B38, C52, P26] may explain another surprising phenomenon, whereby sublethal irradiation of *scid* mice can transiently rescue V(D)J recombination [D12]. While the mechanism of this rescue is still unknown, it suggests that p460 in *scid* is not completely inactive.

86. The importance of the catalytic subunit of DNA-dependent protein kinase in the development of T-cells and B-cells and in protecting animals against cancer has been illustrated by the chance finding of transgenic mice in which the *Xrcc7* gene was knocked out [J20]. These mice, named *slip*, showed a lack of mature lymphocytes, but most remarkably all animals died within 5-6 months from thymic lymphoblastic lymphomas. No tumours of B-cell or myeloid linkage were found, suggesting that *Xrcc7* acts as a tumour suppressor of the T-cell lineage. Treatment of *scid* mice with gamma rays (1 Gy) at 24-48 h after birth led to 86% developing T-cell lymphoma with very short latency, with no other tumour type observed [G35]. Further, mice having both the *scid* defect and a knockout of the *Parp* gene (paragraph 77) also develop very high levels of T-cell lymphoma, suggesting that the carcinogenic effects of a partial defect in double-strand break rejoining is exacerbated by a reduction in repair capacity for other types of DNA damage [M6].

87. The assessment of radiation sensitivity of different tissues has shown that epithelial cells of the intestine and kidney in *scid* mice show a much greater radiosensitization than bone marrow cells from the same animals, compared to the relative sensitivity of these cells in normal animals [H28]. However, the additional sensitivity of *scid* epithelial cells effectively made these cells similar in overall sensitivity to *scid* marrow cells (e.g. at a dose of 2 Gy), since normal epithelial cells are more radioresistant than normal marrow cells. Irradiation of epithelial cells at low dose rate (16 mGy min⁻¹) altered the survival of *scid* cells much less than it did for normal cells [H28], as expected for this type of radiosensitive cell line (see paragraphs 49-50).

88. In addition to the *scid* mouse as a model of defective DNA double-strand break repair, knockout mice for the gene encoding Ku86 have been created [N21, Z9]. These mice weigh 40%-60% less than normal mice, and further weight loss occurred when newborn mice were gamma-irradiated at whole-body doses in excess of 0.25 Gy. Survival of Ku86-deficient mice irradiated at 2-4 months old with gamma rays was also compromised; doses of 3-4 Gy caused 50% mortality in two weeks. Ku86-deficient mice given 2 Gy gamma rays survived for up to 12 weeks but unlike normal mice showed severe hair loss within one month post-irradiation. Examination of tissues at four days post-irradiation showed severe injury to the GI tract at much lower doses than for the normal mice, along with atrophy of lymphoid organs [N21]. In these mice it was found that T- and B-cell development was arrested at an

early stage, as in *scid* mice. The Ku70 subunit has been knocked out in mouse embryonic stem cells, and these cells are sensitive to gamma rays. As might be expected, Ku70-deficient cells are defective in DNA end-binding activity and in V(D)J recombination [G31]. Ku70-deficient mice have similar growth reduction and radiation sensitivity to Ku86-defective mice, but they also develop a high frequency of spontaneous thymic and disseminated T-cell lymphomas [L57].

89. Mice lacking DNA ligase IV do not survive beyond the late embryonic stage of development; ligase IV-deficient embryonic fibroblasts are hypersensitive to gamma rays but not to ultraviolet light and are defective in V(D)J recombination [F20]. Very similar defects have been found in mice and murine fibroblasts lacking the *Xrcc4* gene. Further, lethality in XRCC4- and ligase IV-deficient mouse embryos is associated with severe disruption of the development of the nervous system due to extensive apoptotic cell death [G33]. It is presumed that the enhanced sensitivity of neuronal cells relates to an inability to repair DNA double-strand breaks, although it is not known whether these breaks result from normal metabolism or a specific neuronal process (e.g. a recombination process) required for neuronal function. It is of considerable interest that a single person defective in DNA ligase IV has been identified (paragraphs 43 and 66); this person showed extreme sensitivity to radiation and developed acute lymphoblastic leukaemia at age 14, but was not severely impaired otherwise. Analysis of the ligase IV gene and protein in cells derived from this person showed that ligase function was not completely defective, presumably explaining their relatively normal developmental progress [R38]. This example of repair deficiency is important in revealing that mutations leading to partial activity of the gene product may be permissive for growth and development, but may have undesirable consequences including the possibility of cancer formation.

90. The search for human mutations in the *XRCC4*, 5, 6, and 7 genes among patients known to be compromised in immune functions is beginning to yield some candidates [C28, H26].

91. There is a difference between the "programmed" double-strand breaks generated in the V(D)J recombination process and those caused by radiation damage. The breaks at V(D)J sites have 5'-phosphorylated blunt ends and can be rejoined by a DNA ligase without further processing [S6], while radiation-induced breaks are often not directly ligatable because of extensive damage to the sugars and bases at the break sites (see Section I.B). Since, in the cell, the non-homologous end joining pathway is involved in repairing both types of break, it may be suggested that the exact structure (and possibly complexity) of the breaks does not influence their recognition by the proteins that initiate the rejoining process. However, it seems likely that the context in which the break occurs will influence its repair; in yeast cells, three proteins involved in modulating chromatin structure (Sir2, Sir3, and Sir4) interact with the Ku70 protein and have a role in DNA break rejoining

[T41]. A defect in any of the three *SIR* genes led to increased sensitivity to gamma radiation, providing the homologous recombination pathway was inactive. While the changes in chromatin structure brought about by the Sir proteins are not well understood, it is thought that they may make the broken DNA inaccessible to other DNA-modifying enzymes such as nucleases and thereby protect the damaged DNA from loss. It has also been found that the V(D)J recombination process is restricted to the G₁ phase of the cell cycle [S6]. It is likely that this rejoining process is also cell-cycle-controlled in the repair of radiation-induced double-strand breaks [L51]. Thus, the XR-1 line is highly sensitive and defective in radiation-induced double-strand break repair in G₁ but has nearly normal sensitivity and break repair in late S phase [G5]. One implication of this analysis is that there must be at least two major pathways for the repair of DNA double-strand breaks: the non-homologous end-joining pathway active in G₁ and at least one other pathway operating in other stages of the cell cycle.

92. As noted in paragraph 70, there is evidence suggesting that the homologous recombination repair pathway is active in late S/G₂ stages of the cell cycle; additionally, chick cells defective in the *RAD54* gene were found to be more sensitive than normal cells in this part of the cycle. Chick cell lines were generated with defects in both the homologous recombination repair pathway and the non-homologous end-joining pathway (*RAD54/Ku70*-defective), and these showed increased radiation sensitivity and higher levels of chromosomal aberrations than either single defect [T42].

93. Unlike mammalian cells, in lower eukaryotic organisms such as the budding yeast, repair of DNA double-strand breaks is mainly effected through homologous recombination processes. However, proteins similar to the mammalian Ku70/86 have recently been found in yeast, and these form a dimeric complex binding directly to DNA ends. DNA end-joining in yeast was shown to be impaired when the genes encoding the yeast Ku proteins are mutated, and this process is distinct from the rejoining mediated by homologous recombination. However, it is possible that some activities are shared by both homologous and illegitimate recombination pathways, since it was found that the *RAD50* gene product (paragraph 70) interacts with the yeast Ku proteins [M32]. Mutation of the gene encoding the smaller Ku subunit (*HDF1*) has also been found to lead to shortened telomeres, similarly to *TEL1* mutations (paragraph 64), suggesting that in addition to their end-joining role these proteins may help protect yeast chromosome ends [P21]. Intriguingly, both the Ku and Sir proteins are located at telomeres in undamaged yeast cells, but following the induction of DNA double-strand breaks they relocate to break sites in the genome [M54]. Loss of *HDF1* gene function alone does not, however, lead to radiation sensitivity or a measurable defect in DNA double-strand break repair in yeast, while the combined loss of *HDF1* and homologous-recombination gene functions leads to extreme sensitivity [B35, S50]. Additionally, while the loss of homologous recombination repair in yeast leads to an elevated frequency of chromosomal aberrations, the combined loss of

these two pathways gives a reduction in chromosomal aberration frequency [F24]. This observation has been considered as evidence for the involvement of non-homologous end joining mechanisms in chromosomal aberration formation.

94. While ataxia-telangiectasia cells are not defective in V(D)J recombination, a recurrent feature of the disorder is the appearance of T-cell clones with characteristic chromosomal rearrangements at sites of immunoglobulin and T-cell receptor genes [A7, B2, R4]. Elevated levels of recombination of T-cell receptor genes have been described in lymphocytes from ataxia-telangiectasia patients relative to those from normal individuals [L10]. ATM-deficient mice develop thymic lymphomas and die by 5 months (paragraph 62), but if the gene that normally causes programmed double-strand breaks leading to V(D)J recombination is inactivated, no lymphomas develop [L66]. This striking observation suggests that ATM normally suppresses aberrant recombination events.

95. The rejoining of DNA double-strand breaks, produced enzymatically, has also been examined using small DNA molecules transfected into mammalian cells or exposed to cell extracts. These studies have shown that almost any sort of break end (with flush ends, or with complementary sequence at the ends, or with mismatched ends) can be rejoined by cellular enzymes [N7, P29, R1]. However, even with complementary ends, a fraction of the breaks are rejoined with a loss of sequence around the break sites. This mis-rejoining process was found to occur by a non-conservative recombination mechanism [T10] that appears to differ from the non-homologous end-joining pathway. The mechanism entails deletion of DNA bases between short (2-6 bp) direct repeat sequences, such that one of the repeats is also lost (Figure IV; see also paragraph 182). Occasionally the mis-rejoining can be more complex, with an insertion of DNA at the deletion site. Using substrates that attempt to model more closely DNA double-strand breaks produced by radiation damage, where the sequences at the break-ends are not complementary, breaks were shown to be rejoined by either this repeat-driven mechanism or by a process of blunting the ends before rejoining [M33, R18]. The repeat-driven process of mis-rejoining has been shown to be independent of the Ku proteins [M33]. Recent data with yeast cells reveal that this mis-rejoining mechanism prevails where both the homologous recombination and non-homologous end-joining pathways are knocked out [B35, B36]. This error-prone mechanism constitutes, therefore, a third pathway for the rejoining of double-strand breaks common to many organisms. Interestingly, in extracts from two ataxia-telangiectasia cell lines, the frequency of mis-rejoining by this mechanism was about 20 times higher than in extracts from normal cell lines [G3, N7]. The genetic basis of this short repeat-driven process is unknown, although in principle it may require similar enzymes to those that recombine adjacent homologous sequences within a chromosome. Adjacent sequences with homology of several hundred base pairs recombine in a process termed single-strand annealing in both mammalian cells and yeast; in budding yeast some of these recombination

events are *RAD52*-dependent, but also require the products of the *RAD1* and *RAD10* genes [O5]. *RAD1* and *RAD10* combine to give a single-stranded endonuclease required to snip off the overhanging DNA strands generated by the recombination of adjacent repeats. In mammalian cells, the homologues of *RAD1* and *RAD10* are the nucleotide-excision repair genes *ERCC1* and *XPF*, respectively [W52], and there is evidence that the *ERCC1* gene is involved in recombination-dependent deletion formation in mammalian cells [S85].




Figure IV. Direct-repeat end joining model.

Direct repeats (shown in boxes) are identified on either side of the initial double-strand break. Exonuclease removes opposite strand beyond repeats. Repeats are annealed, leaving extended tails (snip ends), which are removed. Strands are infilled by polymerase and ligated to give the deleted form with only one copy of the repeat.

96. Ataxia-telangiectasia cell lines were found to have elevated rates (by a factor of 30-200) of intrachromosomal recombination in an introduced DNA construct [M8]. These findings suggest that ataxia-telangiectasia cells have a general disturbance in recombination processes, and that the cell extract studies highlight those events leading to sequence loss and rearrangement (see Section IV.B). It seems likely that an elevated level of non-conservative recombination would also lead to an increase in mutation frequency, provided that the mutation system (i.e. target genomic region) allows detection of relatively large deletions and rearrangements. Evidence of an increased spontaneous frequency of mutations in ataxia-telangiectasia relative to normal persons has been found from direct analysis of their blood cells. Measurements made in ataxia-telangiectasia erythrocytes (loss of heterozygosity in the GPA system, paragraph 213) show loss mutations were elevated, on average, by a factor of 10, while ataxia-telangiectasia lymphocytes (*HPRT* gene mutation) show an average elevation of about 4 [C15]. In each system, however, there was a wide range of frequencies, from nearly normal to more than 100-fold higher than normal. These results are also supported by studies of transfection of small target genes

carrying breaks at specific sites (by endonuclease cutting) on shuttle vectors. The proportion of correctly rejoined target genes was lower in ataxia-telangiectasia than normal cell lines [C13, R5, T6]. It is unclear whether this recombination defect is directly related to radiosensitivity in ataxia-telangiectasia, but disturbance of recombination would tend to give genetic instability and ultimately may lead to cancer-proneness.

2. Radiosensitivity and the cell cycle

97. Mammalian cells vary in their radiosensitivity through the cell cycle, generally showing greatest resistance during the period of DNA synthesis (S phase) and least resistance during G_2 /mitosis and late G_1 (where a long G_1 exists) [S37]. To account for this variation in response, different hypotheses were advanced, such as fluctuation in radioprotective substances in the cell and/or duplication of the genetic material in S phase [S38, T27]. However, a number of studies indicate that the fluctuation in response correlates to the ability of cells to repair radiation damage when irradiated in certain cell-cycle phases [I3, W25]. Additionally, it has more recently been found that certain DNA repair-defective cell lines have altered cell-cycle responses (see paragraph 91). These results suggest that the opportunity for repair of damage in relation to DNA synthesis or segregation is an important determinant of response (see paragraph 9). Accordingly, the ability of the cell to optimize repair and to avoid the interaction of damage with other cellular processes by modulating the cell cycle after irradiation may be crucial to recovery and genome stability.

98. It is now well established that some radiosensitive (*rad*) cell lines in yeasts are defective primarily in regulation of the cell cycle. The paradigm for this type of defect is the *rad9* line of budding yeast. Although identified many years ago by sensitivity to x rays and UV light, it was discovered more recently to be defective for a regulatory function (termed a checkpoint) in the G_2 phase of the cell cycle. The loss of this checkpoint allows *rad9* cells carrying DNA damage to proceed into mitosis and die. The normal *RAD9* gene product appears to respond only to damage, since *rad9* mutant cells have normal division kinetics when unirradiated, although they do show spontaneous chromosomal instability [W6, W7]. Irradiated *rad9* cells have been shown to be defective also in extended- G_1 arrest, indicating that the normal *RAD9* gene product operates at more than one transition point in the cell cycle [S9].

99. Numerous other checkpoint genes have now been identified, especially in lower eukaryotes such as yeasts, and homologues of many of these regulatory genes are present in human cells [C42]. It is clear, therefore, that all cells regulate their cycling in relation to a variety of signals to make sure that a proper progression of phases occurs (e.g. that mitosis does not occur until the S phase is finished) but also to ensure that the genome is intact before starting a new phase. Certain gene products, for example, the tumour suppressor p53, appear to be central to this monitoring process in mammalian cells [A21, C43], and the detailed workings of this regulatory network are currently the subject of intense research.

100. It has been known for many years that x irradiation of mammalian cells in extended-G₁ phase (slowly proliferating primary cultures or cells grown until they run out of nutrients ["plateau phase"]) induces a dose-dependent G₁ arrest [L7, N1]. This arrest is evident at doses as low as 100 mGy, but the duration of the arrest is reduced if the G₁ cells are allowed time to recover before restarting the cycle. Kastan et al. [K2] observed increases in the levels of the tumour suppressor protein p53 correlated to a transient G₁ arrest after relatively low doses (500 mGy) of gamma radiation in normal human haematopoietic cells. Cells that were mutant for p53 showed no G₁ arrest but retained the G₂ arrest found typically after irradiation (but see also paragraph 109). The apparent increase in p53 derives from a prolonged half-life of the protein (the protein half-life is normally only about 30 minutes). The link between p53 and G₁ arrest has been generalized for other cell types and for a variety of DNA-damaging agents [F4]. A likely role for p53, given its function as a transcriptional activator, is to activate the genes involved in negative growth control. Radiation-induced G₁ arrest was found to correlate to inhibition of at least two members of the compound family of proteins that controls cell-cycle progression (the cyclin/cdk proteins); this inhibition was dependent on p53 and was mediated by the p53-inducible kinase inhibitor p21 [D4].

101. The signal for p53-dependent G₁ arrest is likely to be DNA breakage; agents causing DNA double-strand breakage (ionizing radiation, bleomycin, DNA topoisomerase inhibitors) are very effective inducers, while UV light appears to cause p53 induction through the processing of base damage into breaks. The introduction of restriction endonucleases into cells has shown that DNA double-strand breakage alone is sufficient for this response [L14, N5]. Microinjection of broken DNA molecules into cell nuclei has suggested that very few double-strand breaks may be required for arrest, and only one or two may be sufficient [H29].

102. Ataxia-telangiectasia cells lack DNA-synthesis arrest (Section II.A.1) and G₁ arrest, and the DNA-damaging agents that strongly induce p53 are also the agents to which ataxia-telangiectasia cells are sensitive [L14, N5]. It has further been shown that p53 induction is reduced and/or delayed in irradiated ataxia-telangiectasia cells and suggested that the ataxia-telangiectasia gene product (ATM) is part of a signalling mechanism that induces p53 [K3, K6, L14]. A similar alteration of p53 response has been found after irradiation of cells derived from patients with Nijmegen breakage syndrome [J21]. ATM has been shown to physically associate with p53; it is responsible for a rapid phosphorylation of a specific residue at the N-terminal end of p53; this phosphorylation occurs within minutes of treatment of cells with ionizing radiation or radiomimetic drugs but not after UV light treatment [B59, C58, K43, S73]. These processes contribute to its increased half-life following irradiation and also lead to the association of p53 with other proteins involved in damage-signalling pathways. In mouse fibroblasts deficient in both ATM and p53 proteins (or ATM and p21), the loss of the G₁/S cell-cycle checkpoint following irradiation is no worse than for either single defect. This again suggests that

ATM and p53 operate in a common pathway of cell-cycle control in response to DNA damage. However, it seems that ATM may regulate only some aspects of p53-dependent responses; for example, the ATM defect leads to premature senescence in mouse fibroblasts, and in this case the combined deficiency of ATM and p53 [W47] or ATM and p21 [W49] alleviates the senescence.

103. The ATM protein has also been shown to interact directly with the proto-oncogene c-Abl [B48, S65], a protein kinase that itself interacts with other important cell-cycle regulators such as the retinoblastoma protein and with RNA polymerase II. When cells are irradiated, c-Abl is activated in normal cells, but this activation is absent in ataxia-telangiectasia cells. Additionally, cyclin-dependent kinases are resistant to inhibition by radiation in ataxia-telangiectasia cells, and this appears to be due to insufficient induction of p21 [B30]. A kinase named CHK1 that links DNA repair with cell-cycle checkpoints in yeast, has recently been found to be conserved in humans and mice and is dependent on ATM for its activity in meiosis. Like ATM (paragraph 62), CHK1 protein localizes along synapsed meiotic chromosomes, probably at the time that they are repairing DNA breaks. It is speculated that CHK1 acts to coordinate signals from both ATM and ATR (paragraph 65) to ensure the correct progression of meiosis [F17]. The activity of a second kinase, CHK2, which is homologous to the yeast RAD53 protein (paragraph 70), has also been shown to be ATM-dependent; this kinase is involved in the negative regulation of the cell cycle [M23]. It seems likely, therefore, that the ATM protein is involved in multiple signal-transduction pathways, by virtue of its interaction with a number of important regulatory proteins.

104. Alterations in the response of cell-cycle regulators in ataxia-telangiectasia cells may be involved in some aspects of the disorder but are unlikely to be responsible for the radiosensitivity of ataxia-telangiectasia. For example, p53-deficient cells have a tendency to be radioresistant rather than radiosensitive [L6], and SV40-transformed ataxia-telangiectasia cell lines in which p53 function is ablated still have the characteristic radiation sensitivity. It is more likely that any compromise of p53 response will affect cancer-proneness; both human patients (Li-Fraumeni syndrome) and *Tp53*-knockout mice show elevated levels of cancer [D3, M4]. Loss of one (*Tp53*^{+/-}) or both (*Tp53*^{-/-}) gene copies in mice also has significant effects on sensitivity to radiation-induced cancers, seen as a much reduced time for tumour onset. When *Tp53*^{+/-} mice were irradiated with gamma rays to a dose of 4 Gy at 7-12 weeks, the tumour onset time was reduced from 70 to 40 weeks, while *Tp53*^{-/-} mice showed a decreased latency only if irradiated when newborn (because at later times their frequency of spontaneous tumour development was already extremely high, masking the effect of irradiation). It is of interest that p53-deficient mice show a high incidence of thymic lymphomas, similar to ATM- and BRCA2-deficient mice (paragraphs 62 and 73), although it should be remembered that this result may be influenced by the fact that all of these mice were produced in the same genetic

background. In the *Tp53*^{+/-} mice, the normal *Tp53* allele was commonly lost in tumour cells; it was notable that in radiation-induced tumours there was both a high rate of loss of the normal allele and a duplication of the mutant allele [K11].

105. Inactivation of mitotic checkpoint genes has been found in tumour cells derived from both BRCA1-deficient and BRCA2-deficient mice (paragraph 73), suggesting that these defects cooperate in cancer progression. Conditional deletion of the *Brcal* gene led to a low frequency and long latency for mammary tumours, but *Tp53*^{+/-} mice carrying this *Brcal* mutation showed a significant acceleration in mammary tumour formation [X7]. It may be significant that the phosphorylation of BRCA1 in response to DNA damage is mediated by ATM [C67], in view of the potential involvement of ATM in breast cancer (paragraph 137 *et seq.*). In BRCA2-deficient cells, mutations in *Tp53* or genes controlling the mitotic checkpoint were implicated in promoting cellular transformation and the development of lymphomas [L64]. Centrosomes, which control the movement of chromosomes during mitosis, are abnormal in both BRCA1- and BRCA2-deficient cells, leading to unequal chromosome segregation [T46, X8].

106. Mice that lack DNA-dependent kinase (*scid* phenotype, paragraph 82) accumulate high levels of p53 because of the presence of “natural” DNA double-strand breaks from unrepaired V(D)J recombination intermediates but show a typical G₁ arrest following irradiation. These mice develop thymic lymphoma with low incidence (15%) and long latency. The importance of p53 expression in protecting the animals from cancer was shown in mice that lack both DNA-dependent kinase and p53; these mice show prolonged survival of lymphocyte precursors and onset of lymphoma/leukaemia by 7-12 weeks of age (mice defective in p53 alone develop lymphoma to 50% incidence at 16-20 weeks). Cell lines derived from these double knockout mice were also about 10 times more resistant to gamma radiation than lines from *scid* mice. It was suggested that p53 may normally limit the survival of cells with broken DNA molecules and therefore that p53 loss promotes genetic instability [G23, N14].

107. Many of these p53-related responses may be associated with other functions of p53, in particular its role in radiation-induced apoptosis (programmed cell death, Section II.B.3). It is clear that DNA- break-inducing agents, including ionizing radiation, kill some cell types through a p53-dependent apoptotic pathway, although this pathway is not involved in the induction of apoptosis by other stimuli [C6, L13]. Alterations in the control of apoptosis may be linked to carcinogenesis, where a loss of p53 function may lead to the survival of precancerous cells, as has been shown in the induction of skin cancer by UV light [Z4].

108. It has also been known for many years that irradiated mammalian cells may arrest in G₂. The G₂ checkpoint is also controlled by specific cyclin/cdk proteins, and gamma radiation rapidly inhibits the kinase component (p34^{cdc2}) at

doses resulting in G₂ arrest [L12]. Conversely, p34^{cdc2} activation accompanies release of a radiation-induced G₂ block by drugs such as caffeine [H14]. In cultured cells, caffeine-induced release of the arrest decreases survival [B49] and increases the proportion of cells dying from apoptosis [B40]. However, it has also been found that caffeine treatment leads to an increased frequency of chromosomal aberrations [L40, T26], although the link between these two effects has been questioned [H18]. These observations suggest that the arrest promotes survival and may reduce the probability of genetic alterations. This view is supported by experiments with arrest-defective *rad9* yeast cells, where the imposition of an artificial G₂ delay promotes survival [W6]. However, in some radiosensitive mammalian cell lines, including ataxia-telangiectasia, enhanced delay in G₂ is commonly linked to enhanced sensitivity [B3, B6]. It seems likely that because of repair defects these cells never progress further in the cycle and die (that is, the cell-cycle arrest is irreversible, because the damage signal is not reduced). The signal for this response could be unrepaired DNA breaks; recent experiments in recombination-deficient yeast carrying a single unrepaired double-strand break in an inessential DNA molecule show that this can be lethal to a majority of cells [B7].

109. Initially there was little evidence of a role for the p53 protein in G₂ arrest (paragraph 100), but more recent data with p53-deficient mice and cells have shown that p53 is required for correct control of entry into mitosis following DNA damage [B60, S74, S75]. Similarly, cells established from patients with germ-line *TP53* mutations (Li-Fraumeni syndrome) have a consistent defect in G₂ response [G36].

3. Apoptosis: an alternative to repair?

110. Early studies of irradiated mammalian cells recognized that there are different forms of cell death [O1]. Attempts to correlate cell death with the formation of chromosomal aberrations concluded that a major cause of death in fibroblastic cells is genetic damage resulting in the loss of chromosome fragments [B22, C26, D8]. The most conclusive experiments were those of Revell and co-workers, showing a close correlation between loss of chromosome fragments, observed as micronuclei, and cell death in individual diploid Syrian hamster BHK21 cells [J13]. It is envisaged in these cells that genetic imbalance, perhaps due to the loss of specific essential genes, results in death once the pre-irradiation levels of their gene products have decayed. However, cells from a number of different developmental lineages are known to undergo rapid cell death, often termed interphase death, after irradiation [H16, P18, Y4].

111. It is now recognized that the balance between cell proliferation and cell death is crucial to the correct development of organisms, and that cell deaths in many tissue lineages are programmed by a genetically -controlled process known as apoptosis. This process was first described as a characteristic breakdown of cellular structures, including DNA degradation [K14, W33]. Apoptosis is initiated through specific cell surface receptors in response to external developmental signals but

may also be induced by DNA damage. Radiation-induced interphase death is considered to be an example of apoptosis [A3, U14, Y3], although the occurrence of apoptosis has more recently been reported in cell lines dying at various times after irradiation [R8, Y5].

112. Initiation of rapid apoptosis in normal cells by radiation appears to be dependent on the induction of the p53 protein [C6, L13], although delayed apoptosis may occur in cell lines deficient in wild-type p53 protein [B27, H37]. While p53-dependent apoptosis does not involve the same receptors as developmentally-regulated apoptosis, thereafter the molecular pathways converge to a common cell death programme. The mechanism of signalling by p53 is not well understood, although p53 can bind DNA at specific sites ("response elements"), and this binding is strongly stimulated by DNA ends, short regions of single-stranded DNA, and short mismatched DNA segments [J9, L28]. The importance of this DNA binding is demonstrated by the fact that the vast majority of *TP53* mutations identified in tumour cells occur in the part of the gene specifying DNA-binding activity [H25]. Relatively stable binding to DNA (half-life >2 h) can allow a variety of subsidiary processes to occur, such as the activation of specific genes. Because p53 has the properties of a transcriptional activator, it may initiate apoptosis by switching on a specific set of genes. Alternatively there is evidence that in some situations p53 may act in the opposite way, repressing genes concerned with cell survival [C23, C32].

113. A second pathway of radiation-induced apoptosis is proposed to work through a membrane-associated signalling system responding to a variety of extracellular stresses. Sphingomyelin and possibly other sphingolipids at the membrane respond to stress by rapidly releasing ceramide, which accumulates and activates protein kinases to initiate cell-cycle arrest and apoptosis. It has been suggested that ceramide acts as a "biostat", measuring and initiating response to cell stresses in the same way as a thermostat measures and regulates temperature [H34]. Radiation can activate components of the ceramide pathway in isolated membranes, suggesting that DNA damage is not required [H35]; however, this suggestion has been challenged [R39], and recently it has been found that another function of the *ATM* gene is to modulate ceramide synthesis following radiation-induced DNA damage [L67]. Lymphoblasts from patients with the Niemann-Pick disorder have an inherited deficiency in acid sphingomyelinase, an enzyme hydrolysing sphingomyelin to give ceramide, and fail to show ceramide accumulation and apoptosis following irradiation. Comparison of mice with a knockout in the gene for acid sphingomyelinase with *Tp53*-knockout mice shows that the initial stages of the sphingomyelin-dependent apoptotic pathway are distinct from those of p53-mediated apoptosis. These two pathways may be prevalent in different tissues; sphingomyelin-dependent apoptosis prevails in the endothelium of the lung and heart and the mesothelium of the pleura and pericardium, while thymic apoptosis is especially dependent on p53. The acid-sphingomyelinase-deficient mice develop normally, showing also that this apoptotic pathway is distinct from develop-

mentally-regulated apoptosis [S55]. In tumour cell lines where *TP53* is mutated, it has also been shown that resistance to radiation correlates to loss of ceramide accumulation [M1].

114. Whether or not the cell goes into apoptosis in response to a given radiation dose is thought to depend on the availability of other proteins that promote or inhibit further steps in the programme. The details of these steps are still being worked out, but it is clear that a diversity of factors influences apoptosis. For example, cells may become susceptible to radiation-induced apoptosis through functional loss of the retinoblastoma (Rb) protein, a cell-cycle regulator, by either mutation of the *RB* gene or the expression of viral oncoproteins that inactivate Rb [S42]. Conversely, the gene product of the *BCL-2* oncogene, first identified at a site of chromosome translocation in B-cell lymphomas, can block radiation-induced apoptosis [R10]. It has been found that *BCL-2* and related proteins localize to cellular membranes, especially of mitochondria, where permeability transitions are important in regulating the apoptotic process [G25]. The competitive formation of protein complexes by members of the *BCL-2* family is suggested to control susceptibility to apoptosis; some of these proteins are antagonists, like *BCL-2*, while others are agonists, and the relative proportions of these proteins determine whether a cell will respond to an apoptotic signal [K5, K19, W34].

115. The extent of apoptosis for a given radiation dose does not appear to be affected by changes in the dose rate or by dose fractionation [H38, L26, L32], although there has been a recent report of a reduction in apoptosis at very low dose rates from gamma rays (<1.5 mGy min⁻¹) [B66]. There is an effect of cell-cycle stage on the extent of apoptosis; in contrast to the "classical" pattern of resistance (Section II.B.2), mammalian cells appear to be resistant in G₁ and sensitive in S phase [L26, L33]. In cells that are relatively resistant to apoptosis, the oxygen enhancement ratio was found to be similar for apoptosis and for clonogenic cell survival [H36]. The relative biological effectiveness (RBE) of fast neutrons for apoptosis in thymocytes was found to be 1 [W16], in accordance with earlier measurements of interphase death in thymocytes and lymphocytes [G15, H19]. However, for intestinal crypt cells of mice, where there is a highly sensitive subpopulation of cells in the stem-cell zone, the RBE for apoptosis was 4 for 14.7 MeV neutrons and 2.7 for 600 MeV neutrons [H38, H39]. It is not known why the RBE varies for different tissues; both thymocytes and intestinal cells show p53-dependent apoptosis [L13, M39], but it is possible that their relative abilities to repair radiation damage differ (see paragraph 87). The proportion of cells dying from apoptosis, relative to other forms of death, was shown to be constant at different doses (range 2-6 Gy) in cells where the main cause of death was apoptosis [L26]. This result does not suggest, at least for doses in excess of 2 Gy, that cells switch on the apoptotic pathway because their repair systems are unable to cope with the level of damage. In human lymphocytes, significant induction of apoptosis by gamma rays could be measured at doses as low as 50 mGy, and some evidence was found that individuals vary in their apoptotic response [B41].

116. It has been suggested that alterations in the p53-dependent apoptotic pathway play an important part in the sensitivity of ataxia-telangiectasia [M38] and Fanconi's anaemia [R20, R34], but this view has not been supported by other studies [B51, D15, E15, K35, T43]. It has been found that p53-mediated apoptosis is normal in irradiated *Atm* knockout mice [B50], but that *ATM* is involved in the regulation of the ceramide-dependent pathway [L67]. Further study of apoptotic responses, especially p53-independent pathways, are required to clarify these responses, preferably in primary cells from these disorders.

4. Summary

117. A link between radiation sensitivity and immune system defects has been revealed from studies of DNA repair genes; the same gene products are used to assemble functional immune genes and to repair radiation-induced DNA breaks by non-homologous end joining. Cells and animals defective in genes in this pathway are extremely sensitive to ionizing radiation and lack low-dose-rate sparing. While non-homologous end joining seems to be the main mechanism in mammalian cells for DNA double-strand break rejoining, at least two other pathways exist, and in lower organisms double-strand breaks are mostly repaired by the homologous recombination pathway.

118. An important component of the non-homologous end-joining pathway, the *XRCC7* gene encoding the catalytic subunit of DNA-dependent kinase, is related in structure to the ataxia-telangiectasia gene product and has been shown to behave as a tumour suppressor. The mechanism of rejoining by this pathway and the relationship of the components of the pathway to other regulatory processes (such as cell-cycle control) are at present the subject of intense research activity.

119. It is clear that the fidelity of rejoining by the different repair pathways is an important determinant of the mutagenic consequences of DNA damage (see also Figure III). Homologous recombination repair is likely to be the only error-free pathway. However, since even this pathway depends on copying information from another DNA strand, the fidelity must relate to the quality of both the information copied and the enzyme (polymerase) used for the copying. Pathways of double-strand break repair based on illegitimate recombination (non-homologous end joining, repeat-driven rejoining) will commonly be error-prone, although in principle the non-homologous end joining pathway can be error-free where the nature of the damage allows this (e.g. repair of a double-strand break formed by the overlap of two single-strand breaks sufficiently separated to allow for templated repair). Repeat-driven rejoining leads to deletion formation (mis-rejoining), and may act as a back-up for damage that is not (or cannot be) repaired by the other pathways. It is still not clear which factors determine the

likelihood that a specific damage site will be repaired by a particular pathway [T44]. The complexity of damage may influence this question (Section I.B), as well as the possibility that radiation damage induces the activity of specific repair enzymes or pathways (Section III.B). In future work it will be important to establish the relative efficiencies of these different pathways at different doses and dose rates and in different cell types, so that mutagenic consequences can be predicted. The potential consequences of error-prone repair pathways at low radiation doses are explored further in Annex G, "*Biological effects at low radiation doses*".

120. When growing cells are irradiated, they arrest their cell-cycle processes, apparently to allow time for repair to be completed satisfactorily. This arrest is part of a checkpoint function that monitors the physical state of DNA at different stages of the cycle. Cells that have lost a checkpoint may be as radiation sensitive as cells that have lost DNA repair capability. Many genes are involved in controlling the cell cycle and determining checkpoints. In mammalian cells, the p53 protein is important in response to DNA breakage and controls both arrest in the G₁ phase and one pathway of programmed cell death (apoptosis). Animals and humans (Li-Fraumeni syndrome) deficient in p53 show elevated levels of cancer; irradiation of p53-deficient mice has a marked effect on the latency period for tumour formation and gives a high incidence of thymic lymphomas. The ataxia-telangiectasia disorder has defects in cell-cycle checkpoint functions, and there is evidence suggesting that *ATM* and p53 operate in a common pathway of cell-cycle control in response to DNA damage.

121. Apoptosis (programmed cell death) is also induced by ionizing radiation, both through the p53-dependent pathway and a membrane-associated signalling pathway. The relative importance of these pathways varies in different tissues; also, the importance of apoptosis as a mechanism of cell death in response to radiation varies with the cell type and developmental stage. The relationship between apoptotic death and radiation injury differs from that for genetic death (loss of essential genes through damage to the genome). Changes in dose rate or dose fractionation do not appear to affect apoptotic responses, and the response of cell-cycle stage is the reverse of the pattern found for genetic death. While it is attractive to consider that in the face of excessive damage, radiation-induced apoptosis is an alternative to DNA repair, the evidence for this possibility is not convincing. Loss of ability to respond to apoptotic stimuli will allow the accumulation of cells that may carry genetic damage and can therefore be a cancer-promoting event [W8]. This phenomenon is seen, for example, in *Tp53*-knockout mice (paragraph 104) and in mice overexpressing the *BCL-2* protein [M21]. In this respect, the correct functioning of apoptotic pathways can be viewed as a complementary mechanism to the repair of DNA damage, removing damaged cells from the population.

III. HUMAN RADIATION RESPONSES

A. CONTRIBUTION OF MUTANT GENES TO HUMAN RADIOSENSITIVITY

122. The term radiosensitivity is used to indicate an abnormally increased response to ionizing radiation of both the whole body (see Section II.A.1) and cells derived from body tissues. Further, in cells the radiation response may be measured in different ways. For example, the ability of cells to grow and form clones (cell survival) following irradiation is commonly used as a measure of sensitivity, but more sensitive assays based on chromosome damage may also be used. In the comparative measurement of cell survival, the terms D_0 (mean lethal dose calculated as the inverse of the slope for exponential survival curves represented semi-logarithmically), D_{bar} (mean lethal dose calculated as the area under the survival curve in linear representation [K34]), or D_{10} (dose to give 10% survival) are commonly used.

123. The range of sensitivity of cells from supposed normal individuals has been measured in survival experiments following multiple doses of acute low-LET radiation. In two major studies with human diploid fibroblasts, the sensitivity to dose was found to vary by a factor of about 2: D_0 range = 0.98-1.6 Gy (mean = 1.22 ± 0.17 Gy) in 34 cell lines tested [C8] and D_0 range = 0.89-1.75 Gy (mean = 1.23 ± 0.23 Gy) in 24 cell lines [L8]. The degree of interexperimental variation for a given cell line was generally small ($\leq 20\%$), although in one of these studies cell lines derived from a mother and daughter showed survival curves that could also vary by a dose factor of about 2 on repeated testing [L8]. However, it was concluded that neither cell culture conditions (including cloning efficiency and population doublings) nor age and sex of the donor were correlated to the observed differences in radiosensitivity [C8, L8]. This suggests that unknown genetic factors that affect radiosensitivity vary in the cells of normal individuals.

124. More recently it has also been possible to measure cell survival with peripheral blood T-lymphocytes, irradiated as resting (G_0 phase) cells, to assess the range of radiosensitivity in humans. G_0 lymphocytes tend to have a more curvilinear response to dose than cycling fibroblasts and, on average, a higher survival at low doses than fibroblasts. Early-passage fibroblast survival data generally show a good fit to a simple exponential, while the G_0 lymphocyte data are better fitted with models including a dose-squared term, such as the quadratic ($-\ln S = \alpha D + \beta D^2$). Using D_0 as a measure, the range of sensitivity to dose of lymphocytes from different individuals can also vary by a factor of about 2 [E7]; however, given the shape of the survival curves, D_0 is difficult to measure accurately, and the range is generally much less than 2 using measures such as D_{bar} or D_{10} [E7, G17, N4].

125. No correlation was found between the measured sensitivities of fibroblasts and lymphocytes from the same individual [G17, K23]. A similar lack of correlation was

found between lymphocyte and fibroblast responses of pretreatment cancer patients, whether the fibroblasts were cycling or in the plateau phase, suggesting that these differences in response were not cell-cycle-dependent [G11]. The differences between fibroblasts and lymphocytes may reflect several other possibilities: modes of death may differ, cell-type-specific factors may affect the expression of genes that modify radiation response (especially after growth in culture), and/or there may be unknown variables in assay conditions. In support of the last possibility, Nakamura et al. [N4], using donors from one ethnic group, found a similar (relatively small) variation in the mean x-ray dose to kill 28 samples of T-lymphocytes from one individual ($D_{10} = 3.66 \pm 0.21$ Gy), very similar to results with samples from 31 different individuals ($D_{10} = 3.59 \pm 0.18$ Gy). Elyan et al. [E7] reported similar data with donors from three ethnic groups, although with a greater overall range of variability. In support of differences in cell-type-specific factors, other cell types such as keratinocytes also show survival that is significantly different from that of fibroblasts taken from the same individuals [S39].

126. The use of experimental conditions promoting recovery from irradiation, such as low-dose-rate irradiation, will, it is suggested, expand the range of sensitivities to dose shown by normal human cells. The exposure of non-growing fibroblasts from 14 different normal individuals to tritiated water (dose rate = 8.5-100 mGy h^{-1}) expanded the range of sensitivity to dose by a factor of more than 3 [L35]. Similarly, Elyan et al. [E7] found that low-dose-rate irradiation (9.8 mGy min^{-1}) of human G_0 lymphocytes from 19 individuals expanded the range of sensitivity to dose by a factor of about 4. When a higher dose rate (28.5 mGy min^{-1}) was used, however, there was little difference between the range of sensitivities measured for fibroblasts and lymphocytes and the range with a dose rate of 4.55 Gy min^{-1} [G11]. The latter result probably reflects incomplete recovery during the low-dose-rate irradiation period [E7, G11].

127. Other assays of cellular sensitivity have been developed based on the measurement of responses such as the growth of cells after single doses of radiation. A "growback" assay following gamma irradiation of lymphoblastoid cells, developed by Gentner and Morrison [G4], detected a wide range of sensitivity in the normal (asymptomatic) population. Of 270 lymphoblastoid lines tested, about 5% of normal lines showed hypersensitivity after acute irradiation, as measured by their overlap with the responses of ataxia-telangiectasia homozygotes, while on the same criterion about 12% were hypersensitive when assayed using chronic irradiation [G4, G13]. However, a wider range of sensitivity was detected using chronic irradiation, and rigorous statistical tests found that the proportion of lines showing a significantly hypersensitive response ($p = 0.05$) was 5%-6% under both irradiation conditions [G14].

128. There have been claims that a number of disorders with an increased incidence of cancer are hypersensitive to x rays, including retinoblastoma, basal-cell naevus syndrome, Gardner's syndrome, and Down's syndrome [A6, M2, W3]. The main method of determining sensitivity differences has been cell survival assays with cultured fibroblasts (as shown in Figure II), in which small numbers of cell cultures derived from normal individuals are compared with those of one or more derived from the patient(s). Unfortunately, such studies have not always given consistent sensitivity differences for the same disorder tested in different laboratories. In part this may be due to the large range of sensitivity found among normal individuals; if the normal lines used in a given laboratory are in the higher part of the survival range, a disorder showing relatively low survival may be classified as sensitive even though it falls within the overall normal range. However, it has been concluded from a statistical analysis of many published survival curves of human fibroblasts that the radiosensitivity of cells from some disorders, calculated as D_{bar} , can be discriminated from that of normal cells [D7]. The disorders include ataxia-telangiectasia homozygotes and heterozygotes, Cockayne's syndrome, Gardner's syndrome, Fanconi's anaemia, and 5-oxoprolinuria homozygotes and heterozygotes.

129. A number of studies had previously suggested that cells from individuals heterozygous for the disorder ataxia-telangiectasia (carriers with one mutated gene, such as the parents of those showing the full symptoms) are more sensitive than normal to the lethal effects of irradiation [A6, A14, C33, K17, P11]. Additionally, ataxia-telangiectasia heterozygotes show an average level of dose-rate sparing that is intermediate between that of lines from ataxia-telangiectasia homozygotes (paragraph 49) and normal persons [P12, W18, W21]. However, even under low-dose-rate conditions, where a number of carriers have been tested, it is evident that their sensitivity range overlaps that of the most sensitive normal individuals [L35, W18]. Survival assays on cells derived from human sources cannot therefore unequivocally distinguish carriers of a mutated AT gene from normal individuals. Similarly, it has been found that protracted radiation exposure will not readily distinguish other disorders suggested to have minor degrees of hypersensitivity [L35]. Inbred mouse strains may show less variation in response from individual to individual, and mice heterozygous for a defective *Atm* gene were found to show a significant reduction in lifespan following irradiation (4 Gy at 2-4 months of age), as well as premature greying of coat colour, compared to normal littermates [B61].

130. The finding of a range of sensitivities to radiation among supposedly normal persons implies that in addition to those with increased sensitivity, there are individuals with greater radiation resistance than average. There is little evidence, however, for specific disorders associated with radiation resistance. It has been noted that cell lines derived from mice deficient in the p53 tumour suppressor are more resistant to radiation than normal (paragraph 104), although this response may be cell-type-dependent [B43]. Similar

studies on fibroblasts derived from patients with the Li-Fraumeni syndrome, defective in p53, highlighted a family carrying a germ-line *TP53* gene mutation in codon 245. Cells from family members showed a radioresistance (D_0 value) that was increased by a factor of 1.2 [B44], compared to an approximately 1.5-fold increase for the p53-deficient mice [L6].

131. There is evidence from cell survival assays that radiation sensitivity at the cellular level may correlate with tissue response of cancer therapy patients. In a study of 811 patients treated with radiotherapy for breast cancer, five showed severe skin responses. The survival of fibroblast cell lines derived from these five patients was compared to six lines derived from women with normal radiotherapy responses (also from the same group of 811 patients). The sensitivity of the severe-response patients ($D_0 = 0.97 \pm 0.11$ Gy) was significantly higher than that of the normal group ($D_0 = 1.16 \pm 0.08$ Gy; see paragraph 123) [L39]. In a prospective study of 21 head-and-neck cancer patients it was found that the sensitivity of their cultured fibroblasts, but not their lymphocytes, correlated with whole-body late effects [G12]. Similarly, a study of fibroblasts derived from 10 radiotherapy patients showed a correlation between cellular sensitivity and late tissue reactions [B42]. The survival of lymphocytes from breast cancer patients suffering severe reactions to radiotherapy has also been studied in conjunction with ataxia-telangiectasia heterozygotes and homozygotes [W21]. Using high-dose-rate irradiation (1.55 Gy min^{-1}), the survival of lymphocytes from severe-response patients could not be distinguished from that of normal-response patients (or other normal controls), but with low-dose-rate irradiation (9.8 mGy min^{-1}) the cell survival of severe-response patients and ataxia-telangiectasia heterozygotes was similar and significantly different from that of normal individuals. Using the growback assay (paragraph 127), lymphoblastoid cell lines derived from newly presented, non-selected cancer patients showed a similar range of radiosensitivities to the normal lines, while patients already known to show some adverse reaction to radiotherapy showed a higher proportion with extreme sensitivity [G14].

132. Assays based on chromosomal damage have commonly shown a higher degree of sensitivity in the detection of differences between individuals. For example, the genetic disorders ataxia-telangiectasia, Bloom's syndrome, and Fanconi's anaemia were described as "chromosomal breakage syndromes" in early studies, showing significantly increased frequencies of chromosomal breaks and exchanges after irradiation [H24]. Chromosomal responses after low-dose-rate irradiation have also been used to examine the sensitivity of ataxia-telangiectasia heterozygotes [W13] and breast cancer patients who react severely to radiotherapy [J12]. Lymphocytes from 5 out of 16 patients (31%) sustaining severe reactions to radiotherapy had dicentric yields outside the control range [J12].

133. An assay based specifically on the chromosomal sensitivity of lymphocytes to irradiation in G_2 has been successful in detecting significant differences between

individuals carrying various cancer-prone disorders. The work of Sanford et al. is based on the measurement of chromatid breaks and gaps a short time after irradiation of lymphocyte cultures; this assay detected enhanced sensitivity in fibroblasts from a number of disorders, including ataxia-telangiectasia [T25], Fanconi's anaemia, Bloom's syndrome, Gardner's syndrome, basal-cell naevus syndrome, xeroderma pigmentosum, and familial polyposis [P2, S5]. In this G_2 assay, cells from cancer patients and tumour cells showed an even higher frequency of chromatid damage [H21, P3]. This assay has some technical difficulties; extensive tests in another laboratory using identical conditions failed to reproduce many of the results, which led to a modified assay that appears to be more encouraging [S28].

134. The modified G_2 chromosomal assay devised by Scott et al. [S27] has been used to assay the sensitivity of lymphocytes from 74 normal individuals, 28 ataxia-telangiectasia heterozygotes, and 50 breast cancer patients. The ataxia-telangiectasia heterozygotes were clearly distinguished in sensitivity from the majority of normal individuals, but about 9% of the normal individuals showed hypersensitivity. Human fibroblast cultures show a similar result, with four ataxia-telangiectasia heterozygotes distinguished from seven normal individuals [M47]. The range of sensitivities of lymphocytes from breast cancer patients was much greater than the range from normal individuals (42% overlapped the ataxia-telangiectasia heterozygote range). It was proposed that in addition to those mutated genes already identified with familial breast cancer, a number of other genes of lower penetrance may also predispose to breast cancer. It was suggested that these genes may be involved in the processing of DNA damage [H21, S27]. Family studies showed that the lymphocytes of first-degree relatives of sensitive individuals were significantly more sensitive to x rays than expectation; segregation analysis of family members (95 individuals in 20 families) suggested that a single gene accounts for most of the variability, but that a second rarer gene with an additive effect on radiosensitivity may also be present [R36].

135. It has also been found that a subset of patients with common variable immune deficiency (CVID) showed enhanced sensitivity to G_2 chromosomal damage [V1]. Among disorders with primary immune deficiency, CVID and ataxia-telangiectasia have the highest reported incidence of tumours, and both disorders show similar types of tumours, suggesting a common risk factor. This finding may relate to the link noted above (Section II.B.1) between immune system dysfunction and radiosensitivity. However, it is clear that the G_2 chromatid assay will detect enhanced levels of damage in cells showing a normal response in survival assays (e.g. xeroderma pigmentosum cells), and it may be argued that it is primarily a means of detecting genetic instability.

136. Another method based on alterations in the sedimentation of human lymphocyte nuclei is claimed to detect particularly sensitive individuals [S4]. When the repair of

damage is measured by return to normal sedimentation properties after irradiation, it was possible to detect individuals sensitive to radiotherapy and patients with autoimmune disease [H5]. Further, patients showing post-therapy complications were found to be more sensitive by this sedimentation assay than those without complications [D2].

137. It has been suggested that individuals heterozygous for the ataxia-telangiectasia defect are also at increased risk of cancer. Swift et al. [S13] compared prospectively the incidence of cancer in 1,599 relatives of ataxia-telangiectasia patients in 161 families in the United States and found an increased relative risk of 3.8 in men and 3.5 in women for all cancers, and an increased relative risk of 5.1 for breast cancer in women. If borne out, this predisposition would contribute significantly to the cancer incidence in the general population: heterozygotes are much more frequent than ataxia-telangiectasia homozygotes (both genes mutant) and from these data could constitute as much as 5%-8% of all adults with cancer [S12]. However, commentary following the publication of the work of Swift et al. [S13] questioned two aspects of their findings. First, bias in the control group was suggested [B24, K22, W14], although this was contested by Swift et al. [S41]. Second, against a background radiation level of 1 mGy a^{-1} , it was considered surprising that an increase in breast cancer could be detected following diagnostic x-ray procedures of $\leq 10 \text{ mGy}$ [B24, L25]. Differences in dose rate might influence the effectiveness of these two sources, and in other studies an acute dose as low as 16 mGy gave a significant increase in breast cancer in normal women aged 5-9 years at exposure [M26]. However, it is clear that a relatively large difference in sensitivity of the ataxia-telangiectasia heterozygotes to the acute diagnostic procedures would be required to attribute the observed cancer incidence to deleterious effects of one mutated copy of the ataxia-telangiectasia gene.

138. Genotyping of 99 ataxia-telangiectasia families using markers tightly linked to the *ATM* gene showed that 25 of 33 women with breast cancer were heterozygous compared with an expectation of 15 of 33 (relative risk = 3.8; 95% CI: 1.1-7.6) [A18]. Two smaller European studies of cancer incidence among ataxia-telangiectasia heterozygotes tend to support the data from the United States, giving an overall relative risk for breast cancer of 3.9 (CI: 2.1-7.2) [E4]. The average relative risk for other cancers was lower at 1.9, with the European studies showing no statistically significant increase over controls. Using these combined data, the proportion of breast cancer cases due to ataxia-telangiectasia carriers would be about 4%. This small proportion is consistent with an inability in two studies [C36, W31] to detect linkage to AT gene markers in breast cancer families (i.e. on the basis of the risk estimates and gene frequency, most cases of familial breast cancer will be caused by other genes) [E4].

139. The identification of the gene mutated in ataxia-telangiectasia (*ATM*, paragraph 56) has led to further attempts to correlate this gene defect with breast cancer incidence or with severity of response to radiotherapy. In a

study of *ATM* mutations in 38 sporadic breast cancers, only rare polymorphisms were detected, none of which would lead to truncation of the gene (paragraph 57), giving no evidence for an increased proportion of ataxia-telangiectasia heterozygotes in breast cancer patients [V8]. The same workers also looked at 88 unrelated primary breast cancer cases from families previously associated with cancer susceptibility and found three *ATM* mutations (3.4%); this number was considered to be higher than expected by chance, but these mutations did not necessarily segregate with cancer incidence in the families [V9]. Screening a further 100 breast cancer cases from families with a history of breast cancer, leukaemia, and lymphoma revealed only one mutation, consistent with minimal involvement of the *ATM* gene [C65]. Fitzgerald et al. [F21] detected heterozygous mutations in 2 of 202 (1%) healthy women with no history of cancer, compared to 2 of 410 (0.5%) patients with early-onset breast cancer, consistent with a lack of association. Similarly, in a study of 18 families associated with a high incidence of breast and gastric cancers, only one *ATM* mutation was found, and this did not cosegregate with the gastric cancer in the family [B58]. In a study of 41 breast cancer patients showing marked normal tissue response to radiotherapy, one was found with a heterozygous mutation in the *ATM* gene (2.4%), compared with none in a comparable number of control patients [S69]. The conclusion that mutation of *ATM* is not a major cause of radiotherapy complications was supported by studies of smaller numbers (15-16) of breast cancer patients showing severe normal tissue damage following radiotherapy, in whom no *ATM* mutations typical of those in heterozygotes were found [A22, R31]. In contrast, a relatively high frequency of *ATM* mutations has been found for a group of prostate cancer patients (3 of 17, 17.5%) with severe late responses to radiotherapy [H47]. The majority of these studies suggest that heterozygosity for the *ATM* gene is not an important cause of breast cancer susceptibility or severe response to radiotherapy, but most do not have sufficient numbers of patients to exclude completely a role for ataxia-telangiectasia heterozygotes.

140. The association of certain forms of lymphocytic leukaemia with mutations in the *ATM* gene has already been noted (paragraph 59). Individuals with less severe mutations in the *ATM* gene, allowing some expression of *ATM* protein (paragraph 57), may also be at greater risk of developing breast cancer [S78]. It has also been observed that loss of heterozygosity (Section IV.B) at chromosome 11q23 is a frequent occurrence in breast carcinomas [C27, H15]. At least two regions of 11q23 are commonly deleted, and one of these includes the *ATM* gene. Other common human malignancies such as lung, cervical, colon, and ovarian carcinomas, as well as neuroblastoma and melanoma, also show an association with loss of heterozygosity at 11q23 and may include the *ATM* gene region [R9]. In a recent study, 40% loss of heterozygosity was found for markers of *ATM* in cases of sporadic invasive ductal breast carcinoma; in the same tumours, markers for the *BRCA1* and *BRCA2* genes showed 31% and 23% loss of heterozygosity, respectively. Loss of heterozygosity of *ATM* correlated with higher grade tumours and a younger age at diagnosis [R32].

141. It is clearly important to detect individuals in the human population who are hypersensitive to radiation and to understand the connection between radiosensitivity and cancer-proneness. While there are a number of indicators of this hypersensitivity, it has been difficult on the basis of cell survival assays to distinguish normal individuals from those carrying mutant genes giving intermediate levels of sensitivity, when the full sensitivity range of normal cells is taken into account (paragraph 128). In the case of ataxia-telangiectasia heterozygotes, newer assays based on G_2 chromosomal sensitivity (paragraphs 133-134), DNA damage levels measured by the Comet assay [D22], enhanced arrest in G_1 [N2, N3], or enhanced arrest in G_2 [H6, L4, L5] all show more promise, but it is still not clear whether any of the assays will detect these heterozygotes exclusively. The discovery of the *ATM* gene (paragraph 56) has made the task of detecting heterozygotes simpler, but because of the large size of this gene it is still a difficult task to screen large numbers of individuals for defects (see paragraph 139). Since many other genes are known to influence cellular radiosensitivity [Z13], it is likely that the molecular cloning of these genes will have a substantial impact on the ability to determine the importance of genetic predisposition to human radiation risk. Indeed, preliminary studies on nine recently-cloned repair genes including *XRCC1* and *XRCC3* (paragraph 78) showed that relatively common alterations in gene sequence exist in normal individuals. An average of 14 percent polymorphic sites yielding protein sequence variations was found in samples of 12-36 individuals, including some individuals who were homozygous for the variant site [M53, S84]. While the functional significance of these variations has yet to be established, their potential to reduce DNA repair capacity may influence individual response to radiation sensitivity and cancer susceptibility.

B. INFLUENCE OF REPAIR ON RADIATION RESPONSES

142. It is known that DNA repair processes influence the sensitivity of mammalian cells and organisms to radiation. If recovery from radiation damage is compromised, as in cell lines defective in double-strand break repair, the slope of the cellular survival curve is increased. Also, while normal cells show a "shouldered" curve, the repair-defective cells commonly show a loss of this feature. In simplistic terms, it is envisaged that repair processes in normal cells increase the chances of survival, but that as an acute dose increases, the amount of damage temporarily saturates the repair capacity of the cell [G6]. There is very good evidence in yeast cells that these aspects of survival are associated primarily with the repair of DNA double-strand breaks [F3]. It has been possible to make this correlation in yeast, because DNA double-strand breaks can be measured accurately, and mutant cells that are temperature-sensitive for double-strand break repair are available. Thus, as the post-irradiation temperature is altered, it is possible to see a concomitant alteration in double-strand break repair and survival. Unfortunately, it

has proved technically difficult to establish similar quantitative correlations in mammalian cells, which have a much larger genome than yeast.

143. Operationally defined measures of cellular recovery, termed potentially-lethal damage repair or sublethal damage repair, have been used for some years in cellular "repair" studies. Potentially-lethal damage repair is measured by the recovery found when cells are held in a non-proliferative state after irradiation, before respreading into fresh growth medium. Sublethal damage repair defines the recovery seen when survival is measured after splitting the dose over a suitable interval. While it has not been clear whether potentially-lethal damage repair and sublethal damage repair measure the same underlying repair processes, it is notable that both are absent in yeast single-gene mutants lacking DNA double-strand break repair (*rad52* series) [R2, R3]. It has also been argued from kinetic data that double-strand break repair is responsible for potentially-lethal damage repair in yeast [F3]. Similarly, with low-dose-rate irradiation, there is good evidence in yeast cells that DNA double-strand break repair is responsible for the dose sparing observed [F3]. Mammalian cell lines with defective double-strand break repair as well as ataxia-telangiectasia cells have also been shown to lack potentially-lethal damage repair (*xrs*, *XR-1*, ataxia-telangiectasia) or low-dose-rate sparing (*xrs*, ataxia-telangiectasia, *irs2*) (see Section II.A.2) [T30]. A correlation has recently been found for human normal and ataxia-telangiectasia cells, irradiated at high or low dose rates, between the amount of DNA double-strand breaks remaining after repair and radiosensitivity (measured as cell survival) [F18].

144. Mammalian cells irradiated with densely ionizing radiation, such as alpha particles, show both a more sensitive response (increased slope of survival curves) and, commonly, the loss of the curve shoulder region. Thus, characteristic increases in RBE are found as the density of ionization (LET) increases. This may well arise from the LET-dependence of damage complexity [A23, G28, G39] (see Section I.B), and refined measurement of DNA breakage following both x irradiation and high-LET particles has shown that there is a comparative excess of small fragments for the high-LET radiations [L53, R26]. However, several studies of mammalian cells using conventional biochemical techniques show that there is little or no difference in the numbers of DNA double-strand breaks induced by low- and high-LET radiations (RBE = ~1) but that fewer of the breaks induced by high-LET radiations are repaired [B52, C53, J4, L52, P27, P28, P37, R27]. Using these methods, a good correlation has been found between the relative number of non-rejoined DNA breaks and the RBE for mammalian cell inactivation over a wide range of LET [G32, R25]. Plasmid DNA, irradiated to give the same amount of damage by gamma rays or by alpha particles, showed much less repair of the alpha-particle damage when exposed to mammalian cell extracts [H41]. It is suggested that damage induced by high-LET radiations is more complex, because of the increased local clustering of ionizations, and therefore less repairable than low-LET radiation damage (see Section I.B). This finding is supported

by measurement of the rejoining kinetics of large fragments of cellular DNA, where rejoining was generally slower after irradiation with high-LET particles than with x rays [L62]. Additionally, using this method the proportion of mis-rejoined fragments increased with x-ray dose (10%-50% over the dose range 5-80 Gy), while similar experiments with high-LET radiation gave a constant 50% of fragments mis-rejoined at all doses tested [L72, K47].

145. In radiosensitive cells, specifically ataxia-telangiectasia cells [C10] and the *xrs* series [T7], the increase in RBE with LET is very much reduced. This result would be expected if the damage caused by densely ionizing radiation is intrinsically more difficult for normal cells to repair than the damage caused by sparsely ionizing radiation. On this basis, radiosensitive cells show little alteration in effectiveness as the LET increases, because they are already inefficient at repairing low-LET radiation damage. This analysis suggests that the RBE-LET relationship is largely caused by the normal cell's ability to repair low-LET radiation damage.

146. Extensive studies in bacteria have shown that some repair processes are inducible by treatment with DNA-damaging agents [W11]. That is, DNA damage causes an increase in the expression or activity of repair enzymes, mitigating the effects of the damage. This process is part of a much wider series of so-called stress responses by which cells adapt to their environment. Knowledge of such inducible processes in mammalian cells is fragmentary, and in some cases the data are controversial. Evidence for the existence of inducible repair processes in response to ionizing radiation damage comes from several different types of experiment. These may be broadly categorized as (a) adaptive response of pre-exposed cells (see also [U2]), (b) refined analysis of survival in low-dose regions, and (c) direct molecular evidence for the inducibility of specific gene products.

147. Pre-exposure to low doses of tritiated thymidine ($18\text{-}37\text{ Bq ml}^{-1}$) or x rays (5-10 mGy) was found to decrease the frequency of chromosomal breaks in proliferating human lymphocytes irradiated subsequently with a higher dose (1.5 Gy) of x rays [O2, S34]. The aberration frequency is reduced to about 60% of cells not receiving pre-exposure [S30]. Pre-exposure to low concentrations of radiomimetic chemicals, such as hydrogen peroxide and bleomycin, can also reduce the effect of a subsequent high dose of x rays [C35, V4, W29]. This adaptive response is, however, stopped by 3-aminobenzamide, an inhibitor of poly(ADP-ribose) polymerase (paragraph 77), which is itself induced in response to DNA breakage by radiation [S34, W23]. These features of the adaptive response have led to the suggestion that low levels of DNA breakage act as a signal for a response mechanism leading to accelerated repair of radiation damage in mammalian cells.

148. Although increased resistance following low radiation doses has been observed in a number of laboratories, it occurs only in the lymphocytes of some individuals [B21, B26, H13, S22, S26]. Also, lymphocytes from several persons with

Down's syndrome [K16] and fibroblasts from homocystinuria patients [Z11] did not show the adaptive response. The adaptive response is usually found when lymphocytes are proliferating, not resting (G_0) [S31, W15], although there are data suggesting that pre-irradiation in G_0 , followed by a challenge dose in G_2 , gives increased resistance [C63, K16]. These observations have led to suggestions that the increased resistance is a result of cell-cycle perturbation in the lymphocyte populations of some individuals following the pre-exposure (i.e. the cells sampled following the second dose are in a more resistant phase of the cycle than if no pre-dose had been given). This idea has received some experimental support [A9, W27], but recently an adaptive response for chromosomal and cell-killing responses has been found in cells other than lymphocytes when irradiated in proliferating or non-proliferating states [I1, L27, M24, S23]. Thus, the causes of the adaptive response remain controversial, but further work with different cell types has outlined the responses affected and the conditions under which the response is observed.

149. Measuring chromatid breaks or chromosome-type exchanges in lymphocytes, the maximum effect on resistance to a second dose occurs within 5-6 h following the pre-exposure [S34, W15]; the phenomenon may persist for up to three cell cycles [C24, S34]. While pre-exposure to relatively high acute x-ray doses (e.g. >200 mGy) does not give the adaptive response [C46], when 500 mGy was given at low dose rate (e.g. 10 mGy min^{-1}) the adaptive response was found. Conversely, there appears to be a minimum dose rate for the response induction by low doses (e.g. for 10 mGy pre-dose, the dose rate must be >50 mGy min^{-1} for full effect) [S33]. A pretreatment with an acute dose of 20 mGy from x rays will confer resistance to chromosome breakage by 150 mGy alpha particles from radon [W28]; however, when acute high-LET radiation was used to give the pre-exposure, there was no increased resistance to a second dose [K16, W24].

150. Pre-exposure to low doses from x or gamma rays has also been shown in several different cell types to decrease the frequency of cell killing [L27, M24, S32], mutation [K13, R12, S20, U17, Z8], apoptosis [F26], and morphological transformation [A15, R33]. However, the last result is still controversial. Using a near-diploid mouse skin cell line (m5S), Sasaki [S23] showed that x-ray-induced cell killing and chromosomal and mutation responses, but not morphological transformation, are mitigated by acute pre-exposure to 20 mGy from x rays. A malignant derivative of m5S cells lacked the adaptive response, but this was restored along with morphological reversion by transfer into these cells of human chromosome 11. The study also showed that chemicals that either activate or inhibit protein kinase C, which has an important role in signal transduction, either mimicked or abolished, respectively, the adaptive response [S23]. There have been reports suggesting that capacity to rejoin radiation-induced DNA double-strand breaks is greater following low-dose pre-exposure [I4, Z8], as well as data implicating activation of antioxidant metabolism in the adaptive response [B65, Z14].

151. The repair of DNA base damage is also a potentially relevant process; a newly developed ultrasensitive assay for thymine glycols (paragraph 17) has shown that pre-exposure to 0.25 Gy from gamma rays increases the repair of this type of base damage from a subsequent dose of 2 Gy. The initial rate of removal of thymine glycols in human cellular DNA was found to be increased by a factor of 2 following pre-exposure [L60].

152. New methods to measure with improved accuracy the survival response of mammalian cells have led to the discovery that the dose-response curve may initially be steeper at low doses (<1 Gy) than predicted from the curve found at higher doses. This response has been characterized as an initial hypersensitivity at doses up to 0.5 Gy, followed by increased radioresistance of the cell population [L23, M17, W32]. The possibility that the hypersensitive response to low doses is the result of differentially sensitive fractions in cell populations has been rejected on the basis of two observations. First, survival curves for radiosensitive tumour cell lines and an ataxia-telangiectasia line appear to show no changes in sensitivity in the low-dose region. Preliminary data for two further radiosensitive lines, one with reduced DNA double-strand break repair and the other reduced excision repair, also show no increased radioresistance at low doses in comparison to their respective parental lines [S35]. Second, it is argued that the hypersensitivity seen would require one fraction of the cell population to have an unreasonably high level of sensitivity (e.g. about 7% of the cells to be >10 times more sensitive than ataxia-telangiectasia cells) [L24]. In parallel to the adaptive response of lymphocytes, recent data also show that pre-exposure to low doses from x rays or low concentrations of hydrogen peroxide increases radiation resistance in the hypersensitive region. Further, the additional resistance from pre-exposure was transitory, requiring time for development and diminishing after two or three cell cycles [M18]. Again, it has been found that low-dose hypersensitivity is absent with high-LET radiation [M17, M19].

153. Multi-laboratory experiments attempting to measure the dose response for chromosomal aberration induction in lymphocytes from two normal donors at low x-ray doses (down to 4 mGy) also appear to identify departures from fitted dose responses that have been attributed to the induction of repair processes [P15]. It was then suggested that the control value in these experiments was excessively high, biasing the result towards non-linearity [E5]. However, analysis of lymphocytes from individuals in Austria before and after exposure to fallout from the Chernobyl accident, where a peak twofold increase in radiation exposure was recorded, also showed dose-response curves departing from linearity (i.e. decreasing or levelling off) at annual doses between 0.3 and 0.5 mGy [P17]. Further *in vitro* studies with x rays and larger numbers of donors have not confirmed a significant departure from linearity at low acute doses (down to 20 mGy), although statistical variation in the small numbers of aberrations detected at lower doses do not allow conclusive statements on dose response [L37, L38]. These further studies did show an excess of multiply-damaged cells in some donors after

irradiation, leading to the possibility that a small subset of lymphocytes is especially sensitive to aberration induction by very low doses. Similar studies with 15 MeV D-T neutrons did not reveal significant departures from linearity at low doses [P16]. Similarly, a study of chromosomal aberration induction in Syrian hamsters injected with ^{137}Cs to give a whole-body dose of about 0.4 mGy, to mimic the Chernobyl fallout exposure, failed to reveal a significant increase above background values [L61]. Thus, the possibility that very low doses give a higher yield of chromosomal aberrations than expected on the basis of a linear extrapolation from high-dose data remains contentious.

154. Reports suggesting that low doses of ionizing radiation induce repair processes *in vivo* pre-date the cellular studies described above. Irradiation of mice with low doses of x or gamma rays has produced evidence of increased resistance to subsequent higher doses to both somatic and germ cells [C24, F6, W26]. Recent data show that the marked cytogenetic adaptive response of mouse germ cells does not influence the response of the somatic or germ cells of the offspring [C25]. Examination of chromosomal aberrations in the lymphocytes of people exposed occupationally to higher-than-average doses of alpha particles from radon (0.01-16 mGy) plus gamma rays (1-3 mGy) showed that the dose-effect curve was not related to that found at higher doses. Individuals subjected to the lowest doses gave a steep increase in aberration frequency with dose, while at higher doses the curve flattened out [P14]. Tuschl et al. [T33] showed that the people exposed to the higher doses (8-16 mGy a^{-1}) of alpha particles gave higher levels of repair (unscheduled DNA synthesis after UV light damage) in lymphocytes, compared to controls. Hospital workers exposed to low levels of x and gamma rays (maximum annual dose = 28 mSv) showed a reduction in the frequency of chromosome aberrations induced by a dose of 2 Gy to their blood lymphocytes, compared to non-exposed controls [B20]. However, it has been found that the lymphocytes of children living in areas contaminated by the Chernobyl accident showed no evidence of increased resistance to x rays [P9, U16], and separate reports suggests that fewer people from the region of the accident showed the adaptive response in lymphocytes compared with a control group [P19, M52].

155. A growing number of specific genes and proteins have been shown to be induced or repressed following irradiation, mostly using relatively high doses from x or gamma rays (2-6 Gy). While these do not as yet form coherent pathway(s) coordinating response to radiation, they do implicate genes and proteins involved in a variety of important molecular processes (see also Section II.B). Induced proteins include oncogenes/ transcription factors (c-jun, c-fos, interleukin-1, and egr-1), proteins involved in cell-cycle regulation (p53 and cyclins A and B), growth factors, and DNA-metabolizing proteins [PCNA, β -polymerase, and poly(ADP-ribose) polymerase], as well as the products of a number of unknown genes [B25, K15, W20]. The development of methods for rapidly screening hundreds or thousands of genes (micro-

arrays) in one experiment, for changes in levels of gene expression, is beginning to show the extent of transcriptional response to radiation damage. For example, using an array of more than 600 genes, gamma irradiation of a myeloid cell line showed induction of 48 genes by factors of two or more, and many of these genes had not been previously reported as radiation-inducible [A25]. Considering those genes which respond differentially in the presence or absence of p53, it was found using micro-arrays that several genes encoding secreted proteins with growth inhibitory functions were upregulated in a p53-dependent fashion following gamma irradiation. Thus, the p53 response to radiation may also be involved in growth inhibitory effects on surrounding cells, as observed in "bystander" effects [K44]. Methods are being developed similarly for assaying large numbers of proteins, to reveal translational and post-translational responses, using 2-dimensional gel systems coupled to mass spectrometry [B64, C69]. These methods will also revolutionize the classification of normal tissue and disease states, in particular cancer, by permitting a molecular description of the complete profile of gene products present [G40]. Such classifications will be invaluable in understanding the mechanistic basis of cancer induction by agents such as ionizing radiation.

156. Increased or decreased levels of certain gene products have also been found after low doses of radiation, within the range inducing the adaptive response [B18]. Doses of less than 500 mGy gamma rays have been found to induce the expression of a variety of stress-response genes [P35, P36], and dose-responses of several of these genes were shown to be approximately linear between 20 and 500 mGy gamma rays [A26]. Some candidate gene products with a potential role in induced radiation resistance have been identified, including a member of the heat shock protein 70 family (PBP74 [S79]), a heat-shock related immunophilin protein (DIR1 [R37]), ribonucleotide reductase [S81], and the MAPK and PKC protein kinases [S80].

C. SUMMARY

157. When the cells of normal individuals are examined for radiosensitivity, as shown by their survival in culture, variation by a factor of 2 is seen; this factor may be extended to 3 or 4 with the use of low-dose-rate irradiation conditions. It has been suggested that this variation has a genetic basis, but it has been technically difficult to establish this. Additionally, a fraction of cancer therapy patients suffers from severe skin reactions, and cells from these patients commonly show a slightly elevated radiation sensitivity.

158. Assays based on chromosomal damage in G_2 cells have also been used to estimate radiation sensitivity in normal individuals and in breast cancer patients, relative to individuals with known radiosensitivity disorders. The response of a relatively large fraction of breast cancer patients overlapped that of carriers of the ataxia-telangiectasia defect, suggesting that a number of genes involved in response to radiation damage may predispose to this form of cancer.

159. It has been suggested that individuals carrying one defective copy of the ataxia-telangiectasia gene (*ATM* heterozygotes) are at increased risk from cancer, especially breast cancer. A number of studies have recently tested this idea and come to the conclusion that *ATM* heterozygosity is not an important cause of breast cancer susceptibility (or severe response to radiotherapy), but commonly too few patients have been tested to exclude completely a role for *ATM*. Specific types of tumours, such as T-cell prolymphocytic leukaemia, are associated with high levels of mutation of the *ATM* gene (paragraph 59). Additionally, recent evidence suggests that the genes involved in familial susceptibility to breast and ovarian cancers (the *BRCA* genes, paragraph 72) are involved in DNA repair processes and lead to radiation sensitivity when defective in mice.

160. Repair processes affect the shape of survival curves, especially through differences in the processing of DNA double-strand breaks, as seen in specific repair-defective lines. The response to double-strand breaks probably also underlies operationally-defined measures of cellular recovery such as potentially-lethal damage repair and sublethal damage repair. High-LET radiations may not induce a higher frequency of DNA double-strand breaks than low-LET radiations, but high-LET radiation damage is much less repairable. The increase in RBE with LET, found for normal cells, is largely determined by the cell's ability to repair low-LET damage.

161. Evidence for the inducibility of repair processes in mammalian cells is fragmentary. Data from pre-exposures

of cells to low radiation doses, as well as refined survival analyses at low doses (<1 Gy), show an altered response suggestive of an induction process but to date have failed to link this response to known inducible processes. However, recent experiments have shown that certain genes in at least two repair pathways are up-regulated following radiation damage (paragraphs 84 and 151). Additionally, refined methods for looking at protein structure and activity are beginning to reveal insights into more subtle modifications following irradiation. Some damage-response proteins may be activated by post-translational modifications (e.g. phosphorylation, paragraph 102), without any change in the amounts of these proteins, as part of a signalling mechanism promoting the repair of DNA damage. Other proteins have been shown to accumulate in the cell to form discrete foci following irradiation (paragraphs 71 and 74); these foci may represent repair protein complexes accumulating at sites of damage. While there has been little exploration of the dose dependence of these events, the data indicate that modification of cellular response mechanisms can occur following irradiation, raising the possibility that dose responses are altered as a result. Methods for data acquisition on the inducibility of gene products are presently being revolutionized by the introduction of micro-array and complementary techniques, so that rapid progress in this research area is to be expected. The potential importance of inducible repair processes in determining responses to low doses of ionizing radiation is considered further in Annex G, “*Biological effects at low radiation doses*”.

IV. MECHANISMS OF RADIATION MUTAGENESIS

A. MUTATION AS A REPAIR-RELATED RESPONSE

162. The cellular processing of radiation-induced damage to DNA by enzymes may result in a return to normal sequence and structure (correct repair). Alternatively, the processing may fail or may cause alterations in DNA, with the consequence of lethality or inherited changes (mutations). It is also possible that some subtle forms of damage may be tolerated by the cell, particularly if it is non-replicating, and lead to persistent lesions in DNA. These lesions would have to be both chemically stable and not be substrates for repair enzymes and may include some minor types of damage such as methylated bases generated by non-enzymatic alkylation [L68].

163. It is likely that simple base damage or loss in the mammalian genome will commonly lead to base-pair substitutions. In recent years the defined production of single types of damage, at specific sites in DNA molecules *in vitro*, has given insights into their consequences. For example, DNA molecules carrying a single site of base loss (abasic site) have been shown to give rise to base-pair substitutions at that site,

either when introduced on shuttle vectors [G8] or when the *RAS* gene was transfected and stable transformed clones selected [K7]. These substitutions (a form of point mutation) will often give rise to alterations in a gene product so that it works less efficiently or not at all.

164. Breaks in DNA, especially double-strand breaks, are thought to lead to larger alterations such as deletions and rearrangements. Some alterations are very large and are seen as chromosomal aberrations. While there is little formal proof of these relationships between breaks and mutation type, it has been shown that the transfer of restriction endonucleases into mammalian cells, causing site-specific DNA double-strand breaks, gives rise to mutations and chromosomal aberrations [T24]. Additionally, the processing of isolated double-strand breaks in defined DNA molecules by human cell extracts show that these may lead to large deletions of surrounding sequence [T10].

165. Reduction in the efficiency or fidelity of damage repair may lead to an increase in genetic change. This is seen strikingly in the recent discovery that loss of DNA mismatch repair capacity is involved in specific forms of cancer.

Mismatch repair is a form of base-excision repair (Section I.B) that removes bases that have been altered or incorrectly placed (by a polymerase) so that the two strands of DNA do not match in base sequence. The cancer connection was first noted as a high frequency of mutation of short-repeat-sequence (microsatellite) DNA in colon cancers, including hereditary non-polyposis colon cancer. It was rapidly established that the genes determining hereditary non-polyposis colon cancer colocalize with mismatch repair genes on human chromosomes 2 and 3 [B12, L15]. More genes with homology to these mismatch repair genes have since been identified, and at least two of these have also been shown to be mutated in the germ line of hereditary non-polyposis colon cancer patients [N10]. Colorectal cancer is one of the most common human cancers and perhaps the most frequent form of hereditary neoplasia; hereditary non-polyposis colon cancer accounts for as much as 5% of all cases of colon cancer, and the involvement of several genes in the phenotype may account for its prevalence [M20].

166. The high frequency of repeat-sequence mutation was recognized as characteristic of loss of the mismatch repair system (MutHLS), known for many years in bacteria and involving the concerted action of three mismatch enzymes to correct errors caused primarily during DNA replication (such as slippage of strands at repeat sequences) [M27]. Patients with hereditary non-polyposis colon cancer are therefore thought to have a mutator phenotype similar to mutant strains of bacteria with defects in mismatch repair. The role of inactive mismatch repair genes in humans is expected to be similar to the part played by tumour-suppressor genes; a germ-line mutation in one of the mismatch repair genes is followed by somatic mutation of the second gene copy during tumour development [H22, L36]. However, the mismatch repair genes are not thought to be directly involved in cancer; rather, they increase genetic instability and the probability that random mutations will affect those genes critical to cancer formation. This idea does not necessarily mean that mutations alone are sufficient for cancer formation; it is interesting to note that a subset of hereditary non-polyposis colon cancer patients carry numerous mutations in their cells and the expected defect in mismatch repair but have unexpectedly few tumours [P10]. Given the proposed mechanism, it is curious also that the increased cancer risk in hereditary non-polyposis colon cancer patients is selective; while other sites are affected, there is, for example, no increased risk for cancers of the breast and lung [W17]. Recent success at breeding transgenic mice defective in mismatch repair genes has shown that they are viable, but a large proportion develop lymphomas and sarcomas at an early age [B16, D9, R11]. While many of these mice die early, commonly succumbing to T-cell lymphoma, those that survive for more than 6 months develop gastrointestinal tumours, suggesting that these mice may be used as a model for human colon cancer [H50].

167. Cell lines lacking mismatch repair also show an increase in mutations at sites other than microsatellites. In colorectal carcinoma lines, shown to have high frequencies of microsatellite variation, the *HPRT* gene mutation rate was also found to be increased by a factor of more than 100 over the

rate in normal human cells [B23, E8]. In one of these lines, known to have a defective human *MLH1* mismatch repair gene, about one quarter of the mutants had point mutations (frameshifts) at a hotspot within a run of guanine bases in the *HPRT* gene [B23]. An extensive study of one mismatch repair-defective line using the *APRT* gene similarly showed an elevation of frameshift mutations at sites of repeat base sequence, as well as AT → TA transversions at sites of secondary DNA structure [H23]. A potentially relevant site for mutations of this type is in the *BAX* gene, involved in promoting apoptosis (Section II.B.3); more than half of 41 colon carcinomas with microsatellite instability also had frameshift mutations in a run of eight guanines in the *BAX* gene [R19]. These findings are consistent with a role for mismatch repair in correcting base misalignments generated during DNA replication in normal cells. Interestingly, lack of mismatch repair also allows cells to become tolerant of certain forms of induced DNA damage. Alkylation of DNA bases normally produces cell-cycle arrest and/or death and can lead to cancer, but mismatch-repair-defective cell lines are highly resistant to the effects of alkylating agents [B28, K12]. Mismatch-repair-deficient mammalian cell lines also show a small but significant increase in resistance to gamma rays [F19], suggesting that repair of certain types of radiation-induced base damage is recognized by the mismatch repair system (e.g. 8-oxoguanine, which mispairs with adenine, paragraph 24). Mouse embryonic stem cells, heterozygous for a defect in the mismatch repair gene *Msh2*, showed resistance to low-dose-rate gamma radiation (0.004 Gy min⁻¹) but not to acute radiation (1 Gy min⁻¹). On the basis of this result, it was speculated that heterozygosity for mismatch repair may also contribute to tumorigenesis in a direct manner, without loss of the other gene copy [D20]. The mechanism of resistance is thought to follow from a reduction in abortive "repair": mismatch repair normally removes a base that is incorrectly incorporated opposite a damaged base but this repair is abortive, because the damaged base remains in place and may lead to lethality.

168. It is to be expected that the genes involved in other functions required for the maintenance of genome stability would lead to a similar effect of increasing mutation frequency, thereby affecting cancer rates. These functions would include the "proofreading" of DNA synthesis by polymerases and the regulation of DNA precursor synthesis. Indeed, it has been shown that mutations in the exonuclease domains of the DNA polymerase δ gene correlate with high mutation rates in some colorectal carcinoma cell lines lacking changes in mismatch repair genes [D6].

169. Repair enzymes may cause mutations by virtue of their imperfect response to damaged DNA, but in the last few years it has become apparent that many organisms have also retained specific enzymes that introduce mutations into damaged DNA. Thus the loss of certain repair functions can be antimutagenic, and indeed mutant strains of bacteria showing no increase in mutant frequency after treatment with DNA-damaging agents have been known for many years [W11]. Recent studies have defined these mutagenic or error-prone DNA repair processes in bacteria and yeast, and have

revealed that similar processes occur in other organisms, including humans [J23]. It has been found that genes coding for special types of DNA polymerases are responsible for many of the small mutations (base substitutions, frameshifts) occurring spontaneously and after treatment with DNA-damaging agents. In bacteria, for example, when DNA damage blocks the normal replication process, an “SOS” response is activated and more than 20 genes are induced [S83]. Among these genes are polymerases (e.g., DNA polymerase V) with a high affinity for damaged DNA, which are able to continue to synthesise DNA for a few bases in the presence of damage (translesion synthesis) and which commonly put in incorrect bases to give a mutation [R35, T45]. Similarly, in yeast, three genes (*REV1*, *REV3*, *REV7*) are required for much DNA-damage-induced mutagenesis, and these have been found to specify mutagenic DNA polymerase activity required for translesion synthesis [N23, N24]. A human gene named *hREV3* has recently been cloned through its homology to the yeast *REV3* gene and been found to have the properties of a mutagenic polymerase [G41]. As well as this error-prone activity, yeast cells have another specialized DNA polymerase (encoded by the *RAD30* gene) that can perform translesion synthesis in an error-free way. Strikingly, a human gene homologous to *RAD30* (*hRAD30*) has been found mutated in a variant form of the human sunlight-sensitivity disorder xeroderma pigmentosum (see paragraph 26). Since individuals with this disorder suffer a high frequency of skin cancer, this result suggests that error-free translesion synthesis is important to protect against sunlight-induced cancers [J24, M49]. While much remains to be learned about the operation of these specialized polymerases, it seems clear that maintaining a balance between error-free and error-prone pathways of translesion synthesis is important in the determination of rates of (point) mutation in cells.

B. THE SPECTRUM OF RADIATION-INDUCED MUTATIONS

170. Ionizing radiation can induce many types of mutation, from small point changes to very large alterations encompassing many genes. From recent studies of mammalian cells, in which a few genes have been examined in some detail, it is clear that radiation is most effective at inducing large genetic changes (large deletions and rearrangements). This mutation spectrum (proportion of different types of mutations) differs from that found spontaneously or the spectra induced by many other DNA-damaging agents (e.g. ultraviolet light, alkylating chemicals). These other agents tend to induce mostly point mutations, independent of the genomic region assessed. However, it has also been found that the proportion of radiation-induced large genetic changes can vary with the genomic region assessed.

171. The size of genetic changes in a given region of the genome is limited by the extent to which that region can tolerate change. The region may contain genes that are essential for the viability of the cell and organism; if an essential gene is altered or lost, the changes incurred will

usually be lethal [E3, T16, T21]. Thus, although the initial occurrence of radiation-induced genetic changes may be similar at different sites in the genome, lethality will limit both the frequency and the apparent size of mutations recovered. Mutations will be recovered in some genes at low frequency, because they or the region they reside in can tolerate little change (certain genes have been found that, within experimental limits, show almost no radiation-induced mutation [T15]), while mutations in other genes may be detected at relatively high frequency because of cellular tolerance to large changes. Knowing this, it is possible to devise mutation detection systems that tolerate very large changes that would be inconsistent with survival in normal diploid cells. The most extreme example of high-frequency mutation detection is in a system devised by Waldren et al. [W12] that uses as the mutational target an accessory human chromosome 11 introduced into a Chinese hamster cell line. Since no genes on the human chromosome are essential for cell survival, very large changes including whole chromosome loss are tolerated, and mutant frequencies are about 100 times greater than for endogenous genes such as *HPRT*.

172. Much of the mutagenic response to ionizing radiation has been measured, for experimental ease, in regions of the genome that are present at the level of one copy per cell. This situation (monosomy) occurs naturally in parts of the genome such as the X and Y chromosomes in males; also, functional monosomy appears to be quite extensive through mechanisms that switch off gene expression in one copy (as in one copy of the X chromosome in females and in imprinted chromosomal regions). However, most chromosomes and genes are present as two copies per cell (disomic; see paragraph 7). Where the loss of function in one copy of a disomic gene is tolerated without harm (e.g. because of a recessive point mutation), it has been found that the other gene copy is mutated at relatively high frequency by ionizing radiation. Since the two gene copies differ in their functionality (that is, they are heterozygous), this mutation process is commonly termed loss of hetero-




Figure V. Effect of gene location (for hemizygous genes) and copy number on mutant frequency in hamster cell lines carrying two copies of the gene (heterozygous) or single copy (hemizygous) [B11].

zygosity. The frequency of mutation in this disomic heterozygous situation has been shown experimentally to be higher than for the same gene in a monosomic situation [E3, M14, Y2], as illustrated in Figure V for the *APRT* gene, where the mutation frequency difference is about 20-fold [B11]. The reason for this higher mutation frequency in the disomic situation appears again to relate to tolerance of the frequent large genetic changes caused by radiation.

Thus, the first copy, despite carrying a point mutation, still has the remainder of that genomic region intact (including any linked essential genes), and large deletions in the second copy can be tolerated because these do not lead to a complete loss of linked essential genes. In a monosomic chromosomal region, however, there is no other copy (or no functioning copy) present, and the mutant cell will not survive large genetic changes (Figure VI).




Figure VI. Consequences of radiation-induced deletion mutation in a hemizygous (monosomic) and heterozygous (disomic) gene.

Loss of function in gene X is selected for, but the mutation also removes parts of adjacent genes, one of which is essential for cell viability. The mutation results in death of the cell carrying the mutation when the genetic region is hemizygous, but not when it is present in a heterozygous state.

173. Where disomic genes are used to study radiation-induced mutation, the mutation frequency will also vary depending on which of the two copies (alleles) is used as the target. It is to be expected that in addition to the target gene, other linked genes will vary in their functional state (heterozygosity). Thus, if an active allele of the target gene is linked on one chromosome to the active allele of an essential gene, deletions of that chromosome will be severely limited. Conversely, linkage of the target gene to an inactive copy of the essential gene will not be limiting if the homologous

chromosome carries the active essential gene and will allow the cell to survive a large deletion. Thus, a 10-fold difference in mutant frequency has been recorded for the two alleles of a heterozygous *TK* gene in lymphoblastoid cells [A11]. This difference in mutant frequency was associated with a class of slow-growth mutants carrying large genetic changes at the site of the mutable allele of the *TK* gene [A12].

174. High mutation frequencies may therefore mainly reflect tolerance to large genetic changes, although higher

frequencies may also arise from additional mutation mechanisms available in heterozygous cells (paragraph 183). Since ionizing radiation is relatively good at inducing large genetic changes, it will be very effective as a mutagen for disomic genes compared with its effectiveness for monosomic genes (also, in general, ionizing radiation is less effective than potent chemical mutagens when used to mutate monosomic genes). It may be that, for example, an individual inherits a point mutation in one copy of a tumour-suppressor gene, which in itself does not lead to deleterious effects because of the "good" copy still present. However, should this person be subjected to an agent like radiation, which is very effective at inducing large genetic changes, the other gene copy can be readily mutated along with much of the surrounding chromosomal region, because that region is shielded by the presence of the other gene copies. In this way radiation can be seen as a very effective mutagen for specific types of mutational change; indeed it could be argued that for recessive gene mutation, individuals are susceptible to agents such as radiation if they already carry point mutations in disomic genes. This phenomenon is an aspect of predisposition of individuals to specific genetic changes and should not be confused with radiosensitivity (paragraph 122).

175. The prediction that large genetic changes would be found in disomic target genes has been borne out in studies of mice [C19]. Screening progeny after x irradiation of male germ cells revealed that very large chromosomal changes (deletions, rearrangements, and complex changes) occur at high frequency in one copy of disomic regions of the genome; these large changes were consistent with reasonable viability and fertility in the progeny. As expected, when these mutant regions are not shielded by a second normal copy of the genes involved, they are invariably lethal to the mice.

C. MOLECULAR ANALYSIS OF RADIATION-INDUCED MUTATIONS

176. While it can be argued that ionizing radiation has a particular mutation spectrum, it has not been found that any one type of mutation is induced specifically by ionizing radiation. However, detailed analysis of mutations (and to some extent of chromosomal aberrations) has given some possible indicators of differences between spontaneously-occurring and radiation-induced mutations, and of differences between densely and sparsely ionizing radiation.

177. Sequence analysis of x- or gamma-ray-induced point mutations in mammalian cells has shown that a variety of types occur, from base-pair substitutions and frameshifts to small deletions. Analysis of substitutions in the *HPRT* and *APRT* genes has shown that all of the 6 possible types occur, although at different frequencies (Table 3) [G10, M12, N11]. A majority of the radiation-induced substitutions were transversions (alteration of the base from a purine to a pyrimidine or vice versa), while many spontaneously-occurring substitutions were G:C → A:T transitions. A larger proportion of frameshift mutations also occurred in the radiation-induced mutants than spontaneously, and a few more of the radiation-induced point mutations were multiple substitutions (more than one base change in close proximity). Using a small gene target transferred into mouse cells, little difference was found in the mutation spectra for spontaneously-occurring and x-ray-induced mutations [K18]. The frequency of small rearrangements and deletions was increased in both spontaneous and radiation-induced mutants, indicating differences in the balance of mutagenic mechanisms at the site of integration of the transgene compared with those at endogenous genes.

Table 3
Comparison of spontaneous and radiation-induced point mutation spectra

Type of mutation	APRT gene / hamster CHO cells [M12]		HPRT gene / human TK6 cells [N11]	
	Spontaneous	Induced by gamma rays (2.5-4 Gy)	Spontaneous	Induced by x rays (2 Gy)
Base substitution	55 (71%)	19 (66%) ^a	10 (55%)	19 (54%)
Transition	31 (40%)	6 (21%)	6 (33%)	7 (20%)
Transversion	24 (31%)	13 (45%)	4 (22%)	12 (34%) ^a
Frameshift	6 (8%)	5 (17%)	3 (17%)	8 (23%)
Small deletion or rearrangement	16 (21%)	5 (17%)	5 (28%)	8 (23%)
Total	77	29	18	35

^a Includes two tandem substitutions.

178. Radiation-induced point mutations were found at sites widely distributed within the gene, while spontaneously-occurring point mutations tended to cluster at certain sites [G10, M12, N11]. Differences in the types of base damage responsible and in the randomness of damage induction are likely to explain these differences.

179. Smaller numbers of large genetic changes have been sequenced because of the difficulty of locating and cloning the breakpoints of large deletions and rearrangements. No new or specific mechanism has been found for the induction of these larger changes. In the hamster *APRT* gene, present in the hemizygous state, the large gamma-ray-induced deletions tend

to have short direct or inverted repeat sequences around their breakpoints and to fall into regions rich in adenine-thymine base pairs. Insertions, however, involved repetitive sequences and were accompanied by short deletions [M13]. It was notable in these studies that large changes extending downstream of the *APRT* gene are not found, suggesting that an essential gene is present in this region.

180. In the human *HPRT* gene, also in the hemizygous state, large deletions are found to extend both upstream and downstream of the gene over a region of about 2 Mb [M16, N9]. Radiation-induced large genetic changes tend to eliminate the whole *HPRT* gene, making their analysis difficult. Therefore a series of flanking markers has been identified to delimit these changes and ultimately to map the positions of the breakpoints. Sequence analysis of a few deletion mutations has again shown the presence of short direct repeats at some breakpoints induced by both x rays and alpha particles, as well as other sequence features in adjacent regions [M15, S18]. The large deletions induced by radiation frequently include sequences adjacent to *HPRT*, and it has been suggested that this may be a signature of radiation-induced events as opposed to spontaneous events [N17]. This type of mutation was equally common with gamma-ray doses of 0.2 or 2 Gy d⁻¹. Use of methods amplifying specific regions of the genome to measure rapidly the frequency of rearrangements following x- or gamma-irradiation, followed by sequencing the breakpoints, has given little indication of the involvement of specific sequence features or clustering of breakpoints [F22, F23].

181. Shuttle vector systems consist of small defined DNA molecules that may be mutated in mammalian cells but rescued into bacteria for rapid analysis of sequence changes. While these facilitate the molecular analysis of induced mutations, they generally have the drawback that the target genes are very small and are flanked by essential sequence. This means that only point mutations are detectable, and the major class of radiation-induced mutations (large deletions and rearrangements) cannot be analysed. However, Lutze and Winegar [L19] devised a large shuttle vector based on Epstein-Barr viral sequence that could be maintained episomally in human lymphoblastoid cells and could detect changes of up to 8 kb or so. Using this vector, the mutation spectra of both x rays (150-600 Gy) and alpha particles (3 Gy) from radon gas were studied [L19, L20]. A larger proportion of deletions was found among alpha-particle mutations (64%) than among x-ray mutations (13%), but in a subsequent study with lower doses of x rays, more deletions were found (41% at 100 Gy, 33% at 20 Gy) [L21]. For both x rays and alpha particles, these deletions were large (>2.4 kb), and their breakpoints were clustered in specific regions of the shuttle vector. The breakpoints were commonly associated with short direct sequence repeats of up to six base pairs. Use of the same vector system in a lymphoblastoid cell line defective in double-strand-break repair gave a slightly higher frequency of deletions after x irradiation, in association with an additional class of small deletion mutations (<2.4 kb) [L21].

182. The presence of sequence features such as short direct repeats of a few base pairs at large deletion junctions suggests that illegitimate recombination (see Section I.B) has driven the mutation process. Recently it has been possible to reconstruct the process of illegitimate recombination in cell-free conditions to show that the process can be associated with a DNA double-strand break. DNA molecules are broken at a specific site, using an endonuclease, and exposed to extracts from human cells for a brief period. Analysis of the extract-treated DNA shows that while the majority of broken molecules are correctly rejoined, a small fraction (about 0.5%) are mis-rejoined to give a deletion. The mis-rejoin mechanism involves the pairing of short direct sequence repeats, situated either side of the break, so that the intervening sequence (including one of the repeats) is deleted [T10]. This mechanism has been found in both radiation-induced mutations and in germinal and somatic-cell mutations in humans. A model for this process is shown in Figure IV. In addition to simple deletion, this cell-free system detected a small fraction of more complex changes, e.g. a large deletion associated with the insertion of several hundred base pairs.

183. In heterozygotes, in addition to the tolerance of large changes, additional mechanisms of mutation may occur. The two gene copies may undergo mitotic recombination or non-disjunction, leading to the appearance of mutations. However, where loss of heterozygosity is measured simply by the presence or absence of the active (wild-type) gene, it is difficult to distinguish mutations occurring by these additional mechanisms from large deletions. In a study of mutation in the human *TK* gene, using both linked marker analysis and densitometry to assess gene copy number, about 50% of the spontaneous mutants appeared to involve recombination, while x-ray-induced mutants were mainly deletions [L29]. Recombination was similarly found to be involved in spontaneous mutation of the *APRT* gene in human cells, but this mechanism was also detected in a number of gamma-ray-induced mutants [F8].

184. While it is difficult to compare somatic-cell mutation data with animal germ-cell data, some recent molecular analyses of the genes used in mouse specific-locus tests indicate possible similarities in the types and mechanisms of mutation. For example, with some dependence on both cell stage and radiation quality, a large proportion of *albino* (*c*) locus mutations on mouse chromosome 7 recovered from germ-cell irradiations are large deletions [R14]. Molecular analysis of radiation-induced mutations at the *c* locus shows that a genomic region of 1.5-2 Mb around the tyrosinase gene may be deleted in mutants without leading to inviability in homozygotes [R13]. This target size is very similar to that of the human *HPRT* gene region used in somatic-cell mutation studies. Also, mutation frequencies per unit dose of low-LET radiation at the *c* locus in mouse germ cells and at the *HPRT* locus in human somatic cells are in order-of-magnitude agreement [T19], given that mutations at the *c* locus occur with about average induced frequency relative to other loci in specific-locus tests [S29]. Comparisons of this type emphasize that the mechanisms of radiation mutagenesis are similar in

somatic and germ cells, and that if measurements are made under similar conditions of mutation tolerance, then the mutation frequencies will be comparable.

185. Comparison of two human lymphoblastoid cell lines derived from the same original cell line has given further insights into the mechanisms of radiation mutagenesis. Line TK6 is more sensitive to the lethal effects of x rays than line WIL2-NS, but the converse is true for mutation response to x rays (although not to chemical mutagens). The increased frequency in WIL2-NS at the hemizygous *HPRT* gene was modest (factor of 4 at 2 Gy), while the increase for the heterozygous *TK* gene is by a factor of 20-50, depending on the *TK* allele used as the target [A13]. Molecular studies have revealed that all *TK* mutants in WIL2-NS arose by loss of the active allele and linked markers (over a 5 Mb region), while point mutations and less extensive deletions were common in the TK6 line. Two copies of the *TK* gene were present on a majority of karyotypes in the WIL2-NS mutants, as expected if these arose from mitotic recombination rather than deletion, while the converse was true for TK6 mutants [X1]. It seems likely, therefore, that the WIL2-NS line has a higher frequency of recombination, leading to a greater ability to survive x-ray damage but incurring a concomitant increase in mutation. Analysis of the p53 status (see paragraph 112) of these two cell lines has shown that p53 protein levels are four times higher in WIL2-NS than in TK6, because of a mutation in exon 7 of the *TP53* gene in WIL2-NS (TK6 is wild-type) and that apoptotic death is substantially delayed in WIL2-NS [X2, Z7].

186. In a study of B-cell precursors from *Tp53*-knockout mice, it was also found that high frequencies of *HPRT* mutations occurred following x irradiation but that these resulted from a preferential survival of mutant clones rather than from a p53-dependent increase in mutation rate [G26]. It was concluded that loss of p53 function allows mutated cells to survive that would otherwise be eliminated by apoptosis. Other recent studies with cells from p53-deficient mice have similarly concluded that they do not have an intrinsically higher-than-average mutation rate [C68, N12, S21].

187. It has been suggested that a relatively unique feature of radiation mutagenesis is the induction of complex genetic changes [M11]. For example, a radiation-induced deletion may be associated with a rearrangement at the same site, and the rearrangement may be an exchange, inversion, or insertion. Additionally, complex changes may be seen as chromosomal exchanges involving several sites in the genome following a single acute radiation treatment [S15]. However, because of their large size and complexity, these changes have been difficult to analyse in cellular genes at the molecular level. Further molecular analyses will be necessary to establish whether or not large mutations and chromosomal aberrations are formed by similar mechanisms. It is clear from studies with one or two genetic regions, and especially the region containing the human *HPRT* gene (Xq26), that genetic changes

identified at the molecular level as mutations can extend to sizes that are visible in the light microscope [S17]. However, the breakpoints of such very large mutations and chromosomal aberrations have not been sequenced.

D. EFFECT OF RADIATION QUALITY

188. In general, high-LET radiation induces a higher frequency of mutants in rodent and human cells, per unit dose, than low-LET radiation [B17, B19, C22, C30, F7, G18, H20, L31, M22, M25, S88, S89, T16, T18, T32, W22]. The RBE varies with LET, peaking at 100-200 keV μm^{-1} , with values as high as 7-10 found for alpha particles and heavy ions in this LET range [C22, T16]. In some mouse cell lines, a relatively small RBE of 2-3 has been found for *HPRT* mutant frequency with alpha particles in the peak range, and when cell survival is taken into account, there is no difference in the effectiveness of alpha particles and x rays. This result appears to be due to the very high effectiveness of low-LET radiation on these cells, as reflected in increased cell killing and mutagenesis for a given dose compared with other rodent lines [B19, I2]. That is, the mutagenic effectiveness of low-LET radiation is more variable than that of high-LET radiation, presumably because cell lines have different abilities to repair low-LET radiation damage (while high-LET damage is less repairable; see Section III.B). An example of this variability is shown in Figure VII [T28].




Figure VII. Fitted curves of mutant frequencies in Chinese hamster V79 cells [T18] and mouse ascites (EAT) cells [I20] induced by x rays and ^{238}Pu alpha particles (see paragraph 172).

189. A study of 19 non-smoking people living in houses with a range of radon concentrations (21-244 Bq m^{-3} in the United Kingdom suggested that there was a correlation of *HPRT* mutant frequency in blood T-lymphocytes with radon concentration. These data suggested that alpha particles in this LET range have a considerably higher effectiveness than found previously in high-dose experiments with human somatic cells. While the estimation of

alpha-particle dose to circulating blood lymphocytes is difficult, the authors estimated an annual doubling dose of 2.1 mSv from their results [B29]. A number of problems attend such measurements, not the least of which is the sample size. In a follow-up study by the same authors with a total of 65 persons from 41 houses in the same town, no significant correlation was found between mutant frequency and radon levels, even for people living in the same house [C34]. In a study in Belgium of 24 people living in houses with radon concentrations exceeding 100 Bq m⁻³, a negative association between radon concentration and *HPRT* mutation frequency was found [A10]. This result led the authors to suggest the induction of repair processes by low levels of radon.

190. The *TP53* tumour-suppressor gene is commonly mutated in many different types of cancers, with a frequency ranging from <10% to >50% depending on tumour type. Analysis of mutations in the *TP53* gene in lung tumours from uranium miners initially gave some hope that these would provide a means of fingerprinting high-LET radiation damage. Vahakangas et al. [V2] sequenced exons 5-9 of the *TP53* gene in 19 lung tumours from miners working in New Mexico. They found several point mutations, although these lacked a class of base substitution (G:C → T:A transversions) and were not at the hotspots described for lung cancer. In contrast, in a larger study of 52 lung tumours from Colorado uranium miners exposed to a dose that was, on average, five times higher, Taylor et al. [T12] sequenced the same exons and found that about half of the substitutions were of the G:C → T:A type in codon 249 (codon = base-pair triplet encoding one subunit of the p53 protein). This transversion is very rare among *TP53* mutations in lung tumours (many of which are presumed to be smoking-associated) and was suggested to be a potential marker for radon-associated lung cancer. At present, apart from the difference in average dose, it is difficult to see why these two *TP53* mutation studies gave such contradictory results. One possibility is that the transversion mutations seen are a result of some other agent in the environment of the mine; if the mine is damp, then fungal toxins similar to aflatoxin B1, known to induce this type of mutation, might be present [V3]. Follow-up experiments looking specifically for the codon 249 transversion in lung cancers from individuals exposed to high domestic radon levels in the United Kingdom [L48] and from German uranium miners [B45] failed to find any examples of this mutation.

191. In a study of thymic lymphomas in RF/J mice induced by gamma rays or by neutrons (0.44 MeV), a similar frequency of lymphomas was associated with mutations activating *RAS* oncogenes (24% for gamma rays, 17% for neutrons). Of the gamma-ray set, 89% were K-*RAS*-activated, with the majority (7 of 8) having a GAT → GGT mutation in codon 12. However, in the neutron set no particular mutation site predominated, and one lymphoma contained a K-*RAS* gene activated by a point mutation in codon 146, a site not previously associated with any human or animal tumour [S36].

192. There are conflicting data on possible differences in the spectrum of large genetic changes induced by high-LET radiation relative to those induced by low-LET radiation. Early cytogenetic studies with diploid human fibroblasts suggested that a greater proportion of *HPRT* mutants induced by high-LET radiation carried large genetic changes than did x-ray-induced mutants [C37, T19]. Studies of specific-locus mutations in mouse spermatogonial stem cells had also concluded, based on a number of criteria, that ²³⁹Pu-induced mutants carried more severe genetic damage than mutants induced by low-LET radiation [R16]. However, classification of *HPRT* mutations by molecular analysis in both diploid human cells and hamster cells has not shown differences in the proportions of large deletions to point mutations at doses of low- or high-LET radiation giving about 20% cell survival [A8, G16, S18, T20]. A confounding factor (commonly underestimated) in determining mutation spectra can be the non-independence of mutants when grown in bulk culture after irradiation [T22], but in the most recent studies [A8, S18] the experimental design ensured the independence of all mutants. Other studies have reported differences: irradiation with 190 keV μm⁻¹ Fe ions gave 82% large deletions of the *HPRT* gene in human TK6 cells relative to 54% for x rays [K21]. High doses of radiation from incorporated ¹²⁵I, giving 1% survival of human TK6 cells, were reported to increase the frequency of deletions or rearrangements when compared with lower doses of ¹²⁵I or with x rays [W22]. It was noted that these high-dose mutations commonly showed loss of part of the *HPRT* gene rather than the entire gene. It has similarly been reported in recent studies with the same cell line that the average size of radon-induced deletions of the *HPRT* gene is not as great as that induced by x rays [B19, C29]. This feature has also been noted for mutations induced at the hemizygous *DHFR* gene in hamster cells; that is, the frequency of total-gene deletions was much higher for gamma-ray- than for alpha-particle-induced mutations, while the reverse was true for intragenic deletions [J11]. In the latter study, it was also found that some alpha-particle-induced deletions shared common breakpoint sites, indicating a non-random distribution. Use of the heterozygous *TK* gene as a target, also in TK6 cells, showed again that the proportion of deletions was similar for x rays and high-LET radiation (4.2 MeV neutrons and argon ions), but that high-LET radiation induced a category of large rearrangements not found for x rays [K20]. It has also been claimed that high LET radiations induce non-contiguous deletions that are not found with x or gamma rays (see also paragraph 208) [S90]. A number of these studies, therefore, suggest there could be differences in the types of large genetic changes induced by high-LET as opposed to low-LET radiation. Much more careful analysis is required before any general statement can be made about the extent or nature of these possible differences.

193. The mutagenic effects of a single alpha-particle traversal were measured with a high-frequency detection system (paragraph 171). Using a microbeam to localize 90 keV μm⁻¹ alpha particles to cell nuclei, a dose-dependent increase in mutant frequency was found for traversals of 1-8 particles, with a twofold increase for 1 alpha particle traversal (cell surviving fraction = 0.8) [H42]. Molecular analysis of

these mutants gave some support for the view that higher doses of high-LET radiation induce larger genetic changes, but this system is unusual in being able to sustain loss of the whole target chromosome without lethality.

194. It has been proposed that the ratio of specific forms of chromosomal aberration may be a fingerprint of exposure to high-LET radiation [B13, S71]. A low ratio (about 6) of interchromosomal to intrachromosomal exchanges was found in several studies of aberration induction by high-LET radiation, while a value of ≥ 15 was found for x or gamma rays. However it was subsequently demonstrated [S54], using a simple two-dimensional chromosome model allowing the prediction of exchange sites relative to track structure, that more densely clustered tracks are not predicted to give a significantly lower ratio of aberration types than random scattering of tracks. More critical assessment of experimental data also does not support this theory [L54, S66]; in particular, reexamination of extensive plant data has ruled out both this theory and any LET-dependence in the ratio of interstitial deletions to inter-arm exchanges [S72]. A different high-LET fingerprint has more recently been proposed by Lucas [L55], who suggested that the proportion of unrejoined ("incomplete") chromosome aberrations will be greater with high-LET radiation, because the local density of DNA double-strand breaks is greater than for low-LET radiation and competition between broken ends leads to a greater chance of incomplete exchange events. Published data are cited that give a ratio of incomplete exchanges to complete exchanges (translocations) of about 9 for low-LET radiations and about 2 for high-LET radiations, with "mixed-LET" radiations giving intermediate values. While this theory has yet to receive critical appraisal in the literature, it may suffer from the same problems as the earlier theory, insofar as only certain sets of data give large differences between high- and low-LET radiations, partly because of difficulties in correctly defining incomplete aberrations. There may be further difficulties with the idea that competition between broken ends will necessarily lead to more incomplete aberrations and with the simple assumption that incomplete aberrations represent unrejoined breaks.

E. NOVEL MECHANISMS OF GENETIC CHANGE

195. Most of the processing of DNA damage by cellular repair enzymes is completed within a few hours of irradiation, including the fixation of mutations. However, there is evidence that cellular responses continue to occur for much longer periods, over many cell generations. These responses include delayed cell death and genetic changes. While these responses may in part be attributed to the time taken for the cell to recover from irradiation, there are in principle several reasons why persistent effects could occur. These include persistence of the damaging agent; persistence of certain forms of DNA damage, i.e. lack of repair; the repair of damage leading to rearrangements of the genome, which themselves upset the correct functioning of the cell (e.g. "position effects" on blocks of genes); and the induction of a

long-lived metabolic disturbance in somatic cells, such that enzymatic activities (e.g. DNA polymerases) involved in the fidelity of maintaining the genome do not function properly.

196. It has been known for many years that cells may take some time to die following irradiation [E2, J7], but more recently emphasis has been placed on lethal mutations that may take effect many cell generations after irradiation [B53, G9, S16]. These two phenomena may in part be aspects of the same response. Delay in cell death in fibroblastic cell lines may be explained by the time taken for cells to show the effects of loss of essential genes through chromosome fragmentation (see paragraph 110). As cells divide, they segregate broken chromosomes (often seen as micronuclei), and the encoded gene products are diluted out of daughter cells, eventually causing death [J7]. In the same way, lethal mutations may cause late-onset death in cells; however, they can arise through more subtle effects on the genome than simple chromosomal fragmentation and loss. Thus, all types of mutation, from point mutation to large genetic changes, may represent lethal mutations if they lead to loss of essential gene products. However, to account for the high frequency of lethal mutations at long times after irradiation, it has also been proposed that some form of persistent genetic instability can be induced by radiation.

197. In addition to delayed death, the mutation frequency in inessential genes such as *HPRT* has been found to be persistently elevated in a large fraction of clones surviving irradiation [C20]. In these experiments, cells irradiated with x rays (12 Gy) were grown as separate clones and examined for both the proportion of cells surviving and for mutant frequency at different times after irradiation. It was found that clones showing reduced survival also commonly had elevated mutant frequencies at the *HPRT* gene for as many as 50-100 cell generations following irradiation. The mutant frequency in individual clones was highly variable but sometimes exceeded $1 \cdot 10^{-3}$. These delayed mutations appeared to be predominantly point mutations [L34].

198. Chromosomal aberrations have also been found to persist following irradiation. One-cell mouse embryos irradiated with x rays or neutrons showed an approximately linear increase in the frequency of chromosomal aberrations per cell in the first, second, and third mitoses post-irradiation. The relatively high frequency of aberrations, especially for neutrons, and the occurrence of chromatid-type aberrations on the third mitosis following irradiation suggested that new aberrations were being produced in post-irradiation cell cycles [W40, W41]. Similar results were obtained with x irradiation of two-cell mouse embryos [W42]. Delayed chromosomal aberrations were not observed in embryos treated with restriction endonucleases, suggesting that lesions other than DNA double-strand breaks are responsible for this effect [W48]. A significant increase in chromosome- and chromatid-type aberrations was also found in cell cultures derived from foetal skin biopsies of mice x irradiated as zygotes [P25]. The study of chromosomal aberrations in lymphocyte and fibroblast clones surviving x irradiation has similarly given evidence of persistent genetic effects [H30, M34, P33]. Analysis of clones

for two months following irradiation showed that >20% had sporadic aberrations as well as transmissible clonal karyotype alterations [H31].

199. Genetic instability in clones of cells surviving irradiation has also been described for alpha-particle irradiation of cultured haematopoietic stem cells from CBA/H mice [K8]. In this case, chromosomal aberrations were measured in cell clones surviving 3 Gy from x rays or 0.25-1 Gy from alpha particles (0.5 Gy from ^{238}Pu alpha particles corresponds to an average of one track per cell). About 50% of the clones surviving alpha-particle irradiation carried aberrations; these were mostly non-identical chromatid-type aberrations, suggesting that they had arisen many generations after irradiation. The frequency of such delayed aberration induction in x-ray survivors was only about 2% [K8]. This form of chromosomal instability in bone marrow cells was also found to be transmissible *in vivo*, by transplanting male cells irradiated with alpha particles into female recipients [W38]. The repopulated haemopoietic system showed instability persisting for up to one year. Alpha particles have also been shown to induce similar delayed chromosomal effects in the bone marrow of two out of four normal humans [K9]. It was suggested that the lack of effect in some individuals reflects genetic determinants that vary in the human population, and additional studies of other inbred mouse strains have also been found to show varying levels of this form of genetic instability. The alpha-particle-induced instability was, however, found to be independent of the p53 status of the cell [K32].

200. Heavy ion (neon, argon, or lead) irradiations have also been found to induce chromosomal instability in cultured human fibroblasts [M35, S53]. Analysis of mass cell cultures for up to 25 passages following irradiation showed that the frequency of aberrations declined at first but then increased until >60% of the cells showed aberrations. In contrast to clonal cell populations, these aberrations showed that the telomeric regions of specific chromosomes (1, 13, and 16) were involved. In a study of alpha particles of different LET, the frequency of micronuclei was found to be increased over that in unirradiated hamster cells seven days after irradiation, even at a dose giving one alpha-particle traversal per cell nucleus [M36]. Calculations indicate that the target size for this effect exceeds the size of the nucleus, suggesting that direct DNA damage by radiation tracks is not causal.

201. It has been speculated that these events indicate that ionizing radiation may induce an "untargeted" mechanism of mutagenesis in cells, as a result of the epigenetic alteration of enzymatic pathways controlling genomic stability [L18]. The idea that DNA may not require a direct hit from radiation to have an increased frequency of genetic changes has received support from the measurement of non-mutational responses. With reference to the process of carcinogenesis in particular, the work of Kennedy et al. [K10, K11] suggests that radiation may induce high-frequency events that predispose the cell towards further (spontaneously-occurring) changes in the process of malignant cell transformation. Furthermore, a study of the induction of sister-chromatid exchanges in

immortalized hamster cells by very low doses of ^{238}Pu alpha particles claimed that induction could be measured at doses (0.3 mGy) where <1% of the cell's nuclei were traversed by an alpha-particle track [N8]. Subsequent studies of primary human fibroblasts confirmed this finding; low doses of alpha particles give three times as many sister-chromatid exchanges [D13] or a five times higher *HPRT* gene mutation frequency [N25] than predicted from the number of nuclei traversed by the particle tracks, giving a considerably larger target size for this effect than expected from nuclear dimensions [D13]. Similarly, the use of a physical barrier to protect one part of a cell population from alpha-particle irradiation showed the expected reduction in killing of mouse bone-marrow stem cells but did not reduce chromosomal instability [L63]. These data are considered as evidence for the existence of a "bystander" effect; that is, damage signals may be transmitted from irradiated to neighbouring unirradiated cells. In a refinement of this type of experiment, an alpha-particle microbeam was used to give precise irradiation of the cytoplasm of cells, without damaging their nuclei. Under these conditions, an average increase of three-fold was found in the mutation frequency in a sensitive human-hamster hybrid cell line (paragraph 171), and the mutation spectrum was similar to that occurring spontaneously [W51]. One explanation of this phenomenon is that reactive oxygen species are generated by radiation in the whole cell and perhaps also in the surrounding medium, and these species or stable reaction products (e.g. lipid peroxides) diffuse into the nucleus and persist to cause chromosomal effects. Evidence for this idea has been found with x or neutron irradiation of bone marrow cells; various indicators of persistent oxygen radical activity were found in cell cultures seven days after irradiation [C40]. It has also been shown that alpha particles can produce factors in culture medium or in cells that cause increases in sister-chromatid exchanges, as noted above; these factors are inhibited by superoxide dismutase, an enzyme that catalyses the conversion of superoxide ions produced in water radiolysis [L17]. Similarly, in the microbeam experiments, reactive oxygen species were implicated in the mutagenic effects of cytoplasmic irradiation [W51]. Further, in human fibroblasts, the modulation of proteins involved in the p53-dependent pathway (Section II.B.2) following very low doses of alpha particles occurs in more cells than have been traversed by an alpha-particle track, apparently through cell-cell contact [A24]. The potential consequences of bystander effects are discussed further in Annex G, "*Biological effects at low radiation doses*".

202. In an attempt to link the delayed appearance of chromosomal aberrations to cancer-proneness, Ponnaiya et al. [P24] measured this form of instability in strains of mice differing in their sensitivity to radiation-induced mammary cancer. Strikingly, cells from the more sensitive strain (BALB/c) showed a marked increase in the frequency of chromatid aberrations after 16 population doublings, while the less sensitive strain (C57BL/6) showed no increase in aberrations over the control level.

203. Higher-than-average frequencies of specific types of chromosomal aberrations have been found after irradiation of

strains of mice that are prone to certain types of cancer. These aberrations are not necessarily the delayed effects of irradiation but nonetheless may indicate that radiation damage to specific parts of the genome contributes disproportionately to cancer induction. As an example, interstitial deletions of mouse chromosome 2 are consistently associated with radiation-induced acute myeloid leukaemia in several inbred strains of mice. The chromosome breakpoints involved in these chromosome 2 deletions are non-randomly distributed and can be detected at high frequency at early times following irradiation, suggesting that they may be an early event in leukaemogenesis [B54, B55, H43, H45, R28, T23]. Investigation of two mouse strains showing large differences in susceptibility to radiation-induced thymic lymphomas showed that at early times following x irradiation the incidence of specific chromosomal aberrations, especially trisomy 15, was much higher in the cancer-prone strain than in the resistant strain [C21]. These examples suggest that certain chromosomal regions may be more sensitive to radiation-induced genetic changes that are associated with cancer, although it is difficult to quantify this contribution precisely without a better understanding of the development of aberrant clones of cells in animal tissues.

204. In some mutation systems very high frequencies of mutation occur both spontaneously and after irradiation. With radiation induction, the amount of initial damage (e.g. the numbers of DNA double-strand breaks per unit dose) in the mutational target may not in these cases be sufficient to account for the numbers of mutations induced. While it is possible to consider that every type of damage (paragraph 15, Table 1) may sometimes lead to mutation, these examples of high-frequency mutation have usually been proposed to arise from a form of genetic instability. For instance, somatic mutation frequencies of coat colour in *Pink-eyed unstable* mice are linear with x-ray dose down to 10 mGy and occur at least 100 times more frequently per unit dose than germ-line mutations in the mouse specific-locus tests [S57]. The *Pink-eyed unstable* male mice have a reduction in the pigment of the coat and eyes, caused by gene duplication interfering with normal pigment production; deletion of this duplication is scored as the mutation, giving normal pigmentation seen as black spots on the grey coat.

205. The spontaneous frequency of mutation at tandem-repeat (VNTR or “minisatellite”) DNA sequences in the germ line of mice is also very high (1%-10% in offspring). The mechanism of mutation at these hypervariable loci does not appear to involve unequal exchange between homologous chromosomes [W30] but rather some form of complex gene conversion process [J10]. The incorporation of human tandem-repeat sequences into yeast cells, to study the mechanisms of variability, has confirmed that such sequences are destabilized in meiosis and that this process depends on the initiation of homologous recombination at a nearby DNA double-strand break [D23]. Irradiation of mouse spermatogonial stem cells by x rays increased the mutation frequency in offspring relative to controls [D10]. More extensive experiments at different stages of spermatogenesis using both single- and multi-locus probes showed no increase in mutation

frequency in post-meiotic spermatids but gave a doubling dose of 0.33 Gy for premeiotic spermatogonial and stem cells. The dose-response for mutation induction by 0.5-1 Gy x rays, combined for spermatogonial and stem cells, was linear [D19]. In contrast, in a separate study using one of the same probes at different germ-cell stages, the irradiation of spermatogonia gave a non-significant increase in mutation frequency, while the frequency in irradiated spermatids was significant. The increase in mutation frequency was not linear with dose, showing little increase above 1 Gy [F5, S19]. Similar data were reported for ²⁵²Cf irradiations (35% gamma rays, 65% neutrons): spermatids again showed the highest induced mutation frequency, and a single dose (1 Gy) to spermatogonia was significantly mutagenic. The RBE for these mutations with ²⁵²Cf was 5.9 for spermatogonia, 2.6 for spermatids, and 6.5 for spermatogonial stem cells [N15]. More work is required to reconcile the discrepancies that are seen for different germ-cell stages. Recently a striking result has been found from tandem-repeat mutation studies following irradiation of male mice with 0.5 Gy ²⁵²Cf neutrons, and the mating of these mice to unirradiated mice through two generations to yield second-generation (F₂) progeny. Remarkably, the F₂ mice showed an elevated frequency of mutation in the repeat sequences inherited from both male and female lines (6-fold and 3.5-fold, respectively), suggesting that genetic instability can be transmitted through the germline [D24]. The induced mutation frequency at these hypervariable sequences seems to be too high, by two orders of magnitude, for direct damage by radiation at the sites of mutation; it was proposed that some indirect mechanism of mutation induction is responsible [D19, F5, S19, D24].

206. Human spontaneous germ-line mutation frequencies at tandem-repeat loci can also be high (1%-7% per gamete). In a pilot study of children of survivors of the atomic bombings (mean gonad dose, generally to only one of the parents = 1.9 Sv) and matched controls, mutation frequency was measured at six tandem-repeat loci. The average mutation frequency was similar in the two groups, at 1.5% per gamete per locus for the exposed gametes and 2% for the unexposed gametes [K26]. A further study of the same children using a probe to detect multiple-repeat loci again showed no increase in mutation rate for the exposed group [S67]. However, a twofold increase in the frequency of mutation at tandem-repeat loci has been reported for children of parents resident in the Mogilev district of Belarus, which was heavily contaminated in the Chernobyl accident [D14]. Four tandem-repeat loci were tested, three of which showed increases by a factor of 1.7-2.0, while one locus (in which only one mutation was found) showed a reduction in the exposed group. The 79 families tested were compared to a control Caucasian population from the United Kingdom, which showed a similar overall distribution of repeat lengths at these loci. In a follow-up study using five additional tandem-repeat loci and including a further 48 families from Mogilev, the same general twofold increase was found when compared with the same (United Kingdom) control population [D1]. The mean dose to the exposed population was calculated to be 27.6±3.3 mSv from ¹³⁷Cs; some evidence for a dose-response relationship was obtained by dividing the exposed group into

those receiving >20 mSv (mutation rate = 0.024) and those receiving <20 mSv (rate = 0.018), compared with the control rate of 0.011. While it is clear that there is a general increase in the mutation rate for several different genomic sites in the exposed population, difficulties of interpretation exist because of the geographical disparity of the control group and the possibility that other environmental agents may be responsible for the increased mutation rate [S67]. As a direct test of the involvement of tandem-repeat loci in radiation-induced carcinogenesis, normal and tumour DNA from post-Chernobyl thyroid carcinomas was examined for mutations in three loci [N22]. Mutations were found in 3 of 17 tumours (18%), with one of these having mutations in all three loci, while none of 20 sporadic thyroid cancers from patients without a history of radiation exposure showed tandem-repeat mutations.

207. Recent studies of “hypermutation” in non-dividing cultures of bacteria placed under stress have also suggested that recombination processes are involved in the formation of small deletions at high frequency [R29]. It is suggested that when bacteria are starved of nutrients, for example, mutations arise at a very high rate in order to survive (if sufficient mutations occur, one of these may be sufficiently favourable to allow the cells to adapt to the prevailing conditions). Evidence supports a model in which cells enter a transient hypermutable state in which DNA double-strand breaks initiate homologous recombination activity (see Figure Id), priming error-prone DNA synthesis. It seems likely that the errors (mutations) arise as a result of a down-regulation of the mismatch repair system (paragraph 166) [H44]. Similarly, specific types of oxidative base damage (e.g. 8-oxoguanine, paragraph 24) may lead to small deletion mutations when mismatch correction is compromised [B56]. The significance of these findings is that the intrinsic mutability of cells is modifiable, and this principle may apply equally to many types of cell, perhaps including those involved in carcinogenesis [R30].

208. Other reported phenomena may also be connected to the induction of genetic instability following irradiation. One of these is coincident mutation: in cells selected for radiation-induced mutation at one genomic site, a high frequency of mutations occurs at other sites. Li et al. [L16] found two mutations at tandem-repeat loci among 50 x-ray-induced *TK* gene mutants, a frequency of 4×10^{-2} and far in excess of expectation based on current knowledge of radiation-induced gene mutation frequencies. No second-site mutations were found in 70 unirradiated clones. Perhaps the clearest example of coincident mutations was found in an extensive study of x-ray-induced mutation in the fungus *Neurospora crassa* [D5]. In a chromosomal region containing 21 genes, about 10% of the radiation-induced mutants had mutations in more than one gene following a single acute dose. However, a yeast cell study has revealed a possible mechanistic basis for non-targeted mutations that occur relatively close to the initial site of damage, without invoking genetic instability [S52]. A DNA double-strand break was placed at a single site in one yeast chromosome, using a site-specific endonuclease, and mutations were measured in a gene situated adjacent to the

break site. After the break had undergone recombination repair, it was found that the adjacent gene had sustained a 300-fold increase in the frequency of point mutations. This result suggests that the break repair process is error-prone, presumably because of a lack of fidelity in DNA repair synthesis, and extends over distances of several hundred DNA base pairs.

209. It is difficult to know how the reports of delayed genetic effects and instability following both low- and high-LET radiation may apply to humans. As indicated above (paragraphs 201-203), a high frequency of radiation-induced genetic or epigenetic changes may contribute to cancer incidence, at least in those instances where an accumulation of somatically-stable changes is required. However, it is still not clear whether genetic instability or a higher-than-normal mutation rate is a necessary for the development of tumours [T38]. In the case of inherited genetic alterations, where germ cells are the target, measurements of mutation frequency made at short intervals after irradiation could underestimate the induced mutation frequency. However, in the case of alpha-particle irradiation, mice injected with ^{239}Pu and subsequently mated to tester stocks (specific-locus method) over many weeks were not found to show increasingly high levels of mutation [R16]. It is, of course, possible that some potential increase in mutations may be balanced by selection against sperm carrying an increased load of genetic damage due to instability. The type of mutation induced may be important; if these are point mutations, they could have a greater chance of transmission and could lead to dominant genetic effects in offspring. The potential importance of genetic instability in carcinogenesis, especially at low radiation doses, is discussed further in Annex G, “*Biological effects at low radiation doses*”.

F. MUTATION FREQUENCIES AND CONSEQUENCES

210. Radiation-induced mutations are always measured against a background of spontaneously-occurring mutations. The mechanisms of spontaneous mutation are numerous; the chemical reactivity of DNA leads to instability, and there are inherent errors in replicating a very large molecule. Some of these mechanisms will overlap those of radiation damage; for example, oxidative damage from metabolic processes in aerobic organisms will give both base damage and strand breaks (paragraph 19). This indicates why many of the types of mutation that occur spontaneously are similar to those formed by ionizing radiation. The cell requires efficient repair processes to cope with endogenous damage; if unrepaired, the damage will lead to base-pair substitutions as well as some larger changes. The increase in mutant frequency found after exposure to ionizing radiation is likely to come from both the additional load of damage similar to that occurring spontaneously (such as DNA base damage and loss) and more complex radiation-induced damage that cannot be handled easily by the cell's battery of repair enzymes (Section I.B).

211. As was stressed in Section IV.B, an important influence on the observed frequency of mutation is the tolerance of the genome site to large changes. When large changes are tolerated, the spectrum of mutations induced by ionizing radiation shows that at least half (and commonly more than half) of the mutations measured shortly after irradiation are large deletions and rearrangements. This spectrum may change with time after irradiation, but as yet there are too few data to comment meaningfully on this possibility. Mutation frequencies have been measured for a few target genes, which have been chosen for their ease of use and especially for their presence in a hemizygous state. While these measurements

seem unlikely to represent the genome as a whole (see Section IV.E), they do give some idea of the variation in frequencies and the reasons for this variation. An interesting comparison can be made of the hamster *APRT* and *HPRT* genes; since both of these code for inessential enzymes of similar function (purine salvage) they should both detect all types of mutant. However, the *APRT* gene is thought to have an essential gene situated downstream, which limits the possibility of detecting very large deletions (paragraph 179). Table 4 shows that the frequencies of both spontaneous and radiation-induced *APRT* mutations are lower than those for *HPRT* by more than an order of magnitude.

Table 4
Mutant frequency and spectrum in two different genes of hamster cells

<i>Agent</i>	<i>Mutant frequency</i> (10^{-6} cells)	<i>Number analysed</i>	<i>Deletions/rearrangements</i> (%)
<i>APRT</i> (autosomal hemizygous) [M13]			
None	0.13	125	7.2
EMS ^a	430	48	0
Gamma rays	1.5-3.0 ^b	85	22.3
<i>HPRT</i> (X-linked hemizygous) [T20]			
None	6.2	44	18
EMS ^a	690	56	0
Gamma rays ^c	38.8	48	71

^a Ethylmethane sulphonate, an alkylating agent.

^b Doses 2.5 and 4 Gy.

^c Dose 5 Gy (both EMS and gamma-ray doses gave 20% cell survival).

212. Whether available mutation systems are sensitive enough to yield useful data on genetic damage in situations of practical importance is questionable. There has been some controversy over attempts to measure, for example, the frequency of *HPRT* gene mutations in the blood lymphocytes of radiotherapy technicians and patients. It was concluded that some of these studies are better at revealing the variables involved in measuring mutation at low doses than at giving reliable data on mutation frequencies (reviewed in [T22]). One study [N16], however, included molecular analysis of the mutations found in lymphocytes of radioimmune therapy patients and showed that a higher proportion with large genetic changes occurred than in controls, with some dose dependence for the fraction of mutants with large changes due to cumulative ¹³¹I activity. Studies of the survivors of the atomic bombings [H32, H33] reported a slight increase, about 10% per Gy, in *HPRT* mutant frequency as estimated dose increased. This frequency increase is considerably lower than found in freshly-irradiated lymphocytes, suggesting that *HPRT* mutants are selected against over the long time period involved. However, it has recently been found that a significant increase in *HPRT* mutant frequency could be detected in combined data from 142 liquidators involved in the Chernobyl accident (24% increase in the liquidator group relative to Russian controls) after adjustment for age and smoking [T49].

213. The problem of selection against mutant cells seems to be less severe in some more recently developed mutation assays. An assay based on the loss of one copy of the cell-surface marker glycoprotein A, encoded by the *GPA* gene, in human erythrocytes gave dose-dependent increases in mutant frequency in survivors of the atomic bombings, and the frequencies per unit dose were similar to those found in human cells irradiated with low-LET radiation under laboratory conditions [K1, K33, L44, L45]. In contrast to the *HPRT* data, there was a positive correlation between the frequencies of chromosomal aberrations and *GPA* mutants in exposed individuals; additionally, survivors with malignant solid tumours showed a significantly higher mutant frequency than those without cancer. This assay will measure chromosome loss (by, for example, non-disjunction) as well as mutation and recombination events in individuals already heterozygous for the *GPA* gene (about half of the human population), but it has the drawback that no molecular analysis of the mutations is possible (mature erythrocytes lack nuclei) [G24]. The *GPA* mutation system has also shown dose-dependent mutation induction in individuals exposed to doses of up to 6 Gy during or following the Chernobyl accident, and the slope of the dose response was very similar for these measurements and the *GPA* mutants measured in survivors of the atomic bombings (at about 25×10^{-6} mutants Gy^{-1}) [J18]. This result has received support from subsequent measurements of *GPA* mutation in liquidators [W53] or

people living in the vicinity of Chernobyl [L71] at the time of the accident. However, use of the *GPA* system to measure mutation in more than 700 Estonian and Latvian workers involved in the Chernobyl clean-up, with estimated median doses of about 100 mGy, did not detect a consistent increase in mutant frequency [B39]. The mutagenic effects of 5.3 MeV alpha particles (average LET = 140 keV μm^{-1}) in a human exposed to ^{232}Th over a 43-year period following thorotrast injection were measurable using the *GPA* gene assay [L11]. These data showed a fivefold increase in *GPA* mutants and correlated to large increases in chromosomal aberration frequency in lymphocytes from the same individual. Other data associated with the consequences of the Chernobyl accident are considered in the Annex J, "Exposures and effects of the Chernobyl accident".

214. Where enough measurements have been made, the average frequency of gene mutation induced by low-LET radiation is similar in cells from different somatic tissues and species. This can be illustrated by a plot of induced *HPRT* mutant frequency against surviving fraction (Figure VIIIa), where differences in intrinsic radiosensitivity are taken into account [T13, T14]. The plot also indicates that there is some consistency in the relationship between mutation and killing, suggesting that these responses derive from similar types of damage, a constant fraction of which is converted to mutations [T17]. It should be noted that this relationship does not say anything about the absolute mutant frequencies for different tissues or organisms; these may differ substantially, as seen, for example, in the response of different germ-cell stages in the mouse [S29].

215. The mutation frequency/survival plot can also be used to show that both the type of radiation and cellular parameters can influence the effectiveness of mutation induction. Thus, for *HPRT* mutation, densely ionizing radiation shows an increase in the effectiveness by a factor of no more than 2; i.e. the RBE for mutation is about twice that for cell killing (Figure VIIIb) [C22, T16, T18]. Similarly, x rays vary in effectiveness with the phase of the cell cycle; an increase in the effectiveness of mutation induction by factors of 2-3 is found relative to cell killing, especially in G_1/S phase [B14, J6]. It is tempting to speculate that the repair systems operational at this point in the cycle (see Section II.B.1) are more prone to recombinational errors.

216. Mutation induction is also subject to dose-rate effects; in general, the effectiveness of low-LET radiations is reduced by factors of 2-4 at low dose rates [T22]. However, it has also been found in specific conditions that the mutagenic effectiveness may remain the same or may increase at low dose rates relative to high dose rates [T22]. For example, if TK6 lymphoblastoid cells are exposed to low dose rates, no change in mutagenic effectiveness is found [K36], while their more radiation-resistant counterpart WIL2-NS (paragraph 185) shows an approximately twofold reduction in effectiveness [F11]. Lack of alteration in mutagenic effectiveness with dose rate has also been seen for radiation-sensitive (DNA-repair-deficient) cell lines [F12]. In addition, at dose rates of low-LET radiation of ≤ 0.5 mGy min^{-1} , rodent cell lines may

show no dose-rate effect [E10, E11, F13] or an increased (inverse) dose-rate effect [C44, C45]. Also, inverse dose-rate effects have been seen with low dose rates of high-LET radiation [K37, N18].





Figure VIII. Mutation-survival relationship for *HPRT* genes.

VIIIa: Similarity of induced mutant frequency in cell lines from human, mouse and hamster [T17];
VIIIb: The increased effectiveness of densely ionizing radiation (^{238}Pu alpha particles) compared to x rays in hamster cells [T18].

217. Some of these dose-rate data point to specific cellular processes influencing the response; in particular, it appears that the response is dependent on the capacity of cells to repair damage to DNA [T22]. It has long been thought that cellular repair capacity is responsible for the reduction in mutagenic effectiveness with dose rate, as seen for example in classical mouse germ-cell data [R6, R7]. Where little or no dose-rate effect is found, the repair capacity of the genetic region or of the whole cell is therefore likely to be impaired. This situation may also apply in some germ-cell stages; spermatozoa of both *Drosophila* and the mouse show no dose-rate effect [R6, T31]. The finding that very low dose rates give little relative effect, or an inverse effect, in some somatic cell lines may indicate that the full repair capacity of a cell requires a minimum level of insult (damage induced per unit

time) before it is brought to bear on that damage. However, it should be noted that chronic irradiation of mouse spermatogonial stem cells has shown no further reduction in specific-locus mutation frequency below 8 mGy min^{-1} with dose rates down to $7 \text{ } \mu\text{Gy min}^{-1}$ [R17].

G. SUMMARY

218. DNA damage caused by ionizing radiations leads to various types of mutation: the smaller mutations, such as base-pair substitutions, appear to result from damage to single bases, while larger changes, such as large deletions and rearrangements, probably arise from DNA double-strand breakage. The spectrum of radiation-induced mutations is dominated by the larger molecular changes if the genetic site assessed will tolerate these without lethality. In view of the propensity to induce large genetic changes, it may be supposed that the average radiation-induced mutation has greater consequences in terms of extensive alterations of the genome than the average spontaneous mutation or those induced by other agents. As yet no distinct mutational fingerprint has been identified for ionizing radiations, although there is a suggestion that the more complex types of molecular change may be overrepresented in the spectrum. High-LET radiations induce a higher mutation frequency per unit dose, but there are insufficient data to establish whether the mutation spectrum is different from that for low-LET radiations.

219. Attempts by cellular enzymes to repair DNA damage are intimately linked to mutation formation. The molecular mechanisms of radiation-induced mutation include illegitimate recombination, but where genes are heterozygous there is evidence that homologous recombination is involved (giving

loss of heterozygosity, commonly seen in some cancers as loss of tumour-suppressor gene function). Some forms of radiation-induced base damage will be repaired by mismatch repair pathways; loss of mismatch repair function can lead to hypermutability, which is linked to specific forms of cancer such as hereditary non-polyposis colon cancer. The involvement of repair processes in mutation formation suggests that the intrinsic mutability of cells is not fixed but will vary with their repair capacity; this concept may have important consequences for the process of carcinogenesis.

220. There are several reports of persistent genetic effects in cells and animals following irradiation, suggesting that genetic instability is induced. A possible explanation for this phenomenon is that stable oxidative reaction products persist and can continue to give genetic damage in subsequent cell generations, although other explanations are possible (paragraph 195). In some mutation systems very high frequencies of mutation (hypermutability) occur spontaneously and after irradiation; on the basis of target size, it is unlikely that these mutations are induced directly by radiation damage at the sites of mutation. Hypermutability also suggests that some form of genetic instability is induced by radiation treatment, but at present it is difficult to know how these data relate to responses such as cancer induction.

221. Methods have been established for measuring mutations with some accuracy in human somatic cells, such as blood lymphocytes, but most mutation systems lack the sensitivity to be used as indicators of genetic damage in cases of low-dose radiation exposure. Some success has been found in measuring mutant frequencies in cells of individuals exposed to relatively high doses of radiation, such as atomic bomb survivors and Chernobyl recovery operation workers.

CONCLUSIONS

222. DNA repair processes have evolved in all biological organisms to combat the deleterious effects of damage to their genetic material. Attempts to repair DNA damage also cause many types of mutation, through inability to properly restore the DNA sequence. Many of the genes involved in the repair of DNA damage in human cells have now been cloned, and their functional analysis has led to considerable progress in our understanding of the ways in which cells and organisms respond to radiation damage. Several different pathways of repair are required to cope with damage from ionizing radiation (Figure III), and the consequences of losing repair capacity can be drastic. The importance of the repair gene function has been revealed in particular by the development of methods to knock out specific genes in experimental animals. Using these methods, it has been found that the loss of repair gene activity is often lethal in early stages of development (Table 2). Where the knockout animals survive the loss of repair capacity, they are commonly very prone to cancer. These findings reinforce more limited studies with rare individuals in the human population, which have suggested for some time that radiosensitivity is linked to cancer

proneness. The converse finding, that mice lacking homologues of the human breast-cancer-predisposing (*BRCA*) genes are radiosensitive, is a further important illustration of this link. Loss of repair gene function is considered to lead to cancer proneness primarily through an increase in genetic instability, although the mechanistic details of this process remain to be elucidated. The precise contribution of loss of repair gene function to the risk of radiation-induced cancer in the population is unknown at present; if calculations are restricted to those rare individuals with recognized repair syndromes, then this contribution will be small, but it is already known that a more subtle variation in these genes occurs widely in the human population.

223. Cellular survival assays indicate a twofold variation in the response of individuals in the general population to acute radiation exposure. This variation may become three- or fourfold when irradiation is given at low dose rates. Specific groups of individuals may show a consistently elevated sensitivity; for example, individuals heterozygous for the ataxia-telangiectasia-mutated (*ATM*) gene, who constitute

about 1 percent of the population, show enhanced radiosensitivity especially when tested in a chromosome-damage assay. This finding has recently been reinforced by similar data with transgenic mice heterozygous for ATM deficiency. Strikingly, using the chromosome-damage assay, one study has shown that about 40 percent of breast cancer patients show a similar enhancement of radiation sensitivity. Recent evidence shows that the enhanced radiosensitivity of breast cancer patients has a genetic basis; it has been suggested that the *ATM* gene is involved in predisposition to breast cancer, but the evidence for this is controversial. It is possible that small reductions in the efficiency of any one of a number of genes involved in radiation response, due to subtle mutations or polymorphisms, will account for the existence of radiosensitive groups of individuals.

224. The analysis of DNA repair processes has revealed that they are part of a complex response system in our cells, which includes genes involved in recognizing and signalling the presence of damage and genes operating checkpoints to ensure that cells do not progress through the cell cycle if they carry DNA damage. The functional analysis of repair genes involved in radiation response has also revealed a link between radiosensitivity and immune dysfunction, because some of the gene products involved in the repair of radiation-induced DNA double-strand breaks also assemble functional immune genes. Thus, radiosensitivity in humans can arise from the loss of a broader spectrum of gene functions than was initially recognized.

225. A reduction in DNA repair capacity may have several consequences for the cellular radiation response, including alteration in the shapes of dose-response curves, the loss of low-dose-rate sparing, and a loss of relative effectiveness for high-LET radiations. It is probable that these consequences arise mainly from an alteration in the ability to repair complex forms of DNA damage, particularly those involving double-strand breaks, since this type of damage is extremely hazardous to the cell. DNA repair capacity is therefore an important component of radiation dose-response, and it is clear that it needs to be considered as a variable in modelling radiation action, especially when attempting to extrapolate to low dose exposure. However, before the effects of repair pathways can be modelled accurately, there is still considerably more to learn about the way in which different pathways contribute to the overall response to radiation. At present it is not known how cells control the use of different repair pathways in responding to damage, and in particular how a balance is achieved between correct (error-free) repair and repair leading to mutation induction. Additionally, more information is needed on the relationship between DNA repair and apoptosis in different cells and tissues to predict the outcome of radiation exposure.

226. There is experimental evidence that pre-irradiation with low doses (5-10 mGy) can increase the resistance of cells to a subsequent higher dose, and that a more sensitive cellular response to radiation exists at low doses than at higher doses. These observations have led to some controversy; they suggest that under some circumstances a resistance factor can be induced by low radiation doses, but the mechanistic basis of

the observations has proved difficult to establish. It has also been found in separate studies that a number of different genes and proteins are induced or repressed by radiation, although few of these appear to be involved directly in the repair of DNA damage. To date little of this characterization has been carried out in a systematic fashion, because it has been conditioned by the availability of cloned genes and proteins. However, new technology based on large-scale gene sequencing coupled to micro-arraying of sequences is beginning to revolutionize this type of study by enabling the assay of hundreds or thousands of gene products at one time. By examining the levels of many gene products from cells before and after irradiation, as well as with time elapsed following irradiation, it will be possible to see how the damage response is coordinated. These methods also permit a molecular description of the differences in levels of gene products present in normal and diseased tissues, including tumour tissue, facilitating an understanding of the mechanistic basis of cancer induction by agents such as ionizing radiation.

227. Many different types of DNA damage are caused by ionizing radiations, ranging from isolated single-strand damage at sites of single ionizations to complex DNA alterations at sites of clustered ionizations. The more complex forms of damage may be unique to the interaction of ionizing radiations with DNA, compared with damage occurring spontaneously or that caused by other DNA-damaging agents. Several attempts have been made to establish whether radiation damage can lead to a distinct fingerprint of genetic changes, but this has proved elusive. It is clear, for example, that the spectrum of gene mutation arising from radiation damage has differences from the spontaneous mutation spectrum, but the overlap in the two spectra is considerable. Again, new technologies based on fluorescent *in situ* hybridization, the use of reporter genes with fluorescent tags, and comparative genomic hybridization will allow a refined view of radiation-induced genetic changes and the hope of future distinctions.

228. Many genetic changes caused by radiation occur within a few hours of giving the dose, but there is experimental evidence to show that some forms of change may occur after much longer times and following many cell divisions. In addition, at some sites in the human genome, the frequency of radiation-induced genetic changes is much higher than would be expected based on direct damage to DNA by radiation tracks. These observations broaden our knowledge of the mechanisms of radiation action, but elucidation is required further before the consequences of these mechanisms for radiation risk can be understood.

229. It is clear that much more knowledge of the structure of the human genome, in particular the disposition of the genes within it and their responses to radiation, is required before it will be possible to predict the average frequency of mutation induced by a given dose of radiation. Additionally, the range of repair capacities present in the human population, including carriers of defective repair genes, will have to be considered to predict mutability on an individual basis. A more complete knowledge of these response mechanisms will allow greater accuracy in the prediction of radiation-induced carcinogenic and hereditary effects.

References

- A1 Alberts, B.M. Protein machines mediate the basic genetic processes. *Trends Genet.* 1: 26-30 (1985).
- A2 Arlett, C.F., S.A. Harcourt, A.R. Lehmann et al. Studies of a new case of Xeroderma pigmentosum (XP3BR) from complementation group G with cellular sensitivity to ionising radiation. *Carcinogenesis* 1: 745-751 (1980).
- A3 Afanasyev, V.N., B.A. Korol, Y.A. Mantsygin et al. Flow cytometry and biochemical analysis of DNA degradation characteristic of two types of cell death. *FEBS Lett.* 194: 347-350 (1986).
- A4 Anderson, C.W. DNA damage and the DNA-activated protein kinase. *Trends Biochem.* 18: 433-437 (1993).
- A5 Alt, F.W., E.M. Oltz, F. Young et al. VDJ recombination. *Immunol. Today* 13: 306-314 (1992).
- A6 Arlett, C.F. and S.A. Harcourt. Survey of radiosensitivity in a variety of human cell strains. *Cancer Res.* 40: 926-932 (1980).
- A7 Aurias, A. and B. Dutrillaux. Probable involvement of immunoglobulin superfamily gene in most recurrent chromosomal rearrangements from ataxia telangiectasia. *Hum. Genet.* 72: 210-214 (1986).
- A8 Aghamohammadi, S.Z., T. Morris, D.L. Stevens et al. Rapid screening for deletion mutations in the hprt gene using the polymerase chain reaction: X-ray and α -particle mutant spectra. *Mutat. Res.* 269: 1-7 (1992).
- A9 Aghamohammadi, S.Z. and J.R.K. Savage. A BrdU pulse double-labelling method for studying adaptive response. *Mutat. Res.* 251: 133-141 (1991).
- A10 Albering, H.J., J.J. Engelen, L. Koulischer et al. Indoor radon, an extrapulmonary genetic risk? *Lancet* 344: 750-751 (1994).
- A11 Amundson, S.A. and H.L. Liber. A comparison of induced mutation at homologous alleles of the tk locus in human cells. *Mutat. Res.* 247: 19-27 (1991).
- A12 Amundson, S.A. and H.L. Liber. A comparison of induced mutation at homologous alleles of the tk locus in human cells. II. Molecular analysis of mutants. *Mutat. Res.* 267: 89-95 (1992).
- A13 Amundson, S.A., F. Xia, K. Wolfson et al. Different cytotoxic and mutagenic responses induced by x-rays in two human lymphoblastoid cell lines derived from a single donor. *Mutat. Res.* 286: 233-241 (1993).
- A14 Arlett, C.F., M.H.L. Green, A. Priestley et al. Comparative human cellular radiosensitivity. I. The effect of SV40 transformation and immortalization on the gamma-irradiation survival of skin derived fibroblasts from normal individuals and from ataxia-telangiectasia patients and heterozygotes. *Int. J. Radiat. Biol.* 54: 911-928 (1988).
- A15 Azzam, E.I., G.P. Raaphorst and R.E.J. Michel. Radiation-induced adaptive response for protection against micronucleus formation and neoplastic transformation in C3H 10T $\frac{1}{2}$ mouse embryo cells. *Radiat. Res.* 138: S28-S31 (1994).
- A16 Aurias, A., J-L. Antonie, R. Assathiany et al. Radiation sensitivity of Bloom's syndrome lymphocytes during S and G₂ phase. *Cancer Genet. Cytogenet.* 16: 131-136 (1985).
- A17 Antoccia, A., R. Ricordy, P. Maraschio et al. Chromosomal sensitivity to clastogenic agents and cell cycle perturbations in Nijmegen breakage syndrome lymphoblastoid cell lines. *Int. J. Radiat. Biol.* 71: 41-49 (1997).
- A18 Athma, P., R. Rappaport and M. Swift. Molecular genotyping shows that ataxia-telangiectasia heterozygotes are predisposed to breast cancer. *Cancer Genet. Cytogenet.* 92: 130-134 (1996).
- A19 Araki, R., M. Itoh, K. Hamatani et al. Normal D-JH rearranged products of the IgH gene in SCID mouse bone marrow. *Int. Immuno.* 8: 1045-1053 (1996).
- A20 Araki, R., A. Fujimori, K. Hamatani et al. Nonsense mutation at Tyr-4046 in the DNA-dependent protein kinase catalytic subunit of severe combined immune deficiency mice. *Proc. Natl. Acad. Sci. U.S.A.* 94: 2438-2443 (1997).
- A21 Agarwal, M.L., W.R. Taylor, M.V. Chernov et al. The p53 network. *J. Biol. Chem.* 273: 1-4 (1998).
- A22 Appleby, J.M., J.B. Barber, E. Levine et al. Absence of mutations in the ATM gene in breast cancer patients with severe responses to radiotherapy. *Br. J. Cancer* 76: 1546-1549 (1997).
- A23 Andreev, S.G., I.K. Khvostunov, D.M. Spitkovsky et al. Clustering of DNA breaks in chromatin fibre: dependence on radiation quality. p. 133-136 in: *Microdosimetry: An Interdisciplinary Approach* (D.T. Goodhead et al., eds.). Roy. Soc. Chem., Cambridge, 1997.
- A24 Azzam, E.I., S.M. de Toledo, T. Gooding et al. Intercellular communication is involved in the bystander regulation of gene expression in human cells exposed to very low fluences of alpha particles. *Radiat. Res.* 150: 497-504 (1998).
- A25 Amundson, S.A., M. Bittner, Y. Chen et al. Fluorescent cDNA microarray hybridization reveals complexity and heterogeneity of cellular genotoxic stress responses. *Oncogene* 18: 3666-3672 (1999).
- A26 Amundson, S.A., K.T. Do and A.J. Fornace. Induction of stress genes by low doses of gamma rays. *Radiat. Res.* 152: 225-231 (1999).
- A27 Au, K.G., S. Clark, J.H. Miller et al. Escherichia coli mutY gene encodes an adenine glycosylase active on G-A mismatches. *Proc. Natl. Acad. Sci. U.S.A.* 86: 8877-8881 (1989).
- A28 Aspinwall, R., D.G. Rothwell, T. Roldan-Arjona et al. Cloning and characterization of a functional human homologue of Escherichia coli endonuclease III. *Proc. Natl. Acad. Sci. U.S.A.* 94: 109-114 (1997).
- B1 Bootsma, D. and J.H.J. Hoeijmakers. Engagement with transcription. *Nature* 363: 114-115 (1993).
- B2 Baer, R., A. Heppell, A.M.R. Taylor et al. The breakpoint of an inversion of chromosome 14 in a T-cell leukemia: sequences downstream of the immunoglobulin heavy chain locus are implicated in tumorigenesis. *Proc. Natl. Acad. Sci. U.S.A.* 84: 9069-9073 (1987).
- B3 Bates, P.R. and M.F. Lavin. Comparison of γ -radiation induced accumulation of ataxia telangiectasia and control cells in G₂ phase. *Mutat. Res.* 218: 165-170 (1989).
- B4 Barbi, G., J.M.J.C. Scheres, D. Schindler et al. Chromosome instability and x-ray hypersensitivity in a microcephalic and growth-retarded child. *Am. J. Med. Genet.* 40: 44-50 (1991).

- B5 Barnes, D.E., A.E. Tomkinson, A.R. Lehmann et al. Mutations in the DNA ligase-I gene of an individual with immunodeficiencies and cellular hypersensitivity to DNA-damaging agents. *Cell* 69: 495-503 (1992).
- B6 Beamish, H. and M.F. Lavin. Radiosensitivity in ataxia-telangiectasia - anomalies in radiation-induced cell cycle delay. *Int. J. Radiat. Biol.* 65: 175-184 (1994).
- B7 Bennett, C.B., A.L. Lewis, K.K. Baldwin et al. Lethality induced by a single site-specific double-strand break in a dispensable yeast plasmid. *Proc. Natl. Acad. Sci. U.S.A.* 90: 5613-5617 (1993).
- B8 Biedermann, K.A., J. Song, A.J. Giaccia et al. *scid* mutation in mice confers hypersensitivity to ionising radiation and a deficiency in DNA double-strand break repair. *Proc. Natl. Acad. Sci. U.S.A.* 88: 1394-1397 (1991).
- B9 Blocher, D., D. Sigut and M.A. Hannan. Fibroblasts from ataxia telangiectasia (AT) and AT heterozygotes show an enhanced level of residual DNA double-strand breaks after low dose-rate gamma-irradiation as assayed by pulsed field gel electrophoresis. *Int. J. Radiat. Biol.* 60: 791-802 (1991).
- B10 Boder, E. Ataxia-telangiectasia: an overview. p. 1-63 in: *Ataxia-telangiectasia: Genetics, Neuropathology and Immunology of a Degenerative Disease of Childhood* (R.A. Gatti and M. Swift, eds.). Liss, New York, 1985.
- B11 Bradley, W.E.C., A. Belouchi and K. Messing. The *aprt* heterozygote/hemizygote system for screening mutagenic agents allows detection of large deletions. *Mutat. Res.* 199: 131-138 (1988).
- B12 Bronner, C.E., S.M. Baker, P.T. Morrison et al. Mutation in the DNA mismatch repair gene homologue Hmlh1 is associated with hereditary non-polyposis colon cancer. *Nature* 368: 258-261 (1994).
- B13 Brenner, D.J. and R.K. Sachs. Chromosomal 'finger-prints' of prior exposure to densely ionizing radiation. *Radiat. Res.* 140: 134-142 (1994).
- B14 Burki, H.J. Ionizing radiation-induced 6-thioguanine resistant clones in synchronous CHO cells. *Radiat. Res.* 81: 76-84 (1980).
- B15 Badie, C., G. Iliakis, N. Foray et al. Defective repair of DNA double-strand breaks and chromosome damage in fibroblasts from a radiosensitive leukemia patient. *Cancer Res.* 55: 1232-1234 (1995).
- B16 Baker, S.M., C.E. Bronner, L. Zhang et al. Male mice defective in the DNA mismatch repair gene PMS2 exhibit abnormal chromosome synapsis in meiosis. *Cell* 82: 309-319 (1995).
- B17 Ban, S., S. Iida, A.A. Awa et al. Lethal and mutagenic effects of ²⁵²Cf radiation in cultured human cells. *Int. J. Radiat. Biol.* 52: 245-251 (1987).
- B18 Boothman, D.A., M. Meyers, E. Odegaard et al. Altered G1 checkpoint control determines adaptive survival responses to ionizing radiation. *Mutat. Res.* 358: 143-153 (1996).
- B19 Bao, C-Y., A-H. Ma, H.H. Evans et al. Molecular analysis of *HPRT* gene mutations induced by α - and X-radiation in human lymphoblastoid cells. *Mutat. Res.* 326: 1-15 (1995).
- B20 Barquinero, J.F., L. Barrios, M.R. Caballin et al. Occupational exposure to radiation induces an adaptive response in human lymphocytes. *Int. J. Radiat. Biol.* 67: 187-191 (1995).
- B21 Bauchinger, M., E. Schmid, H. Braselmann et al. Absence of adaptive response to low-level irradiation from tritiated thymidine and X-rays in lymphocytes of two individuals examined in serial experiments. *Mutat. Res.* 227: 103-107 (1989).
- B22 Bedford, J.S., J.B. Mitchell, H.G. Griggs et al. Radiation-induced cellular reproductive death and chromosome aberrations. *Radiat. Res.* 76: 573-586 (1978).
- B23 Bhattacharyya, N.P., A. Skandalis, A. Ganesh et al. Mutator phenotypes in human colorectal carcinoma cell lines. *Proc. Natl. Acad. Sci. U.S.A.* 91: 6319-6323 (1994).
- B24 Boice, J.D. and R.W. Miller. Risk of breast cancer in ataxia-telangiectasia. *Lancet* 326: 1357-1358 (1992).
- B25 Boothman, D.A., G. Majmudar and T. Johnson. Immediate X-ray-inducible responses from mammalian cells. *Radiat. Res.* 138: S44-S46 (1994).
- B26 Bosi, A. and G. Olivieri. Variability in the adaptive response to ionising radiations in humans. *Mutat. Res.* 211: 13-17 (1989).
- B27 Bracey, T.S., J.C. Miller, A. Preece et al. Gamma-radiation-induced apoptosis in human colorectal adenoma and carcinoma cell lines can occur in the absence of wild type p53. *Oncogene* 10: 2391-2396 (1995).
- B28 Branch, P., G. Aquilina, M. Bignami et al. Defective mismatch binding and a mutator phenotype in cells tolerant to DNA damage. *Nature* 362: 652-654 (1993).
- B29 Bridges, B.A., J. Cole, C.F. Arlett et al. Possible association between mutant frequency in peripheral lymphocytes and domestic radon concentrations. *Lancet* 337: 1187-1189 (1991).
- B30 Beamish, H., R. Williams, P. Chen et al. Defect in multiple cell cycle checkpoints in ataxia-telangiectasia post-irradiation. *J. Biol. Chem.* 271: 20486-20493 (1996).
- B31 Byrd, P.J., C.M. McConville, P. Cooper et al. Mutations revealed by sequencing the 5' half of the gene for ataxia telangiectasia. *Hum. Mol. Genet.* 5: 145-149 (1996).
- B32 Bentlos, N.J., D.A. Holzman, K.S. Keegan et al. The *Schizosaccharomyces pombe rad3* checkpoint gene. *EMBO J.* 15: 6641-6651 (1996).
- B33 Barlow, C., S. Hirotsumi, R. Paylor et al. *Atm*-deficient mice: a paradigm of ataxia telangiectasia. *Cell* 86: 159-171 (1996).
- B34 Benson, F.E., A. Stasiak and S.C. West. Purification and characterization of the human Rad51 protein, an analogue of *E. coli* RecA. *EMBO J.* 13: 5764-5771 (1994).
- B35 Boulton, S.J. and S.P. Jackson. *Saccharomyces cerevisiae* Ku70 potentiates illegitimate DNA double-strand break repair and serves as a barrier to error-prone DNA repair pathways. *EMBO J.* 15: 5093-5103 (1996).
- B36 Boulton, S.J. and S.P. Jackson. Identification of a *Saccharomyces cerevisiae* Ku80 homologue: roles in DNA double strand break rejoining and in telomeric maintenance. *Nucleic Acids Res.* 24: 4639-4648 (1996).
- B37 Blunt, T., D. Gell, M. Fox et al. Identification of a nonsense mutation in the carboxy-terminal region of DNA-dependent protein kinase catalytic subunit in the *scid* mouse. *Proc. Natl. Acad. Sci. U.S.A.* 93: 10285-10290 (1996).
- B38 Bosma, G.C., M. Fried, R.P. Custer et al. Evidence of functional lymphocytes in some (leaky) *scid* mice. *J. Exp. Med.* 167: 1016-1033 (1988).
- B39 Bigbee, W.L., R.H. Jensen, T. Veidebaum et al. Biodosimetry of Chernobyl cleanup workers from Estonia and Latvia using the glycophorin A *in vivo* somatic cell mutation assay. *Radiat. Res.* 147: 215-224 (1997).
- B40 Bernhard, E.J., R.J. Muschel, V.J. Bakanauskas et al. Reducing the radiation-induced G₂ delay causes HeLa cells to undergo apoptosis instead of mitotic death. *Int. J. Radiat. Biol.* 69: 575-584 (1996).
- B41 Boreham, D.R., K.L. Gale, S.R. Maves et al. Radiation-induced apoptosis in human lymphocytes: potential as a biological dosimeter. *Health Phys.* 71: 685-691 (1996).

- B42 Burnet, N.G., J. Nyman, I. Turesson et al. The relationship between cellular radiation sensitivity and tissue response may provide the basis for individualizing radiotherapy schedules. *Radiother. Oncol.* 33: 228-238 (1994).
- B43 Biard, D.S.F., M. Martin, Y. Le Rhun et al. Concomitant *p53* gene mutation and increased radiosensitivity in rat lung embryo epithelial cells during neoplastic development. *Cancer Res.* 54: 3361-3364 (1994).
- B44 Bech-Hansen, T., W.A. Blattner, B.M. Sell et al. Transmission of in vitro radioresistance in a cancer-prone family. *Lancet* i: 1335-1337 (1981).
- B45 Bartsch, H., M. Hollstein, R. Mustonen et al. Screening for putative radon-specific *p53* mutation hotspot in German uranium miners. *Lancet* 346: 121 (1995).
- B46 Beckman, K.B. and B.N. Ames. Oxidative decay of DNA. *J. Biol. Chem.* 272: 19633-19636 (1997).
- B47 Barlow, C., M. Liyanage, P.B. Moens et al. Partial rescue of prophase I defects of *Atm*-deficient mice by *p53* and *p21* null alleles. *Nature Genet.* 17: 462-466 (1997).
- B48 Baskaran, R., L.D. Wood, L.L. Whitaker et al. Ataxia-telangiectasia mutant protein activates c-Able tyrosine kinase in a response to ionizing radiation. *Nature* 387: 516-519 (1997).
- B49 Busse, P.M., S.K. Bose, R.W. Jones et al. The action of caffeine on X-irradiated HeLa cells. II Synergistic lethality. *Radiat. Res.* 71: 666-677 (1977).
- B50 Barlow, C., K.D. Brown, C-X. Deng et al. *Atm* selectively regulates distinct *p53*-dependent cell-cycle checkpoint and apoptotic pathways. *Nature Genet.* 17: 453-456 (1997).
- B51 Brown, J.M. Correspondence re: M.S. Meyn, Ataxia telangiectasia and cellular responses to DNA damage. *Cancer Res.* 57: 2313-2315 (1997).
- B52 Blocher, D. DNA double-strand break repair determines the RBE of alpha-particles. *Int. J. Radiat. Biol.* 54: 761-771 (1988).
- B53 Beer, J.Z. and I. Szumiel. Slow clones, reduced clonogenicity and intracolon recovery in X-irradiated L5178Y-S cell cultures. *Radiat. Environ. Biophys.* 33: 125-139 (1994).
- B54 Bouffler, S.D., G. Breckon and R. Cox. Chromosomal mechanisms in murine radiation acute myeloid leukaemogenesis. *Carcinogenesis* 17: 655-659 (1996).
- B55 Breckon, G., D. Papworth and R. Cox. Murine radiation myeloid leukaemogenesis: a possible role for radiation-sensitive sites on chromosome 2. *Genes Chromosomes Cancer* 3: 367-375 (1991).
- B56 Bridges, B.A. and A.R. Timms. Mutation in *Escherichia coli* under starvation conditions: a new pathway leading to small deletions in strains defective in mismatch correction. *EMBO J.* 16: 3349-3356 (1997).
- B57 Berdal, K.G., R.F. Johansen and E. Seeberg. Release of normal bases from intact DNA by a native DNA repair enzyme. *EMBO J.* 17: 363-367 (1998).
- B58 Bay, J.O., M. Grancho, D. Pernin et al. No evidence for constitutional ATM mutation in breast/gastric cancer families. *Int. J. Oncol.* 12: 1385-1390 (1998).
- B59 Banin, S., L. Moyal, S-Y. Shieh et al. Enhanced phosphorylation of *p53* by ATM in response to DNA damage. *Science* 281: 1674-1677 (1998).
- B60 Bouffler, S.D., C.J. Kemp, A. Balmain et al. Spontaneous and ionizing radiation-induced chromosomal abnormalities in *p53*-deficient mice. *Cancer Res.* 55: 3883-3889 (1995).
- B61 Barlow, C., M.A. Eckhaus, A.J. Schaffer et al. *Atm* haploinsufficiency results in increased sensitivity to sublethal doses of ionizing radiation in mice. *Nature Genetics* 21: 359-360 (1999).
- B62 Bullrich, F., D. Rasio, S. Kitada et al. ATM mutations in B-cell chronic lymphocytic leukemia. *Cancer Res.* 59: 24-27 (1999).
- B63 Barlow, C., P.A. Dennery, M.K. Shigenaga et al. Loss of the ataxia-telangiectasia gene product causes oxidative damage in target organs. *Proc. Natl. Acad. Sci. U.S.A.* 96: 9915-9919 (1999).
- B64 Blackstock, W.P. and M.P. Weir. Proteomics: quantitative and physical mapping of cellular proteins. *Trends Biotechnol.* 17: 121-127 (1999).
- B65 Bravard, A., C. Luccioni, E. Moustacchi et al. Contribution of antioxidant enzymes to the adaptive response to ionizing radiation of human lymphoblasts. *Int. J. Radiat. Biol.* 75: 639-645 (1999).
- B66 Boreham, D.R., J.A. Dolling, S.R. Maves et al. Dose-rate effects for apoptosis and micronucleus formation in gamma-irradiated human lymphocytes. *Radiat. Res.* 153: 579-86 (2000).
- C1 Caporossi, D., B. Porfirio, B. Nicoletti et al. Hypersensitivity of lymphoblastoid lines derived from ataxia telangiectasia patients to the induction of chromosomal aberrations by etoposide (VP-16). *Mutat. Res.* 290: 265-272 (1993).
- C2 Caldecott, K.W., C.K. McKeown, J.D. Tacker et al. An interaction between the mammalian DNA repair protein XRCC1 and DNA ligase III. *Mol. Cell. Biol.* 14: 68-76 (1994).
- C3 Chen, P., F.P. Imray and C. Kidson. Gene dosage and complementation analysis of ataxia telangiectasia lymphoblastoid lines assayed by induced chromosome aberrations. *Mutat. Res.* 129: 165-172 (1984).
- C4 Chessa, L., P. Petrinelli, A. Antonelli et al. Heterogeneity in ataxia-telangiectasia: classical phenotype associated with intermediate cellular radiosensitivity. *Am. J. Med. Genet.* 42: 741-746 (1992).
- C5 Chen, F., A. Nastasi, Z. Shen et al. Cell-cycle dependent expression of mammalian homologs of yeast DNA double-strand break repair genes *Rad51* and *Rad52*. *Mutat. Res.* 384: 205-211 (1997).
- C6 Clarke, A.R., C.A. Purdie, D.J. Harrison et al. Thymocyte apoptosis induced by *p53*-dependent and independent pathways. *Nature* 362: 849-852 (1993).
- C7 Cox, R., G.P. Hosking and J. Wilson. Ataxia telangiectasia: evaluation of radiosensitivity in cultured skin fibroblasts as a diagnostic test. *Arch. Dis. Childhood* 53: 386-390 (1978).
- C8 Cox, R. and W.K. Masson. Radiosensitivity in cultured human fibroblasts. *Int. J. Radiat. Biol.* 38: 575-576 (1980).
- C9 Cox, R., W.K. Masson, R.R. Weichselbaum et al. The repair of potentially lethal damage in X-irradiated cultures of normal and ataxia-telangiectasia human fibroblasts. *Int. J. Radiat. Biol.* 39: 357-365 (1981).
- C10 Cox, R. A cellular description of the repair defect in ataxia-telangiectasia. p.141-153 in: *Ataxia-telangiectasia - A Cellular and Molecular Link between Cancer, Neuropathology and Immune Deficiency* (B.A. Bridges and D.G. Harnden, eds.). Wiley, Chichester, 1982.
- C11 Cornforth, M.N. and J.S. Bedford. On the nature of a defect in cells from individuals with ataxia-telangiectasia. *Science* 227: 1589-1591 (1985).
- C12 Conley, M.E., N.B. Spinner, B.S. Emanuel et al. A chromosome breakage syndrome with profound immunodeficiency. *Blood* 67: 1251-1256 (1986).
- C13 Cox, R., P.G. Debenham, W.K. Masson et al. Ataxia-telangiectasia: a human mutation giving high-frequency misrepair of DNA double-stranded scissions. *Mol. Biol. Med.* 3: 229-244 (1986).

- C14 Cornforth, M.N. and J.S. Bedford. A quantitative comparison of potentially lethal damage repair and the rejoining of interphase chromosome breaks in low passage normal human fibroblasts. *Radiat. Res.* 111: 385-405 (1987).
- C15 Cole, J. and T.R. Skopek. Somatic mutant frequency, mutation rates and mutational spectra in the human population in vivo. *Mutat. Res.* 304: 33-105 (1994).
- C16 Collins, A.R. Mutant rodent cell lines sensitive to ultraviolet light, ionizing radiation and cross-linking agents: a comprehensive survey of genetic and biochemical characteristics. *Mutat. Res.* 293: 99-118 (1993).
- C17 Costa, N.D. and J. Thacker. Response of radiation-sensitive human cells to defined DNA breaks. *Int. J. Radiat. Biol.* 64: 523-529 (1993).
- C18 Curry, C.J.R., J. Tsai, H.T. Hutchinson et al. AT_{FRESNO}: a phenotype linking ataxia telangiectasia with the Nijmegen breakage syndrome. *Am. J. Hum. Genet.* 45: 270-275 (1989).
- C19 Cattanaach, B.M., M.D. Burtenshaw, C. Rasberry et al. Large deletions and other gross forms of chromosome imbalance compatible with viability and fertility in the mouse. *Nature Genet.* 3: 56-61 (1993).
- C20 Chang, W.P. and J.B. Little. Persistently elevated frequency of spontaneous mutations in progeny of CHO clones surviving X-irradiation: association with delayed reproductive death phenotype. *Mutat. Res.* 270: 191-199 (1992).
- C21 Chen, Y., E. Kubo, T. Sado et al. Cytogenetic analysis of thymocytes during early stages after irradiation in mice with different susceptibilities to radiation-induced lymphomagenesis. *J. Radiat. Res.* 37: 267-276 (1996).
- C22 Cox, R. and W.K. Masson. Mutation and inactivation of cultured mammalian cells exposed to beams of accelerated heavy ions. III. Human diploid fibroblasts. *Int. J. Radiat. Biol.* 36: 149-160 (1979).
- C23 Caelles, C., A. Helmberg and M. Karin. p53-dependent apoptosis in the absence of transcriptional activation of p53-target genes. *Nature* 370: 220-223 (1994).
- C24 Cai, L. and S.Z. Liu. Induction of cytogenetic adaptive response of somatic and germ cells in vivo and in vitro by low dose X-irradiation. *Int. J. Radiat. Biol.* 58: 187-194 (1990).
- C25 Cai, L. and P. Wang. Induction of a cytogenetic adaptive response in germ cells of irradiated mice with very low-dose rate of chronic gamma-irradiation and its biological influence on radiation-induced DNA or chromosomal damage and cell killing in their male offspring. *Mutagenesis* 10: 95-100 (1995).
- C26 Carrano, A.V. Chromosome aberrations and radiation-induced cell death. II. Predicted and observed cell survival. *Mutat. Res.* 17: 355-366 (1973).
- C27 Carter, S., M. Negrini, D.R. Gillum et al. Loss of heterozygosity at 11q22-23 in breast cancer. *Cancer Res.* 54: 6270-6274 (1994).
- C28 Cavazzana-Calvo, M., F. La Deist, G. de Saint Basil et al. Increased radiosensitivity of granulocyte macrophage colony-forming units and skin fibroblasts in human autosomal recessive severe combined immunodeficiency. *J. Clin. Invest.* 91: 1214-1218 (1993).
- C29 Chaudhry, M.A., Q. Jiang, M. Ricanati et al. Characterization of multilocus lesions in human cells exposed to X radiation and radon. *Radiat. Res.* 145: 31-38 (1996).
- C30 Chen, D.J., G.F. Strinste and N. Tokita. The genotoxicity of α -particles in human embryonic skin fibroblasts. *Radiat. Res.* 100: 321-327 (1984).
- C31 Chen, J.W., A.E. Tompkinson, W. Ramos et al. Mammalian DNA ligase. III: molecular cloning, chromosomal localization, and expression in spermatocytes undergoing meiotic recombination. *Mol. Cell. Biol.* 15: 5412-5422 (1995).
- C32 Chen, M., J. Quintans, Z. Fuks et al. Suppression of *Bcl-2* messenger RNA production may mediate apoptosis after ionizing radiation, tumor necrosis factor α , and ceramide. *Cancer Res.* 55: 991-994 (1995).
- C33 Cole, J., C.F. Arlett, M.H.L. Green et al. Comparative human cellular radiosensitivity. II. The survival following gamma-irradiation of unstimulated (G_0) T-lymphocytes, T-lymphocyte lines, lymphoblastoid cell lines and fibroblasts from normal donors, from ataxia-telangiectasia patients and from ataxia-telangiectasia heterozygotes. *Int. J. Radiat. Biol.* 54: 929-943 (1988).
- C34 Cole, J., M.H.L. Green, B.A. Bridges et al. Lack of evidence for an association between the frequency of mutants or translocations in circulating lymphocytes and exposure to radon gas in the home. *Radiat. Res.* 145: 61-69 (1996).
- C35 Cortes, F., I. Dominguez, J. Pinero et al. Adaptive response in human lymphocytes conditioned with hydrogen peroxide before irradiation with x-rays. *Mutagenesis* 5: 555-557 (1990).
- C36 Cortessis, V., S. Ingles, R. Millikan et al. Linkage analysis of DRD2, a marker linked to the ataxia-telangiectasia gene, in 64 families with premenopausal bilateral breast cancer. *Cancer Res.* 53: 5083-5086 (1993).
- C37 Cox, R. and W.K. Masson. Do radiation-induced thioguanine-resistant mutants of cultured mammalian cells arise by HGPRT gene mutation or X-chromosome rearrangement? *Nature* 276: 629-630 (1978).
- C38 Cimprich, K.A., T.B. Shin, C.T. Keith et al. cDNA cloning and gene mapping of a candidate human cell cycle checkpoint protein. *Proc. Natl. Acad. Sci. U.S.A.* 93: 2850-2855 (1996).
- C39 Caldecott, K.W., S. Aoufouchi, P. Johnson et al. XRCC1 polypeptide interacts with DNA polymerase β and possibly poly(ADP-ribose) polymerase, and DNA ligase III is a novel molecular 'nick-sensor' in vitro. *Nucleic Acids Res.* 24: 4387-4394 (1996).
- C40 Clutton, S.M., K.M.S. Townsend, C. Walker et al. Radiation-induced genomic instability and persisting oxidative stress in primary bone marrow cultures. *Carcinogenesis* 17: 1633-1639 (1996).
- C41 Cooper, P.K., T. Nospikel, S.G. Clarkson et al. Defective transcription-coupled repair of oxidative base damage in cockayne syndrome patients from XP group G. *Science* 275: 990-993 (1997).
- C42 Carr, A.M. and M.F. Hoekstra. The cellular responses to DNA damage. *Trends Cell Biol.* 5: 32-40 (1995).
- C43 Cox, L.S. and D.P. Lane. Tumour suppressors, kinases and clamps: how p53 regulates the cell cycle in response to DNA damage. *Bioessays* 17: 501-508 (1995).
- C44 Crompton, N.E.A., F. Zoelzer, E. Schneider et al. Increased mutant induction by very low dose-rate gamma-irradiation. *Naturwissenschaften* 72: S439 (1985).
- C45 Crompton, N.E.A., B. Barth and J. Kiefer. Inverse dose-rate effect for the induction of 6-thioguanine resistant mutants in Chinese hamster V79-S cells by ^{60}Co γ -rays. *Radiat. Res.* 124: 300-308 (1990).
- C46 Chernikova, S.V., V.Y. Gotlib and I.I. Pelevina. Influence of low dose irradiation on the radiosensitivity. *Radiat. Biol. Radioecol.* 33: 537-541 (1994).

- C47 Charlton, D.E., H. Nikjoo and J.L. Humm. Calculation of initial yields of single- and double-strand breaks in cell nuclei from electrons, protons and alpha particles. *Int. J. Radiat. Biol.* 56: 1-19 (1989).
- C48 Cliby, W.A., C.J. Roberts, K.A. Cimprich et al. Over-expression of a kinase-inactive ATR protein causes sensitivity to DNA-damaging agents and defects in cell cycle checkpoints. *EMBO J.* 17: 159-169 (1998).
- C49 Cleaver, J.E. and W.F. Morgan. Poly(ADP-ribose) polymerase: a perplexing participant in cellular responses to DNA breakage. *Mutat. Res.* 257: 1-18 (1991).
- C50 Custer, R.P., G.C. Bosma and M.J. Bosma. Severe combined immunodeficiency (SCID) in the mouse: pathology, reconstitution, neoplasms. *Am. J. Pathol.* 120: 464-477 (1985).
- C51 Critchlow, S.E., R.P. Bowater and S.P. Jackson. Mammalian DNA double-strand break repair protein XRCC4 interacts with DNA ligase IV. *Curr. Biol.* 7: 588-598 (1997).
- C52 Carroll, A.M. and M.J. Bosma. T-lymphocyte development in scid mice is arrested shortly after the initiation of T-cell receptor δ gene recombination. *Genes Dev.* 5: 1357-1366 (1991).
- C53 Coquerelle, T.M., K.F. Weibezahn and C. Lucke-Huhle. Rejoining of double-strand breaks in normal human and ataxia-telangiectasia fibroblasts after exposure to ^{60}Co gamma-rays, ^{241}Am alpha-particles or bleomycin. *Int. J. Radiat. Biol.* 51: 209-218 (1987).
- C54 Carney, J.P., R.S. Maser, H. Olivares et al. The hMre11/Rad50 protein complex and Nijmegen breakage syndrome: linkage of double-strand break repair to the cellular DNA damage response. *Cell* 93: 477-486 (1998).
- C55 Cadet, J., T. Douki and J-L. Ravenat. Artifacts associated with the measurement of oxidised DNA bases. *Environ. Health Perspect.* 105: 1034-1039 (1997).
- C56 Chaudhry, M.A. and M. Weinfeld. The action of *Escherichia coli* endonuclease III on multiply damaged sites in DNA. *J. Mol. Biol.* 249: 914-922 (1995).
- C57 Chaudhry, M.A. and M. Weinfeld. Reactivity of human apurinic/apyrimidinic endonuclease and *Escherichia coli* exonuclease III with bistranded abasic sites in DNA. *J. Biol. Chem.* 272: 15650-15655 (1997).
- C58 Canman, C.E., D-S. Lim, K.A. Cimprich et al. Activation of the ATM kinase by ionizing radiation and phosphorylation of p53. *Science* 281: 1677-1679 (1998).
- C59 Connor, F., D. Bertwistle, P.J. Mee et al. Tumorigenesis and a DNA repair defect in mice with a truncating *Brca2* mutation. *Nature Genet.* 17: 423-430 (1997).
- C60 Cunningham, R.P., S.M. Saporito, S.G. Spitzer et al. Endonuclease IV (*nfo*) mutant of *Escherichia coli*. *J. Bacteriol.* 168: 1120-1127 (1986).
- C61 Chester, N., F. Kuo, C. Kozak et al. Stage-specific apoptosis, developmental delay, and embryonic lethality in mice homozygous for a targeted disruption of the murine Bloom's syndrome gene. *Genes Dev.* 12: 3382-3393 (1998).
- C62 Clarke, A.A., N.J. Philpott, E.C. Godon-Smith et al. The sensitivity of fanconi anaemia group C cells to apoptosis induced by mitomycin C is due to oxygen radical generation not DNA crosslinking. *Br. J. Haematol.* 96: 240-247 (1997).
- C63 Cotes, F., I. Dominguez, M.J. Flores et al. Differences in the adaptive response to radiation damage in Go lymphocytes conditioned with hydrogen peroxide or low-dose x-rays. *Mutat. Res.* 311: 157-163 (1994).
- C64 Cedervall, B., R. Wong, N. Albright et al. Methods for the quantification of DNA double-strand breaks determined from the distribution of DNA fragment sizes measured by pulsed-field gel electrophoresis. *Radiat. Res.* 143: 8-16 (1995).
- C65 Chen, J., G.G. Birkholtz, P. Lindblom et al. The role of ataxia-telangiectasia heterozygotes in familial breast cancer. *Cancer Res.* 58: 1376-1379 (1998).
- C66 Cartwright, R., C.E. Tambini, P.J. Simpson et al. The XRCC2 repair gene from human and mouse encodes a novel member of the recA/RAD51 family. *Nucleic Acids Res.* 26: 3084-3089 (1998).
- C67 Cortez, D., Y. Wang, J. Qin et al. Requirement of ATM-dependent phosphorylation of Brca1 in the DNA damage response to double-strand breaks. *Science* 286: 1162-1166 (1999).
- C68 Chuang, Y.Y.E., Q. Chen et al. Radiation-induced mutations at the autosomal thymidine kinase locus are not elevated in p53-null cells. *Cancer Res.* 59: 3073-3076 (1999).
- C69 Celis, J.E., M. Ostergaard, N.A. Jensen et al. Human and mouse proteomics databases: novel resources in the protein universe. *FEBS Lett.* 430: 64-72 (1998).
- D1 Dubrova, Y.E., V.N. Nesterov, N.G. Krouchinsky et al. Further evidence for elevated human minisatellite mutation rate in Belarus eight years after the Chernobyl accident. *Mutat. Res.* 381: 267-278 (1997).
- D2 Deeley, J.O.T. and J.L. Moore. Nuclear lysate sedimentation measurements of peripheral blood lymphocytes from radiotherapy patients. *Int. J. Radiat. Biol.* 56: 963-973 (1989).
- D3 Donehower, L.A., M. Harvey, B.L. Slagle et al. Mice deficient for p53 are developmentally normal but susceptible to spontaneous tumours. *Nature* 356: 215-221 (1992).
- D4 Dulic, V., W.K. Kaufmann, S.J. Wilson et al. p53-dependent inhibition of cyclin-dependent kinase activities in human fibroblasts during radiation-induced G₁ arrest. *Cell* 76: 1013-1023 (1994).
- D5 de Serres, F.J. X-ray induced specific locus mutations in the *ad-3* region of two-component heterokaryons of *Neurospora crassa* II. More extensive genetic tests reveal an unexpectedly high frequency of multiple locus mutations. *Mutat. Res.* 210: 281-290 (1989).
- D6 da Costa, L.T., B. Liu, W. el-Deiry et al. Polymerase δ variants in RER colorectal tumours. *Nature Genet.* 9: 10-11 (1995).
- D7 Deschavanne, P.J., D. Debieu, B. Fertil et al. Re-evaluation of *in vitro* radiosensitivity of human fibroblasts of different genetic origins. *Int. J. Radiat. Biol.* 50: 279-293 (1986).
- D8 Dewey, W.C., S.C. Furman and H.H. Miller. Comparison of lethality and chromosomal damage induced by X-rays in synchronized Chinese hamster cells *in vitro*. *Radiat. Res.* 43: 561-581 (1970).
- D9 Dewind, N., M. Dekker, A. Berns et al. Inactivation of the mouse MSH2 gene results in mismatch repair deficiency, methylation tolerance, hyperrecombination and predisposition to cancer. *Cell* 82: 321-330 (1995).
- D10 Dubrova, Y.E., A.J. Jeffreys and A.M. Malashenko. Mouse minisatellite mutations induced by ionising radiation. *Nature Genet.* 5: 92-94 (1993).
- D11 Danska, J.S., D.P. Holland, S. Mariathasan et al. Biochemical and genetic defects in the DNA-dependent protein kinase in murine *scid* lymphocytes. *Mol. Cell. Biol.* 16: 5507-5517 (1996).

- D12 Danska, J.S., F. Pflumio, C. Williams et al. Rescue of T-cell specific V(D)J recombination in SCID mice by DNA-damaging agents. *Science* 266: 450-455 (1994).
- D13 Deshpande, A., E.H. Goodwin, S.M. Bailey et al. Alpha-particle-induced sister chromatid exchange in normal human lung fibroblasts: evidence for an extranuclear target. *Radiat. Res.* 145: 260-267 (1996).
- D14 Dubrova, Y.E., V.N. Nestorov, N.G. Krouchinsky et al. Human minisatellite mutation rate after the Chernobyl accident. *Nature* 380: 683-686 (1996).
- D15 Duchaud, E., A. Ridet, D. Stoppa-Lyonnet et al. Dereglated apoptosis in ataxia-telangiectasia - association with clinical stigmata and radiosensitivity. *Cancer Res.* 56: 1400-1404 (1996).
- D16 Dianov, G., A. Price and T. Lindahl. Generation of single-nucleotide repair patch following excision of uracil residues from DNA. *Mol. Cell. Biol.* 12: 1605-1612 (1992).
- D17 Dolganov, G.M., R.S. Maser, A. Novikov et al. Human Rad50 is physically associated with hMre11: identification of a conserved multiprotein complex implicated in recombinational DNA repair. *Mol. Cell. Biol.* 16: 4832-4841 (1996).
- D18 de Winter, J.P., Q. Waisfisz, M.A. Rooimans et al. The Fanconi anaemia group G gene *FANCG* is identical with *XRCC9*. *Nature Genet.* 20: 281-283 (1998).
- D19 Dubrova, Y.E., M. Plumb, J. Brown et al. Stage specificity, dose response, and doubling dose for mouse minisatellite germ-line mutation induced by acute radiation. *Proc. Natl. Acad. Sci. U.S.A.* 95: 6251-6255 (1998).
- D20 DeWeese, T.L., J.M. Shipman, N.A. Larrier et al. Mouse embryonic stem cells carrying one or two defective *Msh2* alleles respond abnormally to oxidative stress inflicted by low-level radiation. *Proc. Natl. Acad. Sci. U.S.A.* 95: 11915-11920 (1998).
- D21 di Fagagna, F.d'A., M.P. Hande W-M. Tong et al. Functions of poly(ADP-ribose) polymerase in controlling telomere length and chromosomal stability. *Nature Genet.* 23: 76-80 (1999).
- D22 Djuzenova, C.S., D. Schindler, H. Stopper et al. Identification of ataxia telangiectasia heterozygotes, a cancer-prone population, using the single-cell gel electrophoresis (Comet) assay. *Lab. Invest.* 79: 699-705 (1999).
- D23 Debrauwere, H., J. Buard, J. Tessier et al. Meiotic instability of human minisatellite *CEBI* in yeast requires DNA double-strand breaks. *Nature Genetics* 23: 367-371 (1999).
- D24 Dubrova, Y.E., M. Plumb, B. Gutierrez et al. Transgenerational mutation by radiation. *Nature* 405:37 (2000).
- E1 Ejima, Y. and M.S. Sasaki. Enhanced expression of X-ray and UV-induced chromosome aberrations by cytosine arabinoside in ataxia telangiectasia cells. *Mutat. Res.* 159: 117-123 (1986).
- E2 Elkind, M.M., A. Han and K.W. Volz. Radiation response of mammalian cells grown in culture. IV. Dose dependence of division delay and postirradiation growth of surviving and non-surviving Chinese hamster cells. *J. Natl. Cancer Inst.* 30: 705-721 (1963).
- E3 Evans, H.H., J. Mencl, M.-F. Horng et al. Locus specificity in the mutability of L5178Y mouse lymphoma cells: the role of multilocus lesions. *Proc. Natl. Acad. Sci. U.S.A.* 83: 4379-4383 (1986).
- E4 Easton, D.F. Cancer risks in A-T heterozygotes. *Int. J. Radiat. Biol.* 66: S177-S182 (1994).
- E5 Edwards, A.A., D.C. Lloyd and J.S. Prosser. Chromosome aberrations in human lymphocytes: a radiobiological review. p. 423-432 in: *Low Dose Radiation* (K.F. Baverstock and J.W. Stather, eds.). Taylor and Francis, London, 1989.
- E6 Ellis, N.A., J. Groden, T-Z. Ye et al. The Bloom's syndrome gene product is homologous to RecQ helicases. *Cell* 83: 655-666 (1995).
- E7 Elyan, S.A.G., C.M.L. West, S.A. Roberts et al. Use of low-dose-rate irradiation to measure the intrinsic radiosensitivity of human T-lymphocytes. *Int. J. Radiat. Biol.* 64: 375-383 (1993).
- E8 Eshleman, J.R., E.Z. Lang, G.K. Bowerfind et al. Increased mutation rate at the *HPRT* locus accompanies micro-satellite instability in colon cancer. *Oncogene* 10: 33-37 (1995).
- E9 Elson, A., Y. Wang, C.J. Daugherty et al. Pleiotropic defects in ataxia-telangiectasia protein-deficient mice. *Proc. Natl. Acad. Sci. U.S.A.* 93: 13084-13089 (1996).
- E10 Evans, H.H., M-F. Horng, J. Mencl et al. The influence of dose rate on the lethal and mutagenic effects of X-rays in proliferating L5178Y cells differing in radiation sensitivity. *Int. J. Radiat. Biol.* 47: 553-562 (1985).
- E11 Evans, H.H., M. Nielsen, J. Mencl et al. The effect of dose rate on X-radiation induced mutant frequency and the nature of DNA lesions in mouse lymphoma L5178Y cells. *Radiat. Res.* 122: 316-325 (1990).
- E12 Essers, J., R.W. Hendriks, S.M.A. Swagemakers et al. Disruption of the mouse *RAD54* reduces ionizing radiation resistance and homologous recombination. *Cell* 89: 195-204 (1997).
- E13 Engelward, B.P., G. Weeda, M.D. Wyatt et al. Base excision repair-deficient mice lacking the Aag alkyladenine DNA glycosylase. *Proc. Natl. Acad. Sci. U.S.A.* 94: 13087-13092 (1997).
- E14 Eilam, R., Y. Peter, A. Elson et al. Selective loss of dopaminergic nigro-striatal neurons in brains of *Atm*-deficient mice. *Proc. Natl. Acad. Sci. U.S.A.* 95: 12653-12656 (1998).
- E15 Enns, L., R.D.C. Barley, M.C. Paterson et al. Radiosensitivity in ataxia telangiectasia fibroblasts is not associated with deregulated apoptosis. *Radiat. Res.* 150: 11-16 (1998).
- F1 Friedberg, E.C. Deoxyribonucleic acid repair in the yeast *Saccharomyces cerevisiae*. *Microbiol. Rev.* 52: 70-102 (1988).
- F2 Fiorilli, M., A. Antonelli, G. Russo et al. Variant of ataxia-telangiectasia with low level radiosensitivity. *Hum. Genet.* 70: 274-277 (1985).
- F3 Frankenberg-Schwager, M. and D. Frankenberg. DNA double-strand breaks: their repair and relationship to cell killing in yeast. *Int. J. Radiat. Biol.* 58: 569-575 (1990).
- F4 Fritsche, M., C. Haessler and G. Brandner. Induction of nuclear accumulation of the tumor-suppressor protein p53 by DNA-damaging agents. *Oncogene* 8: 307-318 (1993).
- F5 Fan, Y.J., Z.W. Wang, S. Sadamoto et al. Dose response of radiation induction of germline mutation at a hypervariable mouse minisatellite locus. *Int. J. Radiat. Biol.* 68: 177-183 (1995).
- F6 Fomenko, L.A., I.K. Kozhanovskaia and A.I. Gaziev. Formation of micronuclei in bone marrow cells of chronically irradiated mice and subsequent acute exposure to gamma radiation. *Radiobiologia* 31: 701-715 (1991).
- F7 Fuhrman Conti, A.M., G. Francone, M. Volonte et al. Induction of 8-azaguanine resistant mutants in human cultured cells exposed to 31 MeV protons. *Int. J. Radiat. Biol.* 53: 467-476 (1988).

- F8 Fujimori, A., A. Tachibana and K. Tatsumi. Allelic losses at the *aprt* locus of human lymphoblastoid cells. *Mutat. Res.* 269: 55-62 (1992).
- F9 Fanconi Anaemia/Breast Cancer Consortium. Positional cloning of the Fanconi anaemia group A gene. *Nature Genet.* 14: 324-328 (1996).
- F10 Foray, N., C.F. Arlett and E.P. Malaise. Dose-rate effect on induction and repair rate of radiation-induced DNA double-strand breaks in a normal and an ataxia-telangiectasia human fibroblast cell line. *Biochimie* 77: 900-905 (1995).
- F11 Furuno-Fukushi, I., K. Tatsumi, M. Takahagi et al. Quantitative and qualitative effect of γ -ray dose rate on mutagenesis in human lymphoblastoid cells. *Int. J. Radiat. Biol.* 70: 209-217 (1996).
- F12 Furuno-Fukushi, I. and H. Matsudaira. Mutation induction by different dose rates of γ -rays in radiation-sensitive mutants of mouse leukaemia cells. *Radiat. Res.* 120: 370-374 (1989).
- F13 Furuno-Fukushi, I., A.M. Ueno and H. Matsudaira. Mutation induction by very low dose rate gamma-rays in cultured mouse leukemia cells L5178Y. *Radiat. Res.* 115: 273-280 (1988).
- F14 Fields, C., M.D. Adams, O. White et al. How many genes in the human genome? *Nature Genet.* 7: 345-346 (1994).
- F15 Friedberg, E.C., G.C. Walker and W. Siede. *DNA Repair and Mutagenesis*. ASM Press, Washington, 1995.
- F16 Foray, N., A. Priestley, G. Alsbeih et al. Hypersensitivity of ataxia telangiectasia fibroblasts to ionizing radiation is associated with a repair deficiency of DNA double-strand breaks. *Int. J. Radiat. Biol.* 72: 271-283 (1997).
- F17 Flaggs, G., A.W. Plug, K.M. Dunks et al. Atm-dependent interactions of a mammalian Chk1 homolog with meiotic chromosomes. *Curr. Biol.* 7: 977-986 (1997).
- F18 Foray, N., A. Priestley, G. Alsbeih et al. Hypersensitivity of ataxia telangiectasia fibroblasts to ionizing radiation is associated with a repair deficiency of DNA double-strand breaks. *Int. J. Radiat. Biol.* 72: 271-283 (1997).
- F19 Fritzell, J.A., L. Narayanan, S.M. Baker et al. Role of DNA mismatch repair in the cytotoxicity of ionizing radiation. *Cancer Res.* 57: 5143-5147 (1997).
- F20 Frank, K.M., J.M. Sekiguchi, K.J. Seidl et al. Late embryonic lethality and impaired V(D)J recombination in mice lacking DNA ligase IV. *Nature* 396: 173-177 (1998).
- F21 Fitzgerald, M.G., J.M. Bean, S.R. Hegde et al. Heterozygous ATM mutations do not contribute to early onset of breast cancer. *Nature Genet.* 15: 307-310 (1997).
- F22 Forrester, H.B., I.R. Radford et al. Selection and sequencing of interchromosomal rearrangements from gamma-irradiated normal human fibroblasts. *Int. J. Radiat. Biol.* 75: 543-551 (1999).
- F23 Forrester, H.B., R.F. Yeh et al. A dose response for radiation-induced intrachromosomal DNA rearrangements detected by inverse polymerase chain reaction. *Radiat. Res.* 152: 232-238 (1999).
- F24 Friedl, A.A., M. Kiechle, B. Fellerhof et al. Radiation-induced chromosome aberrations in *Saccharomyces cerevisiae*: influence of DNA repair pathways. *Genetics* 148: 975-988 (1998).
- F25 Fukumura, R., R. Araki, A. Fujimori et al. Murine cell line SX9 bearing a mutation in the *dna-pkcs* gene exhibits aberrant V(D)J recombination not only in the coding joint but also the signal joint. *J. Biol. Chem.* 273: 13058-13064 (1998).
- F26 Filippovitch, I.V., N.I. Sorokina, N. Robillard et al. Radiation-induced apoptosis in human tumor cell lines: adaptive response and split-dose effect. *Int. J. Cancer* 77: 76-81 (1998).
- G1 Game, J.C. DNA double-strand breaks and the *RAD50-RAD57* genes in *Saccharomyces*. *Cancer Biol.* 4: 73-83 (1993).
- G2 Gatti, R.A., I. Berkel, E. Boder et al. Localization of an ataxia-telangiectasia gene to chromosome 11q22-23. *Nature* 336: 577-580 (1988).
- G3 Ganesh, A., P. North and J. Thacker. Repair and misrepair of site-specific DNA double-strand breaks by human cell extracts. *Mutat. Res.* 299: 251-259 (1993).
- G4 Gentner, N.E. and D.P. Morrison. Determination of the proportion of persons in the population-at-large who exhibit abnormal sensitivity to ionizing radiation. p.253-262 in: *Low Dose Radiation* (K.F. Baverstock and J.W. Stather, eds.). Taylor and Francis, London, 1989.
- G5 Giaccia, A., R. Weinstein, J. Hu et al. Cell-cycle-dependent repair of double-strand breaks in a γ -ray sensitive Chinese hamster cell. *Somat. Cell Mol. Genet.* 11: 485-491 (1985).
- G6 Goodhead, D.T. Saturable repair models of radiation action in mammalian cells. *Radiat. Res.* 104: S58-S67 (1985).
- G7 Gottlieb, T.M. and S.P. Jackson. The DNA-dependent protein kinase: requirement for DNA ends and association with ku antigen. *Cell* 72: 131-142 (1990).
- G8 Gentil, A., J.B. Cabralneto, R. Mariagesamson et al. Mutagenicity of a unique apurinic/aprimidinic site in mammalian cells. *J. Mol. Biol.* 227: 981-984 (1992).
- G9 Gorgojo, L. and J.B. Little. Expression of lethal mutations in progeny of irradiated mammalian cells. *Int. J. Radiat. Biol.* 55: 619-630 (1989).
- G10 Grosovsky, A.J., J.G. de Boer, P.J. de Jong et al. Base substitutions, frameshifts and small deletions constitute ionising radiation-induced point mutations in mammalian cells. *Proc. Natl. Acad. Sci. U.S.A.* 85: 185-188 (1988).
- G11 Geara, F.B., L.J. Peters, K.K. Ang et al. Intrinsic radiosensitivity of normal human fibroblasts and lymphocytes after high- and low-dose-rate irradiation. *Cancer Res.* 52: 6348-6352 (1992).
- G12 Geara, F.B., L.J. Peters, K.K. Ang et al. Prospective comparison of *in vitro* normal cell radiosensitivity and normal tissue reactions in radiotherapy patients. *Int. J. Radiat. Oncol. Biol. Phys.* 27: 1173-1179 (1993).
- G13 Gentner, N.E. Increased definition of abnormal radiosensitivity using low dose rate testing. *Adv. Radiat. Biol.* 16: 293-302 (1992).
- G14 Gentner, N.E. Communication to the UNSCEAR Secretariat (1995).
- G15 Gerachi, J.P., P.D. Thrower, K.L. Jackson et al. The RBE of cyclotron fast neutrons for interphase death in rat thymocytes *in vitro*. *Int. J. Radiat. Biol.* 25: 403-405 (1974).
- G16 Gibbs, R.A., J. Camakaris, G.S. Hodgson et al. Molecular characterization of ^{125}I decay and X-ray induced HPRT mutants in CHO cells. *Int. J. Radiat. Biol.* 51: 193-199 (1987).
- G17 Green, M.H.L., C.F. Arlett, J. Cole et al. Comparative human cellular radiosensitivity: III. γ -radiation survival of cultured skin fibroblasts and resting T-lymphocytes from the peripheral blood of the same individual. *Int. J. Radiat. Biol.* 59: 749-765 (1991).
- G18 Griffiths, S.D., S.J. Marsden, E.G. Wright et al. Lethality and mutagenesis of B lymphocyte progenitor cells following exposure to α -particles and X-rays. *Int. J. Radiat. Biol.* 66: 197-205 (1994).

- G19 Gilad, S., R. Khosravi, D. Shkedy et al. Predominance of null mutations in ataxia telangiectasia. *Hum. Mol. Genet.* 5: 433-439 (1996).
- G20 German, J. Bloom syndrome: a mendelian prototype of somatic mutational disease. *Medicine* 72: 393-406 (1993).
- G21 Gschwend, M., O. Levrán, L. Kruglyak et al. A locus for Fanconi anaemia on 16q determined by homozygosity mapping. *Am. J. Hum. Genet.* 59: 377-384 (1996).
- G22 Greenwell, P.W., S.L. Kronmal, S.E. Porter et al. *TELL1*, a gene involved in controlling telomere length in *S. cerevisiae*, is homologous to the human ataxia telangiectasia gene. *Cell* 82: 823-829 (1995).
- G23 Guidos, C.J., C.J. Williams, I. Grandal et al. V(D)J recombination activates a p53-dependent DNA damage checkpoint in *scid* lymphocyte precursors. *Genes Dev.* 10: 2038-2054 (1996).
- G24 Grant, S.G. and W.L. Bigbee. In vivo somatic mutation and segregation at the human glycophorin A (*GPA*) locus: phenotypic variation encompassing both gene-specific and chromosomal mechanisms. *Mutat. Res.* 288: 163-172 (1993).
- G25 Golstein, P. Controlling cell death. *Science* 275: 1081-1082 (1997).
- G26 Griffiths, S.D., A.R. Clarke, L.E. Healy et al. Absence of p53 permits propagation of mutant cells following genotoxic damage. *Oncogene* 14: 523-531 (1997).
- G27 Goodhead, D.T. Track structure considerations in low dose and low dose rate effects of ionizing radiation. *Adv. Radiat. Biol.* 16: 7-44 (1992).
- G28 Goodhead, D.T. Initial events in the cellular effects of ionizing radiations: clustered damage in DNA. *Int. J. Radiat. Biol.* 65: 7-17 (1994).
- G29 Goodhead, D.T. and H. Nikjoo. Clustered damage in DNA: estimates from track-structure simulations. *Radiat. Res.* 148: 485-486 (1997).
- G30 Grawunder, U., M. Wilm, X. Wu et al. Activity of DNA ligase IV stimulated by complex formation with XRCC4 protein in mammalian cells. *Nature* 388: 492-495 (1997).
- G31 Gu, Y., S. Jin, Y. Gao et al. Ku70-deficient embryonic stem cells have increased ionizing radiosensitivity, defective DNA end-binding activity, and inability to support V(D)J recombination. *Proc. Natl. Acad. Sci. U.S.A.* 94: 8076-8081 (1997).
- G32 Goodhead, D.T., J. Thacker and R. Cox. Non-rejoining DNA breaks and cell inactivation. *Nature* 272: 379-380 (1978).
- G33 Gao, Y., Y. Sun, K.M. Frank et al. A critical role for DNA end-joining proteins in both lymphogenesis and neurogenesis. *Cell* 95: 891-902 (1998).
- G34 Gowen, L.C., A.V. Avrutskaya, A.M. Latour et al. *BRCA1* required for transcription coupled repair of oxidative DNA damage. *Science* 281: 1009-1012 (1998).
- G35 Gurley, K.E., K. Vo and C.J. Kemp. DNA double-strand breaks, p53, and apoptosis during lymphomagenesis in *scid/scid* mice. *Cancer Res.* 58: 3111-3115 (1998).
- G36 Goi, K., M. Takagi, S. Iwata et al. DNA damage-associated dysregulation of the cell cycle and apoptosis control in cells with germ-line p53 mutation. *Cancer Res.* 57: 1895-1902 (1997).
- G37 Gayther, S.A., W. Warren, S. Mazoyer et al. Germline mutations in the *BRCA1* gene in breast and ovarian cancer families provide evidence for a genotype-phenotype correlation. *Nature Genet.* 11: 428-433 (1995).
- G38 Gayther, S.A., J. Mangion, P. Russell et al. Variation of risks of breast and ovarian cancer associated with different mutations of the *BRCA2* gene. *Nature Genet.* 15: 103-105 (1997).
- G39 Goodhead, D.T., J. Thacker and R. Cox. Effects of radiations of different qualities on cells: molecular mechanisms of damage and repair. *Int. J. Radiat. Biol.* 63: 543-556 (1993).
- G40 Golub, T.R., D.K. Slonim, P. Tamayo et al. Molecular classification of cancer: class discovery and class prediction by gene expression monitoring. *Science* 286: 531-537 (1999).
- G41 Gibbs, P.E., W.G. McGregor, V.M. Maher et al. A human homolog of the *Saccharomyces cerevisiae* REV3 gene, which encodes the catalytic subunit of DNA polymerase zeta. *Proc. Natl. Acad. Sci. U.S.A.* 95: 6876-6880 (1998).
- H1 Hanawalt, P.C. Heterogeneity of DNA repair at the gene level. *Mutat. Res.* 247: 203-211 (1991).
- H2 Hozak, P., A.B. Hassan, D.A. Jackson et al. Visualization of replication factories attached to a nucleoskeleton. *Cell* 73: 361-373 (1993).
- H3 Hoeijmakers, J. Nucleotide excision repair II from yeast to mammals. *Trends Genet.* 9: 211-217 (1993).
- H4 Hutchinson, F. Chemical changes induced in DNA by ionising radiation. *Prog. Nucleic Acid Res. Mol. Biol.* 32: 115-154 (1985).
- H5 Harris, G., W.A. Cramp, J.C. Edwards et al. Radiosensitivity of peripheral blood lymphocytes in autoimmune disease. *Int. J. Radiat. Biol.* 47: 689-699 (1985).
- H6 Hannan, M.A., M. Kunhi, M. Einspenner et al. Post-irradiation DNA synthesis inhibition and G₂ phase delay in radiosensitive body cells from non-Hodgkin's lymphoma patients: an indication of cell cycle defects. *Mutat. Res.* 311: 265-276 (1994).
- H7 Henner, W.D. and M.E. Blaska. Hypersensitivity of cultured ataxia telangiectasia cells to etoposide. *J. Natl. Cancer Inst.* 76: 1107-1011 (1986).
- H8 Hendrickson, E.A., D.G. Schatz and D.T. Weaver. The *scid* gene encodes a transacting factor that mediates the rejoining event of Ig gene rearrangement. *Genes Dev.* 2: 817-829 (1988).
- H9 Hecht, F. and B.K. Hecht. Cancer in ataxia-telangiectasia patients. *Cytogenet. Cell Genet.* 46: 9-19 (1990).
- H10 Hendrickson, E.A., X-Q. Qin, E.A. Bump et al. A link between double-strand break related repair and V(D)J recombination: the *scid* mutation. *Proc. Natl. Acad. Sci. U.S.A.* 88: 4061-4065 (1991).
- H11 Hickson, I.D. and A.L. Harris. Mammalian DNA repair - use of mutants hypersensitive to cytotoxic agents. *Trends Genet.* 4: 101-106 (1988).
- H12 Hsieh, C.L., C.F. Arlett and M.R. Lieber. V(D)J recombination in ataxia telangiectasia, Bloom's syndrome and a DNA ligase I-associated immunodeficiency disorder. *J. Biol. Chem.* 268: 20105-20109 (1993).
- H13 Hain, J., R. Janussi and W. Burkart. Lack of adaptive response to low doses of ionising radiation in human lymphocytes from 5 different donors. *Mutat. Res.* 283: 137-144 (1992).
- H14 Hain, J., N.E.A. Crompton, W. Burkart et al. Caffeine release of radiation induced S and G₂ phase arrest in V79 hamster cells: increase of histone messenger RNA levels and p34^{cdc2} activation. *Cancer Res.* 53: 1507-1510 (1994).
- H15 Hampton, G.M., A. Mannermaa, R. Winqvist et al. Loss of heterozygosity in sporadic human breast carcinoma: a common region between 11q22 and 11q23. *Cancer Res.* 54: 4586-4589 (1994).
- H16 Harris, A.W. and J.W. Lowenthal. Cells of some cultured lymphoma lines are killed rapidly by X-rays and bleomycin. *Int. J. Radiat. Biol.* 42: 111-116 (1982).

- H17 Hartley, K.O., D. Gell, G.C.M. Smith et al. DNA-dependent protein kinase catalytic subunit: a relative of phosphatidylinositol 3-kinase and the ataxia telangiectasia gene product. *Cell* 82: 849-856 (1995).
- H18 Harvey, A.N. and J.R.K. Savage. A case of caffeine-mediated cancellation of mitotic delay without enhanced breakage in V79 cells. *Mutat. Res.* 304: 203-209 (1994).
- H19 Hedges, M.J. and S. Hornsey. The effect of X-rays and neutrons on lymphocyte death and transformation. *Int. J. Radiat. Biol.* 33: 291-300 (1978).
- H20 Hei, T.K., E.J. Hall and C.A. Waldren. Mutation induction and relative biological effectiveness of neutrons in mammalian cells. *Radiat. Res.* 115: 281-291 (1988).
- H21 Helzlsouer, K.J., E.L. Harris, R. Parshad et al. Familial clustering of breast cancer: possible interaction between DNA repair proficiency and radiation exposure in the development of breast cancer. *Int. J. Cancer* 64: 14-17 (1995).
- H22 Hemminki, A., P. Peltomaki and J.P. Mecklin. Loss of the wild type MLH1 gene is a feature of hereditary non-polyposis colorectal cancer. *Nature Genet.* 8: 405-410 (1994).
- H23 Hess, P., G. Aquilina, E. Dogliotti et al. Spontaneous mutations at aprt locus in a mammalian cell line defective in mismatch recognition. *Somat. Cell Mol. Genet.* 20: 409-421 (1994).
- H24 Higurashi, M. and P.E. Conen. In vitro chromosomal radiosensitivity in 'chromosomal breakage syndromes. *Cancer* 32: 380-383 (1973).
- H25 Hollstein, M., K. Rice, M.S. Greenblatt et al. Database of p53 gene somatic mutations in human tumors and cell lines. *Nucleic Acids Res.* 22: 3551-3555 (1994).
- H26 Huo, Y.K., Z. Wang, J-H. Hong et al. Radiosensitivity of ataxia-telangiectasia, X-linked agammaglobulinemia, and related syndromes using a modified colony survival assay. *Cancer Res.* 54: 2544-2547 (1994).
- H27 Hays, S.L., A.A. Firmenich and P. Berg. Complex formation in yeast double-strand break repair: participation of Rad51, Rad52, Rad55 and Rad57 proteins. *Proc. Natl. Acad. Sci. U.S.A.* 92: 6925-6929 (1995).
- H28 Hendry, J.H. and T-N. Jiang. Differential radiosensitizing effect of the *scid* mutation among tissues, studied using high and low dose rates; implications for prognostic indicators in radiotherapy. *Radiother. Oncol.* 33: 209-216 (1994).
- H29 Huang, L-C., K.C. Clarkin and G.M. Wahl. Sensitivity and selectivity of the DNA damage sensor responsible for activating p53-dependent G₁ arrest. *Proc. Natl. Acad. Sci. U.S.A.* 93: 4827-4832 (1996).
- H30 Holmberg, K., S. Falt, A. Johansson et al. Clonal chromosome aberrations and genomic instability in X-irradiated human T-lymphocyte cultures. *Mutat. Res.* 286: 321-330 (1993).
- H31 Holmberg, K., A.E. Meijer, G. Auer et al. Delayed chromosomal instability in human T-lymphocyte clones exposed to ionizing radiation. *Int. J. Radiat. Biol.* 68: 245-255 (1995).
- H32 Hakoda, M., M. Akiyama, S. Kyoizumi et al. Measurement of in vivo HGPRT-deficient mutant cell frequency using a modified method for cloning human peripheral blood T-lymphocytes. *Mutat. Res.* 197: 161-169 (1988).
- H33 Hirai, Y., Y. Kusunoki, S. Kyoizumi et al. Mutant frequency at the *HPRT* locus in peripheral blood T-lymphocytes of atomic bomb survivors. *Mutat. Res.* 329: 183-196 (1995).
- H34 Hannun, Y.A. Functions of ceramide in coordinating cellular responses to stress. *Science* 274: 1855-1859 (1996).
- H35 Haimovitz-Friedman, A., C. Kan, D. Ehleiter et al. Ionizing radiation acts on cellular membranes to generate ceramide and initiate apoptosis. *J. Exp. Med.* 180: 525-535 (1994).
- H36 Hopcia, K.L., Y.L. McCarey, F.C. Sylvester et al. Radiation-induced apoptosis in HL60 cells: oxygen effect, relationship between apoptosis and loss of clonogenicity and dependence of time of apoptosis on radiation dose. *Radiat. Res.* 145: 315-323 (1996).
- H37 Han, Z., D. Chatterjee, D.M. He et al. Evidence for a G₂ checkpoint in p53-independent apoptosis induction by X-irradiation. *Mol. Cell. Biol.* 15: 5849-5857 (1995).
- H38 Hendry, J.H., C.S. Potten, C. Chadwick et al. Cell death (apoptosis) in the mouse small intestine after low doses: effects of dose-rate, 14.7 MeV neutrons and 600 MeV (maximum energy) neutrons. *Int. J. Radiat. Biol.* 42: 611-620 (1982).
- H39 Hendry, J.H., C.S. Potten and A. Merritt. Apoptosis induced by high- and low-LET radiations. *Radiat. Environ. Biophys.* 34: 59-62 (1995).
- H40 Hakem, R., J.L. de la Pomba, C. Sirard et al. The tumour suppressor gene *Brcal* is required for embryonic cellular proliferation in the mouse. *Cell* 85: 1009-1023 (1996).
- H41 Hodgkins, P., P. O'Neill, D. Stevens et al. The severity of alpha-particle induced DNA damage is revealed by exposure to cell-free extracts. *Radiat. Res.* 146: 660-667 (1996).
- H42 Hei, T.K., L-J. Wu, S-X. Liu et al. Mutagenic effects of a single and an exact number of α particles in mammalian cells. *Proc. Natl. Acad. Sci. U.S.A.* 94: 3765-3770 (1997).
- H43 Hayata, I., M. Seki, K. Yoshida et al. Chromosomal aberrations observed in 52 mouse myeloid leukaemias. *Cancer Res.* 43: 367-373 (1983).
- H44 Harris, R.S., G. Feng, C. Thulin et al. Mismatch repair protein MutL becomes limiting during stationary-phase mutation. *Genes Dev.* 11: 2426-2437 (1997).
- H45 Hayata, I., T. Ichikawa and Y. Ichikawa. Specificity in chromosomal abnormalities in mouse bone marrows induced by the difference of the condition of irradiation. *Proc. Jpn. Acad.* 63: 289-292 (1987).
- H46 Haber, J.E. The many faces of Mre11. *Cell* 95: 583-586 (1998).
- H47 Hall, E.J., P.B. Schiff, G.E. Hanks et al. A preliminary report: frequency of A-T heterozygotes among prostate cancer patients with severe late responses to radiation therapy. *Cancer J. Sci. Am.* 4: 385-389 (1998).
- H48 Herzog, K-H., M.J. Chong, M. Kapsetaki et al. Requirement for Atm in ionizing radiation-induced cell death in the developing central nervous system. *Science* 280: 1089-1091 (1998).
- H49 Harrison, L., Z. Hatahet, A.A. Purmal et al. Multiply damaged sites in DNA: interactions with Escherichia coli endonucleases III and VIII. *Nucleic Acids Res.* 26: 932-941 (1998).
- H50 Heyer, J., K. Yang, M. Lipkin et al. Mouse models for colorectal cancer. *Oncogene* 18: 5325-5333 (1999).
- H51 Harada, Y-N., N. Shiomi, M. Koike et al. Postnatal growth failure, short life span, and early onset of cellular senescence and subsequent immortalization in mice lacking the xeroderma pigmentosum Group G gene. *Mol. Cell. Biol.* 19: 2366-2372 (1999).
- I1 Ikushima, T. Radio-adaptive response: characterization of a cytogenetic repair induced by low-level ionising radiation in cultured Chinese hamster cells. *Mutat. Res.* 227: 241-246 (1989).

- I2 Iliakis, G. The mutagenicity of α -particles in Ehrlich ascites tumor cells. *Radiat. Res.* 99: 52-58 (1984).
- I3 Iliakis, G. and M. Nuesse. Evidence that repair and expression of potentially lethal damage cause the variations in cell survival after X irradiation observed through the cell cycle in Ehrlich ascites tumour cells. *Radiat. Res.* 95: 87-107 (1983).
- I4 Ikushima, T., H. Aritomi and J. Morisita. Radioadaptive response: efficient repair of radiation-induced DNA damage in adapted cells. *Mutat. Res.* 358: 193-198 (1996).
- J1 Jaspers, N.G.J., R.D. Taalman and C. Baan. Patients with an inherited syndrome characterized by immunodeficiency, microcephaly and chromosomal instability: genetic relationship to ataxia-telangiectasia. *Am. J. Hum. Genet.* 42: 66-73 (1988).
- J2 Jaspers, N.G.J., R.A. Gatti, C. Baan et al. Genetic complementation analysis of ataxia telangiectasia and Nijmegen breakage syndrome: a survey of 50 patients. *Cytogenet. Cell Genet.* 49: 259-263 (1988).
- J3 Jeggo, P.A., J. Tesmer and D.J. Chen. Genetic analysis of ionising radiation sensitive mutants of cultured mammalian cell lines. *Mutat. Res.* 254: 125-133 (1990).
- J4 Jenner, T.J., C.M. Delara, P. O'Neill et al. Induction and rejoining of DNA double-strand breaks in V79-4 mammalian cells following gamma-irradiation and alpha-irradiation. *Int. J. Radiat. Biol.* 64: 265-273 (1993).
- J5 Jones, N.J., R. Cox and J. Thacker. Isolation and cross-sensitivity of X-ray sensitive mutants of V79-4 hamster cells. *Mutat. Res.* 183: 279-286 (1987).
- J6 Jostes, R.F., K.M. Bushnell and W.C. Dewey. X-ray induction of 8-azaguanine-resistant mutants in synchronous Chinese hamster ovary cells. *Radiat. Res.* 83: 146-161 (1980).
- J7 Joshi, G.P., W.J. Nelson, S.H. Revell et al. Discrimination of slow growth from non-survival among small colonies of diploid Syrian hamster cells after chromosome damage induced by a range of X-ray doses. *Int. J. Radiat. Biol.* 42: 283-296 (1982).
- J8 Jackson, S.P. Ataxia-telangiectasia at the crossroads. *Curr. Biol.* 5: 1210-1212 (1995).
- J9 Jayaraman, L. and C. Prives. Activation of p53 sequence-specific DNA binding by short single strands of DNA requires the p53 C-terminus. *Cell* 81: 1021-1029 (1995).
- J10 Jeffreys, A.J., K. Tamaki, A. MacLeod et al. Complex gene conversion events in germline mutation at human minisatellites. *Nature Genet.* 6: 136-145 (1994).
- J11 Jin, Y., T-A. Yie and A.M. Carothers. Non-random deletions at the dihydrofolate reductase locus of Chinese hamster ovary cells induced by α -particles simulating radon. *Carcinogenesis* 16: 1981-1991 (1995).
- J12 Jones, L.A., D. Scott, R. Cowan et al. Abnormal radiosensitivity of lymphocytes from breast cancer patients with excessive normal tissue damage after radiotherapy: chromosome aberrations after low-dose-rate irradiation. *Int. J. Radiat. Biol.* 67: 519-528 (1995).
- J13 Joshi, G.P., W.J. Nelson, S.H. Revell et al. X-ray induced chromosome damage in live mammalian cells, and improved measurements of its effects on their colony-forming ability. *Int. J. Radiat. Biol.* 41: 161-181 (1982).
- J14 Johnson, R.D. and L.S. Symington. Functional differences and interactions among the putative RacA homologs Rad51, Rad55 and Rad57. *Mol. Cell. Biol.* 15: 4843-4850 (1995).
- J15 Joenje, H., C. Mathew and E. Gluckman. Fanconi's anaemia research: current status and prospects. *Eur. J. Cancer* 31A: 268-272 (1995).
- J16 Joenje, H., J.R. Lo Ten Foe, A. Oostra et al. Classification of Fanconi anaemia patients by complementation analysis: evidence for a fifth genetic subgroup. *Blood* 86: 2156-2160 (1995).
- J17 Joenje, H., F. Arwert, A.W. Eriksson et al. Oxygen-dependence of chromosomal aberrations in Fanconi's anaemia. *Nature* 290: 142-143 (1981).
- J18 Jensen, R.H., R.G. Langlois, W.L. Bigbee et al. Elevated frequency of glycophorin A mutations in erythrocytes from Chernobyl accident victims. *Radiat. Res.* 141: 129-135 (1995).
- J19 Joenje, H., A.B. Oostra, M. Wijker et al. Evidence for at least eight Fanconi anemia genes. *Am. J. Hum. Genet.* 61: 940-944 (1997).
- J20 Jhappan, C., H.C. Morse, R.D. Fleischmann et al. DNA-PKcs: a T-cell tumour suppressor encoded at the mouse *scid* locus. *Nature Genet.* 17: 483-486 (1997).
- J21 Jongmans, W., M. Vuillaume, K. Chrzanowska et al. Nijmegen Breakage Syndrome cells fail to induce the p53-mediated DNA damage response following exposure to ionizing radiation. *Mol. Cell. Biol.* 17: 5016-5022 (1997).
- J22 Jongmans, W., M. Artuso, M. Vuillaume et al. The role of ataxia telangiectasia and the DNA-dependent protein kinase in the p53-mediated cellular response to ionising radiation. *Oncogene* 13: 1133-1138 (1996).
- J23 Johnson, R.E., M.T. Washington, S. Prakash et al. Bridging the gap: a family of novel DNA polymerases that replicate faulty DNA. *Proc. Natl. Acad. Sci. U.S.A.* 96: 12224-12226 (1999).
- J24 Johnson, R.E., C.M. Kondratyck, S. Prakash et al. HRAD30 mutations in the variant form of xeroderma pigmentosum. *Science* 285: 263-265 (1999).
- K1 Kyoizumi, S., M. Akiyama, J.B. Cologne et al. Somatic cell mutations at the glycophorin A locus in erythrocytes of atomic bomb survivors: implications for radiation carcinogenesis. *Radiat. Res.* 146: 43-52 (1996).
- K2 Kastan, M.B., O. Onyekwere, D. Sidransky et al. Participation of p53 protein in the cellular response to DNA damage. *Cancer Res.* 51: 6304-6311 (1991).
- K3 Kastan, M.B., Q.M. Zhan, W.S. Eldeiry et al. A mammalian cell cycle checkpoint pathway utilizing p53 and GADD45 is defective in ataxia-telangiectasia. *Cell* 71: 587-597 (1992).
- K4 Kaufmann, W.K. and S.J. Wilson. G₁ arrest and cell-cycle-dependent clastogenesis in UV-irradiated human fibroblasts. *Mutat. Res.* 314: 67-76 (1994).
- K5 Kroemer, G. The proto-oncogene Bcl-2 and its role in regulating apoptosis. *Nature Med.* 3: 614-620 (1997).
- K6 Khanna, K.K. and M.F. Lavin. Ionizing radiation and UV induction of p53 protein by different pathways in ataxia-telangiectasia cells. *Oncogene* 8: 3307-3312 (1993).
- K7 Kamiya, H., M. Suzuki, Y. Komatsu et al. An abasic site analogue activates a c-Ha-ras gene by a point mutation at modified and adjacent positions. *Nucleic Acids Res.* 20: 4409-4415 (1992).
- K8 Kadhim, M.A., D.A. Macdonald, D.T. Goodhead et al. Transmission of chromosomal instability after plutonium α -particle irradiation. *Nature* 355: 738-740 (1992).
- K9 Kadhim, M.A., S.A. Lorimore, M.D. Hepburn et al. Alpha-particle induced chromosomal instability in human bone marrow cells. *Lancet* 344: 987-988 (1994).
- K10 Kennedy, A.R., M. Fox, G. Murphy et al. Relationship between X-ray exposure and malignant transformation in C3H10T $\frac{1}{2}$ cells. *Proc. Natl. Acad. Sci. U.S.A.* 77: 7262-7266 (1980).

- K11 Kennedy, A.R. and J.B. Little. Evidence that a second event in X-ray induced oncogenic transformation *in vitro* occurs during cellular proliferation. *Radiat. Res.* 99: 228-248 (1984).
- K12 Kat, A., W.G. Thilly, W.H. Fang et al. An alkylation-tolerant, mutator cell line is deficient in strand-specific mismatch repair. *Proc. Natl. Acad. Sci. U.S.A.* 90: 6424-6428 (1993).
- K13 Kelsey, K.T., A. Memisoglu, D. Frenkel et al. Human lymphocytes exposed to low doses of X-rays are less susceptible to radiation-induced mutagenesis. *Mutat. Res.* 263: 197-201 (1991).
- K14 Kerr, J.F.R., A.H. Wyllie and A.R. Currie. Apoptosis: a basic biological phenomenon with wide-ranging implication in tissue kinetics. *Br. J. Cancer* 26: 239-257 (1972).
- K15 Keyse, S.M. The induction of gene expression in mammalian cells by radiation. *Cancer Biol.* 4: 119-128 (1993).
- K16 Khandogina, E.K., G.R. Mutovin, S.V. Zvereva et al. Adaptive response in irradiated human lymphocytes: radiobiological and genetical aspects. *Mutat. Res.* 251: 181-186 (1991).
- K17 Kidson, C., P. Chen and P. Imray. Ataxia-telangiectasia heterozygotes: dominant expression of ionising radiation-sensitive mutants. p. 141-153 in: *Ataxia-telangiectasia - A Cellular and Molecular Link between Cancer, Neuropathology and Immune Deficiency* (B.A. Bridges and D.G. Harnden, eds.). Wiley, Chichester, 1982.
- K18 Kimura, H., H. Higuchi, H. Iyehara-Ogawa et al. Sequence analysis of X-ray-induced mutations occurring in a cDNA of the human *hprt* gene integrated into mammalian chromosomal DNA. *Radiat. Res.* 134: 202-208 (1993).
- K19 Korsmeyer, S.J. Regulators of cell death. *Trends Genet.* 11: 101-105 (1995).
- K20 Kronenberg, A. and J.B. Little. Molecular characterization of thymidine kinase mutants of human cells induced by densely ionising radiation. *Mutat. Res.* 211: 215-224 (1989).
- K21 Kronenberg, A., S. Gauny, K. Criddle et al. Heavy ion mutagenesis: linear energy transfer effects and genetic linkage. *Radiat. Environ. Biophys.* 34: 73-78 (1995).
- K22 Kuller, L.H. and B. Modan. Risk of breast cancer in ataxia-telangiectasia. *Lancet* 326: 1357 (1992).
- K23 Kushihiro, J., N. Nakamura, S. Kyoizumi et al. Absence of correlations between radiosensitivities of human lymphocytes and skin fibroblasts from the same individuals. *Radiat. Res.* 122: 326-332 (1990).
- K24 Kubota, Y., R.A. Nash, A. Klungland et al. Reconstitution of DNA base-excision repair with purified human proteins: interaction between DNA polymerase β and the XRCC1 protein. *EMBO J.* 15: 6662-6670 (1996).
- K25 Komatsu, K., N. Kubota, M. Gallo et al. The *scid* factor on human chromosome 8 restores V(D)J recombination in addition to double-strand break repair. *Cancer Res.* 55: 1774-1779 (1995).
- K26 Kodaira, M., C. Satoh, K. Hiyamam et al. Lack of effects of atomic bomb radiation on genetic instability of tandem-repetitive elements in human germ cells. *Am. J. Hum. Genet.* 57: 1275-1283 (1995).
- K27 Keegan, K.S., D.A. Holzman, A.W. Plug et al. The ATR and ATM protein kinases associate with different sites along meiotically pairing chromosomes. *Genes Dev.* 10: 2423-2437 (1996).
- K28 Kanaar, R., C. Troelstra, S.M.A. Swagemakers et al. Human and mouse homologs of the *Saccharomyces cerevisiae* *RAD54* DNA repair gene: evidence for functional conservation. *Curr. Biol.* 6: 828-838 (1996).
- K29 Kubota, Y., R.A. Nash, A. Klungland et al. Reconstitution of DNA base excision repair with purified human proteins: interaction between DNA polymerase β and the XRCC1 protein. *EMBO J.* 15: 6662-6670 (1996).
- K30 Kuhn, E.M. Effects of X-irradiation in G₁ and G₂ on Bloom's syndrome and normal chromosomes. *Hum. Genet.* 54: 335-341 (1980).
- K31 Kataoka, M., M. Kawamura, K. Hamamoto et al. Radiation-induced oesophageal stricture in a case of Bloom's syndrome. *Clin. Oncol.* 1: 47-48 (1989).
- K32 Kadhim, M.A., C.A. Walker, M.A. Plumb et al. No association between p53 status and α -particle-induced chromosomal instability in human lymphoblastoid cells. *Int. J. Radiat. Biol.* 69: 167-174 (1996).
- K33 Kyoizumi, S., N. Nakamura, M. Hakoda et al. Detection of somatic mutations at the glycophorin A locus in erythrocytes of atomic bomb survivors using a single beam flow sorter. *Cancer Res.* 49: 581-588 (1989).
- K34 Kellerer, A.M., E.J. Hall, H.H. Rossi et al. RBE as a function of neutron energy. II. Statistical analysis. *Radiat. Res.* 65: 172-186 (1976).
- K35 Kruyt, F.A.E., L.M. Dijkmans, T.K. van den Berg et al. Fanconi anemia genes act to suppress a cross-linker-inducible p53-independent apoptosis pathway in lymphoblastoid cell lines. *Blood* 87: 938-948 (1996).
- K36 Koenig, F. and J. Kiefer. Lack of dose-rate effect for mutation induction by gamma-rays in human TK6 cells. *Int. J. Radiat. Biol.* 54: 891-897 (1988).
- K37 Kronenberg, A. and J.B. Little. Mutagenic properties of low doses of X-rays, fast neutrons and selected heavy ions in human cells. p. 554-559 in: *Low Dose Radiation* (K.F. Baverstock and J.W. Stather, eds.). Taylor and Francis, London, 1989.
- K38 Kolodner, R.D. Mismatch repair: mechanisms and relationship to cancer susceptibility. *Trends Biochem. Sci.* 20: 397-401 (1995).
- K39 Karow, J.K., R.K. Chakraverty and I.D. Hickson. The Bloom's syndrome gene product is a 3'-5' DNA helicase. *J. Biol. Chem.* 272: 30611-30614 (1997).
- K40 Kupfer, G.M., D. Naf, A. Suliman et al. The Fanconi anaemia proteins, FAA and FAC, interact to form a nuclear complex. *Nature Genet.* 17: 487-490 (1997).
- K41 Kooistra, R., K. Vreeken, J.B.M. Zonneveld et al. The *Drosophila melanogaster* *RAD54* homolog, *DmRAD54*, is involved in the repair of radiation damage and recombination. *Mol. Cell. Biol.* 17: 6097-6104 (1997).
- K42 Klungland, A. and T. Lindahl. Second pathway for completion of human DNA base excision repair: reconstitution with purified proteins and requirement for DNase IV (FEN1). *EMBO J.* 16: 3341-3348 (1997).
- K43 Khanna, K.K., K.E. Keating, S. Kozlov et al. ATM associates with and phosphorylates p53: mapping the region of interaction. *Nature Genet.* 20: 398-400 (1998).
- K44 Komarova, E.A., L. Diatchenko, O.W. Rokhlin et al. Stress-induced secretion of growth inhibitors: a novel tumour suppressor function of p53. *Oncogene* 17: 1089-1096 (1998).
- K45 Kobayashi, S., M. Nishimura, Y. Shimada et al. Increased sensitivity of *scid* heterozygous mice to ionizing radiation. *Int. J. Radiat. Biol.* 72: 537-545 (1997).
- K46 Klungland, A., M. Hoss, D. Gunz et al. Base excision repair of oxidative damage activated by XPG protein. *Mol. Cell.* 3: 33-42 (1999).
- K47 Kuhne, M., K. Rothkamm and M. Lobrich. No dose dependence of DNA double-strand break misrejoining following alpha-particle irradiation. *Int. J. Radiat. Biol.* (2000, in press).

- L1 Lehmann, A.R., B.A. Bridges, P.C. Hanawalt et al. Workshop on processing of DNA damage. *Mutat. Res.* 364: 245-270 (1996).
- L2 Lindahl, T. Instability and decay of the primary structure of DNA. *Nature* 362: 709-715 (1993).
- L3 Lindahl, T. and D.E. Barnes. Mammalian DNA ligases. *Annu. Rev. Biochem.* 61: 251-281 (1992).
- L4 Lavin, M.F., P. Le Poidevin and P. Bates. Enhanced levels of G₂ phase delay in ataxia-telangiectasia heterozygotes. *Cancer Genet. Cytogenet.* 60: 183-187 (1992).
- L5 Lavin, M.F., I. Bennett, J. Ramsay et al. Identification of a potentially radiosensitive subgroup among patients with breast cancer. *J. Natl. Cancer Inst.* 86: 1627-1634 (1994).
- L6 Lee, J.M. and A. Bernstein. p53 mutations increase resistance to ionizing radiation. *Proc. Natl. Acad. Sci. U.S.A.* 90: 5742-5746 (1993).
- L7 Little, J.B. Delayed initiation of DNA synthesis in irradiated human diploid cells. *Nature* 218: 1064-1065 (1968).
- L8 Little, J.B., J. Nove, L.C. Strong et al. Survival of human diploid fibroblasts from normal individuals after X-irradiation. *Int. J. Radiat. Biol.* 54: 899-910 (1988).
- L9 Lieber, M.R., J.E. Hesse, S. Lewis et al. The defect in murine severe combined immune deficiency: joining of signal sequences but not coding segments in V(D)J recombination. *Cell* 55: 7-16 (1988).
- L10 Lipkowitz, S., M-H. Stern and I.R. Kirsch. Hybrid T-cell receptor genes formed by interlocus recombination in normal and ataxia-telangiectasia lymphocytes. *J. Exp. Med.* 172: 409-418 (1990).
- L11 Littlefield, L.G., L.B. Travis, A.M. Sayer et al. Cumulative genetic damage in hematopoietic stem cells in a patient with a 40-year exposure to alpha particles emitted by thorium dioxide. *Radiat. Res.* 148: 135-144 (1997).
- L12 Lock, R.B. and W.E. Ross. Inhibition of p34^{cdc2} kinase activity by etoposide or irradiation as a mechanism of G₂ arrest in Chinese hamster ovary cells. *Cancer Res.* 50: 3761-3766 (1990).
- L13 Lowe, S.W., E.M. Schmitt, S.W. Smith et al. p53 is required for radiation-induced apoptosis in mouse thymocytes. *Nature* 362: 847-849 (1993).
- L14 Lu, X. and D.P. Lane. Differential induction of transcriptionally active p53 following UV or ionizing radiation - defects in chromosome instability syndromes? *Cell* 75: 765-778 (1993).
- L15 Leach, F.S., N.C. Nicolaides, N. Papadopoulos et al. Mutations of a mutS homolog in hereditary nonpolyposis colorectal cancer. *Cell* 75: 1215-1225 (1993).
- L16 Li, C-Y., D.W. Yandell and J.B. Little. Evidence for coincident mutations in human lymphoblast clones selected for functional loss of a thymidine kinase gene. *Mol. Carcinog.* 5: 270-277 (1992).
- L17 Lehnert, B.E. and E.H. Goodwin. Extracellular factor(s) following exposure to α particles can cause sister chromatid exchange in normal human cells. *Cancer Res.* 57: 2164-2171 (1997).
- L18 Little, J. DNA damage, recombination and mutation. p. 123-129 in: *Molecular Mechanisms in Radiation Mutagenesis and Carcinogenesis* (K.H. Chadwick, R. Cox, H.P. Leenhouts et al., eds.). European Commission, Brussels, 1994.
- L19 Lutze, L. and R.A. Winegar. pHAZE: a shuttle vector system for the detection and analysis of ionising radiation-induced mutations. *Mutat. Res.* 245: 305-310 (1990).
- L20 Lutze, L.H., R.A. Winegar, R. Jostes et al. Radon-induced deletions in human cells - role of nonhomologous strand rejoining. *Cancer Res.* 52: 5126-5129 (1992).
- L21 Lutze, L.H., J.E. Cleaver and R.A. Winegar. Factors affecting the frequency, size and location of ionising-radiation-induced deletions in human cells. p. 41-46 in: *Molecular Mechanisms in Radiation Mutagenesis and Carcinogenesis* (K.H. Chadwick, R. Cox, H.P. Leenhouts et al., eds.). European Commission, Brussels, 1994.
- L22 Lane, D.P. p53, guardian of the genome. *Nature* 358: 15-16 (1992).
- L23 Lambin, P., B. Marples, B. Fertil et al. Hypersensitivity of a human tumour cell line to very low radiation doses. *Int. J. Radiat. Biol.* 63: 639-650 (1993).
- L24 Lambin, P., B. Fertil, E.P. Malaise et al. Multiphasic survival curves for cells of human tumor cell lines: induced repair or hypersensitive subpopulation? *Radiat. Res.* 138: S32-S36 (1994).
- L25 Land, C.E. Risk of breast cancer in ataxia-telangiectasia. *Lancet* 326: 1359-1360 (1992).
- L26 Langley, R.E., S.G. Quartuccio, P.T. Keannealey et al. Effect of cell-cycle stage, dose rate and repair of sublethal damage on radiation-induced apoptosis in F9 teratocarcinoma cells. *Radiat. Res.* 144: 90-96 (1995).
- L27 Laval, F. Pretreatment with oxygen species increases the resistance of mammalian cells to hydrogen peroxide and γ -rays. *Mutat. Res.* 201: 73-79 (1988).
- L28 Lee, S., B. Elenbaas, A. Levine et al. p53 and its 14 kDa c-terminal domain recognize primary DNA damage in the form of insertion/deletion mismatches. *Cell* 81: 1013-1020 (1995).
- L29 Li, C-Y., D.W. Yandell and J.B. Little. Molecular mechanisms of spontaneous and induced loss of heterozygosity in human cells in vitro. *Somat. Cell Mol. Genet.* 18: 77-87 (1992).
- L30 Li, Z., T. Ottervel, Y. Gao et al. The *XRCC4* gene encodes a novel protein involved in DNA double-strand break repair and V(D)J recombination. *Cell* 83: 1079-1089 (1995).
- L31 Liber, H.L., P.K. LeMotte and J.B. Little. Toxicity and mutagenicity of X-rays and [¹²⁵I]dUrd or [³H]TdR incorporated in the DNA of human lymphoblast cells. *Mutat. Res.* 111: 387-404 (1983).
- L32 Ling, C.C., C.H. Chen and W.X.L. Li. Apoptosis induced at different dose rates: implications for the shoulder regions of the cell survival curves. *Radiother. Oncol.* 32: 129-136 (1994).
- L33 Ling, C.C., M. Guo, C.H. Chen et al. Radiation-induced apoptosis: effects of cell age and dose fractionation. *Cancer Res.* 55: 5207-5212 (1995).
- L34 Little, J.B. Changing views of cellular radiosensitivity. *Radiat. Res.* 140: 299-311 (1994).
- L35 Little, J.B. and J. Nove. Sensitivity of human diploid fibroblast cell strains from various genetic disorders to acute and protracted radiation exposure. *Radiat. Res.* 123: 87-92 (1990).
- L36 Liu, B., R.E. Parsons, S.R. Hamilton et al. hMSH2 mutations in hereditary nonpolyposis colorectal cancer kindreds. *Cancer Res.* 54: 4590-4594 (1994).
- L37 Lloyd, D.C., A.A. Edwards, A. Leonard et al. Frequencies of chromosomal aberrations induced in human blood lymphocytes by low doses of X-rays. *Int. J. Radiat. Biol.* 53: 49-55 (1988).
- L38 Lloyd, D.C., A.A. Edwards, A. Leonard et al. Chromosomal aberrations in human lymphocytes induced in vitro by very low doses of X-rays. *Int. J. Radiat. Biol.* 61: 335-343 (1992).
- L39 Loeffler, J.S., J.R. Harris, W.K. Dahlberg et al. *In vitro* radiosensitivity of human diploid fibroblasts derived from women with unusually sensitive clinical responses to definitive radiation therapy for breast cancer. *Radiat. Res.* 121: 227-231 (1990).

- L40 Lucke-Huhle, C., L. Hieber and R.D. Wegner. Caffeine-mediated release of alpha-irradiation induced G₂ arrest increase the yield of chromosome aberrations. *Int. J. Radiat. Biol.* 43: 123-132 (1983).
- L41 Lakin, N.D., P. Weber, T. Stankovic et al. Analysis of the ATM protein in wild-type and ataxia telangiectasia cells. *Oncogene* 13: 2707-2716 (1996).
- L42 Lim, D-S. and P. Hasty. A mutation in mouse *rad51* results in an early embryonic lethal that is suppressed by a mutation in *p53*. *Mol. Cell. Biol.* 16: 7133-7143 (1996).
- L43 Lo Ten Foe, J.R., M.A. Roomians, L. Bosnoyan-Collins et al. Expression cloning of a cDNA for the major Fanconi anaemia gene, *FAA*. *Nature Genet.* 14: 320-323 (1996).
- L44 Langlois, R.G., W.L. Bigbee, S. Kyoizumi et al. Evidence for increased somatic cell mutations at the glycophorin A locus in atomic bomb survivors. *Science* 236: 445-448 (1987).
- L45 Langlois, R.G., M. Akiyama, Y. Kusunoki et al. Analysis of somatic cell mutations at the glycophorin A locus in atomic bomb survivors: a comparative study of assay methods. *Radiat. Res.* 136: 111-117 (1993).
- L46 Liebertrau, W., A. Budde, A. Savoia et al. p53 activates Fanconi anemia group C gene expression. *Hum. Mol. Genet.* 6: 277-283 (1997).
- L47 Lonn, U., S. Lonn, U. Nylen et al. An abnormal profile of DNA replication intermediates in Bloom's syndrome. *Cancer Res.* 50: 3141-3145 (1990).
- L48 Lo, Y.M.D., S. Darby, L. Noakes et al. Screening for codon 249 p53 mutation in lung cancer associated with domestic radon exposure. *Lancet* 345: 60 (1995).
- L49 Levran, O., T. Erlich, N. Magdalena et al. Sequence variation in the Fanconi anemia gene *FAA*. *Proc. Natl. Acad. Sci. U.S.A.* 94: 13051-13056 (1997).
- L50 Liu, C.Y., A. Flesken-Nitikin, S. Li. et al. Inactivation of the mouse *Brcal* gene leads to failure in the morphogenesis of the egg cylinder in early post-implantational development. *Genes Dev.* 10: 1835-1843 (1996).
- L51 Lee, S.E., R.A. Mitchell, A. Cheng et al. Evidence for DNA-PK-dependent and independent DNA double-strand break repair pathways in mammalian cells as a function of the cell cycle. *Mol. Cell. Biol.* 17: 1425-1433 (1997).
- L52 Lobrich, M., B. Rydberg and P.K. Cooper. DNA double-strand breaks induced by high-energy neon and iron ions in human fibroblasts. II. Probing individual *NotI* fragments by hybridization. *Radiat. Res.* 139: 142-151 (1994).
- L53 Lobrich, M., P.K. Cooper and B. Rydberg. Non-random distribution of DNA double-strand breaks induced by particle irradiation. *Int. J. Radiat. Biol.* 70: 493-503 (1996).
- L54 Lucas, J.N., F.S. Hill, A.M. Chen et al. A rapid method for measuring pericentric inversions using fluorescence *in situ* hybridization (FISH). *Int. J. Radiat. Biol.* 71: 29-33 (1997).
- L55 Lucas, J.N. Cytogenetic signature for ionizing radiation. *Int. J. Radiat. Biol.* 73: 15-20 (1998).
- L56 Liu, N. and P.E. Bryant. Response of ataxia telangiectasia cells to restriction endonuclease induced DNA double-strand breaks 1. Cytogenetic characterization. *Mutagenesis* 8: 503-510 (1993).
- L57 Li, G., H. Ouyang, X. Li et al. *Ku70*: a candidate tumor suppressor gene for murine T cell lymphoma. *Mol. Cell* 2: 1-8 (1998).
- L58 Luo, L., F. Lu, S. Hart et al. Ataxia-telangiectasia and T-cell leukemias: no evidence for somatic *ATMi* mutation in sporadic T-ALL or for hypermethylation of the *ATM-NPAT/E14* bidirectional promoter in T-PLL. *Cancer Res.* 58: 2293-2297 (1998).
- L59 Liu, N., J.E. Lamerdin, J.D. Tucker et al. The human *XRCC9* gene corrects chromosomal instability and mutagen sensitivities in CHO UV40 cells. *Proc. Natl. Acad. Sci. U.S.A.* 94: 9232-9237 (1997).
- L60 Le, X.C., J.Z. Xing, J. Lee et al. Inducible repair of thymine glycol detected by an ultrasensitive assay for DNA damage. *Science* 280: 1066-1069 (1998).
- L61 Lloyd, D.C., P. Finnon, A.A. Edwards et al. Chromosome aberrations in syrian hamsters following very low radiation doses in vivo. *Mutat. Res.* 377: 63-68 (1997).
- L62 Lobrich, M., P.K. Cooper and B. Rydberg. Joining of correct and incorrect DNA ends at double-strand breaks produced by high-linear energy transfer radiation in human fibroblasts. *Radiat. Res.* 150: 619-626 (1998).
- L63 Lorimore, S.A., M.A. Kadhim, D.A. Pocock et al. Chromosomal instability in the descendants of unirradiated surviving cells after α -particle irradiation. *Proc. Natl. Acad. Sci. U.S.A.* 95: 5730-5733 (1998).
- L64 Lee, H., A.H. Trainer, L.S. Friedman et al. Mitotic checkpoint inactivation fosters transformation in cells lacking the breast cancer susceptibility gene, *Brca2*. *Mol. Cell* 4: 1-10 (1999).
- L65 Liu, N., J.E. Lamerdin, R.S. Tebbs et al. XRCC2 and XRCC3, new human RAD51-family members, promote chromosome stability and protect against DNA cross-links and other damages. *Mol. Cell* 1: 783-793 (1998).
- L66 Liao, M.J. and T. van Dyke. Critical role for *Atm* in suppressing V(D)J recombination-driven thymic lymphoma. *Genes Dev.* 13: 1246-1250 (1999).
- L67 Liao, W.C., A. Haimovitz-Friedman, R.S. Persaud et al. Ataxia telangiectasia-mutated gene product inhibits DNA damage-induced apoptosis via ceramide synthase. *J. Biol. Chem.* 274: 17908-17917 (1999).
- L68 Lindahl, T. Repair of intrinsic DNA lesions. *Mutat. Res.* 238: 305-311 (1990).
- L69 Lindahl, T. and R.D. Wood. Quality control by DNA repair. *Science* 286: 1897-1905 (1999).
- L70 Lehmann, A.R. Nucleotide excision repair and the link with transcription. *Trends Biochem. Sci.* 20: 402-405 (1995).
- L71 Livingston, G.K., R.H. Jensen, E.B. Silberstein et al. Radiobiological evaluation of immigrants from the vicinity of Chernobyl. *Int. J. Radiat. Biol.* 72: 703-713 (1997).
- L72 Lobrich, M., M. Kuhne, J. Wetzel et al. Joining of correct and incorrect DNA double-strand break ends in normal human and ataxia telangiectasia fibroblasts. *Genes, Chromosomes Cancer* 27: 59-68 (2000).
- M1 Michael, J.M., M.F. Lavin and D.J. Watters. Resistance to radiation-induced apoptosis in burkitt's lymphoma cells is associated with defective ceramide signaling. *Cancer Res.* 57: 3600-3605 (1997).
- M2 Mullenders, L.H.F., H. Vrieling, J. Venema et al. Hierarchies of DNA repair in mammalian cells: biological consequences. *Mutat. Res.* 250: 223-228 (1991).
- M3 Murray, A.W. Creative blocks - cell-cycle checkpoints and feedback controls. *Nature* 359: 599-604 (1992).
- M4 Malkin, D., F.P. Li, I.C. Strong et al. Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas and other neoplasms. *Science* 250: 1233-1238 (1990).
- M5 McKinnon, P.J. Ataxia-telangiectasia: an inherited disorder of ionising radiation sensitivity in man. *Hum. Genet.* 75: 197-208 (1987).
- M6 Morrison, C., G.C.M. Smith, L. Stingl et al. Genetic interaction between PARP and DNA-PK in V(D)J recombination and tumorigenesis. *Nature Genet.* 17: 479-482 (1997).

- M7 McWhir, J., J. Selfridge, D.J. Harrison et al. Mice with DNA repair gene (*ERCC1*) deficiency have elevated levels of p53, liver nuclear abnormalities and die before weaning. *Nature Genet.* 5: 217-223 (1993).
- M8 Meyn, M.S. High spontaneous intrachromosomal recombination rates in ataxia-telangiectasia. *Science* 260: 1327-1330 (1993).
- M9 Menissier de Murcia, J.M., C. Niedergang, C. Trucco et al. Requirement of poly(ADP-ribose) polymerase in recovery from DNA damage in mice and in cells. *Proc. Natl. Acad. Sci. U.S.A.* 94: 7303-7307 (1997).
- M10 Muris, D.F.R., O. Bezzubova, J.M. Buerstedde et al. Cloning of human and mouse genes homologous to RAD52: a yeast gene involved in DNA repair and recombination. *Mutat. Res.* 315: 295-305 (1994).
- M11 Meuth, M. The structure of mutation in mammalian cells. *Biochim. Biophys. Acta* 1032: 1-17 (1990).
- M12 Miles, C. and M. Meuth. DNA sequence determination of gamma-radiation-induced mutations of the hamster *aprt* locus. *Mutat. Res.* 227: 97-102 (1989).
- M13 Miles, C., G. Sargent, G. Phear et al. DNA sequence analysis of gamma radiation-induced deletions and insertions at the *aprt* locus of hamster cells. *Mol. Carcinog.* 3: 233-242 (1990).
- M14 Moore, M.M., A. Amtower, G.H.S. Strauss et al. Genotoxicity of gamma-irradiation in L5178Y mouse lymphoma cells. *Mutat. Res.* 174: 149-154 (1986).
- M15 Morris, T. and J. Thacker. Formation of large deletions by illegitimate recombination in the *HPRT* gene of primary human fibroblasts. *Proc. Natl. Acad. Sci. U.S.A.* 90: 1392-1396 (1993).
- M16 Morris, T., W. Masson, B. Singleton et al. Analysis of large deletions in the *HPRT* gene of primary human fibroblasts using the polymerase chain reaction. *Somat. Cell Mol. Genet.* 19: 9-19 (1993).
- M17 Marples, B. and M.C. Joiner. The response of Chinese hamster V79 cells to low radiation doses: evidence of enhanced sensitivity of the whole cell population. *Radiat. Res.* 133: 41-51 (1993).
- M18 Marples, B. and M.C. Joiner. The elimination of low dose hypersensitivity in Chinese hamster V79-379A cells by pretreatment with X-rays or hydrogen peroxide. *Radiat. Res.* 141: 160-169 (1995).
- M19 Marples, B., G.K.Y. Lam, H. Zhou et al. The response of Chinese hamster V79-379A cells exposed to negative pions: evidence that increased radioresistance is dependent on linear energy transfer. *Radiat. Res.* 138: S81-S84 (1994).
- M20 Marra, G. and C.R. Boland. Hereditary non-polyposis colorectal cancer: the syndrome, the genes, and historical perspectives. *J. Natl. Cancer Inst.* 87: 1114-1125 (1995).
- M21 McDonnell, T.J. and S.J. Korsmeyer. Progression from lymphoid hyperplasia to high-grade malignant lymphoma in mice transgenic for the t(14;18). *Nature* 349: 254-256 (1991).
- M22 Metting, N.F., S.T. Palayoor, R.M. Macklis et al. Induction of mutations by bismuth-212 α -particles at two genetic loci in human B-lymphoblasts. *Radiat. Res.* 132: 339-345 (1992).
- M23 Matsuoka, S., M. Huang and S.J. Elledge. Linkage of ATM to cell cycle regulation by the Chk2 protein kinase. *Science* 282: 1893-1897 (1998).
- M24 Meyers, M., R.A. Schea, A.E. Petrowski et al. Role of X-ray inducible genes and proteins in adaptive survival response. p. 263-270 in: *Low Dose Irradiation and Biological Defence Mechanisms* (T. Sugahara et al., eds.). Elsevier, 1992.
- M25 Miyazaki, N. and Y. Fujiwara. Mutagenic and lethal effects of [5-¹²⁵I]iodo-2'-deoxyuridine incorporated into DNA of mammalian cells and their RBEs. *Radiat. Res.* 88: 456-465 (1981).
- M26 Modan, B., A. Chetrit, E. Alfandary et al. Increased risk of breast cancer after low-dose irradiation. *Lancet* 1: 629-631 (1989).
- M27 Modrich, P. Mechanisms and biological effects of mismatch repair. *Annu. Rev. Genet.* 25: 229-253 (1991).
- M28 Maser, R.S., K.J. Mosen, B.E. Nelms et al. hMre11 and hRad50 nuclear foci are induced during the normal cellular response to DNA double-strand breaks. *Mol. Cell. Biol.* 17: 6087-6096 (1997).
- M29 Metcalfe, J.A., J. Parkhill, L. Campbell et al. Accelerated telomere shortening in ataxia telangiectasia. *Nature Genet.* 13: 350-353 (1996).
- M30 McConville, C.M., T. Stankovic, P.J. Byrd et al. Mutations associated with variant phenotypes in ataxia-telangiectasia. *Am. J. Hum. Genet.* 59: 320-330 (1996).
- M31 Morrow, D.W., D.A. Tagle, Y. Shiloh et al. *TELI*, an *S. cerevisiae* homolog of the human gene mutated in ataxia telangiectasia, is functionally related to the yeast checkpoint gene *MEC1*. *Cell* 82: 831-840 (1995).
- M32 Milne, G.T., S. Jin, K.B. Shannon et al. Mutations in two Ku homologs define a DNA end-joining repair pathway in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 16: 4189-4198 (1996).
- M33 Mason, R.M., J. Thacker and M.P. Fairman. The joining of non-complementary DNA double-strand breaks by mammalian extracts. *Nucleic Acids Res.* 24: 4946-4953 (1996).
- M34 Marder, B.A. and W.F. Morgan. Delayed chromosomal instability induced by DNA damage. *Mol. Cell. Biol.* 13: 6667-6677 (1993).
- M35 Martins, M.B., L. Sabatier, M. Ricoul et al. Specific chromosome instability induced by heavy ions: a step towards transformation of human fibroblasts? *Mutat. Res.* 285: 229-237 (1993).
- M36 Manti, L., M. Jamali, K.M. Prise et al. Genomic instability in Chinese hamster cells after exposure to X-rays or alpha particles of different mean linear energy transfer. *Radiat. Res.* 147: 22-28 (1997).
- M37 Morita, T., Y. Yoshimura, A. Yamamoto et al. A mouse homolog of the *Escherichia coli recA* and *Saccharomyces cerevisiae RAD51* genes. *Proc. Natl. Acad. Sci. U.S.A.* 90: 6577-6580 (1993).
- M38 Meyn, M.S. Ataxia-telangiectasia and cellular responses to DNA damage. *Cancer Res.* 55: 5991-6001 (1995).
- M39 Merritt, A.J., C.S. Potten, C.J. Kemp et al. The role of *p53* in spontaneous and radiation-induced apoptosis in the gastrointestinal tract of normal and *p53*-deficient mice. *Cancer Res.* 54: 614-617 (1994).
- M40 Mullaart, E., P.H.M. Lohman, F. Berends et al. DNA damage metabolism and aging. *Mutat. Res.* 237: 189-210 (1990).
- M41 Matsuura, S., C. Weemaes, D. Smeets et al. Genetic mapping using microcell-mediated chromosome transfer suggest a locus for Nijmegen breakage syndrome at chromosome 8q21-24. *Am. J. Hum. Genet.* 60: 1487-1494 (1997).
- M42 Mizuta, R., J.M. LaSalle, H-L. Cheng et al. RAB22 and RAB163/mouse BRCA2: proteins that specifically interact with the RAD51 protein. *Proc. Natl. Acad. Sci. U.S.A.* 94: 6927-6932 (1997).
- M43 Myung, K., D.M. He, S.E. Lee et al. KARP-1: a novel leucine zipper protein expressed from the Ku86 autoantigen locus is implicated in the control of DNA-dependent protein kinase activity. *EMBO J.* 16: 3172-3184 (1997).

- M44 Myung, K., C. Braastad, D.M. He et al. KARP-1 is induced by DNA damage in a p53- and ataxia telangiectasia mutated-dependent fashion. *Proc. Natl. Acad. Sci. U.S.A.* 95: 7664-7669 (1998).
- M45 Matsuura, S., H. Tauchi, A. Nakamura et al. Positional cloning of the gene for Nijmegen breakage syndrome. *Nature Genet.* 19: 179-181 (1998).
- M46 Marmorstein, L.Y., T. Ouchi and S.A.A. Aaronson. The *BRCA2* gene product functionally interacts with p53 and Rad51. *Proc. Natl. Acad. Sci. U.S.A.* 95: 13869-13874 (1998).
- M47 Mitchell, E.L. and D. Scott G2 chromosomal radiosensitivity in fibroblasts of ataxia-telangiectasia heterozygotes and a Li-Fraumeni syndrome patient with radiosensitive cells. *Int. J. Radiat. Biol.* 72: 435-438 (1997).
- M48 Morimatsu, M., G. Donoho and P. Hasty. Cells deleted for *BRCA2* COOH terminus exhibit hypersensitivity to gamma-radiation and premature senescence. *Cancer Res.* 58: 3441-3447 (1998).
- M49 Masutani, C., R. Kusumoto, A. Yamada et al. The XPV (xeroderma pigmentosum variant) gene encodes human DNA polymerase ϵ . *Nature* 399: 700-704 (1999).
- M50 Moynahan, M.E., J.W. Chiu, B.H. Koller et al. Brca1 controls homology-directed DNA repair. *Mol. Cell* 4: 511-518 (1999).
- M51 Maki, H. and M. Sekiguchi. MutT protein specifically hydrolyses a potent mutagenic substrate for DNA synthesis. *Nature* 355: 273-275 (1992).
- M52 Makedonov, G.P., L.V. Tshovrebova, S.V. Unzhakov et al. Radioadaptive response in lymphocytes from the children from polluted Bryansk regions after Chernobyl accident. *Radiat. Biol. Radioecol.* 37: 640-644 (1997).
- M53 Mohrenweiser, H.W. and I.M. Jones. Variation in DNA repair is a factor in cancer susceptibility: a paradigm for the promises and perils of individual and population risk estimation. *Mutat. Res.* 400: 15-24 (1998).
- M54 Martin, S.G., T. Laroche, N. Suka et al. Relocalization of telomeric Ku and SIR proteins in response to DNA strand breaks in yeast. *Cell* 97: 621-633 (1999).
- N1 Nagasawa, H., J.B. Robertson, C.S. Arundel et al. The effect of X-irradiation on the progression of mouse 10T $\frac{1}{2}$ cells released from density-inhibited cultures. *Radiat. Res.* 97: 537-545 (1984).
- N2 Nagasawa, H., S.A. Latt, M.E. Lalande et al. Effects of X-irradiation on cell-cycle progression, induction of chromosomal aberrations and cell killing in ataxia telangiectasia fibroblasts. *Mutat. Res.* 148: 71-82 (1985).
- N3 Nagasawa, H., K.H. Kraemer, Y. Shiloh et al. Detection of ataxia telangiectasia heterozygous cell lines by post-irradiation cumulative labelling index: measurements with coded samples. *Cancer Res.* 47: 398-402 (1987).
- N4 Nakamura, N., R. Sposto, J.-I. Kushiuro et al. Is inter-individual variation of cellular radiosensitivity real or artifactual? *Radiat. Res.* 125: 326-330 (1991).
- N5 Nelson, W.G. and M.B. Kastan. DNA strand breaks - the DNA template alterations that trigger p53-dependent DNA damage response pathways. *Mol. Cell. Biol.* 14: 1815-1823 (1994).
- N6 Nove, J., J.B. Little, P.J. Mayer et al. Hypersensitivity of cells from a new chromosomal-breakage syndrome to DNA-damaging agents. *Mutat. Res.* 163: 255-262 (1986).
- N7 North, P., A. Ganesh and J. Thacker. The rejoining of double-strand breaks in DNA by human cell extracts. *Nucleic Acids Res.* 18: 6205-6210 (1990).
- N8 Nagasawa, H. and J.B. Little. Induction of sister chromatid exchanges by extremely low doses of α -particles. *Cancer Res.* 52: 6394-6396 (1992).
- N9 Nicklas, J.A., T.C. Hunter, J.P. O'Neill et al. Fine structure mapping of the human *hprt* gene region on the X-chromosome (Xq26). *Am. J. Hum. Genet.* 47 (Suppl.): A193 (1990).
- N10 Nicolaidis, N.C., N. Papadopoulos, B. Liu et al. Mutations of two PMS homologues in hereditary nonpolyposis colon cancer. *Nature* 371: 75-80 (1994).
- N11 Nelson, S.L., C.R. Giver and A.J. Grosovsky. Spectrum of X-ray-induced mutations in the human *hprt*-gene. *Carcinogenesis* 15: 495-502 (1994).
- N12 Nishino, H., A. Knoll, V.L. Buettner et al. p53 wild-type and p53 nullizygous Big Blue transgenic mice have similar frequencies and patterns of observed mutation in liver, spleen and brain. *Oncogene* 11: 263-270 (1995).
- N14 Nacht, M., A. Strasser, Y.R. Chan et al. Mutations in *p53* and *SCID* genes cooperate in tumorigenesis. *Genes Dev.* 10: 2055-2066 (1996).
- N15 Niwa, O., Y.-J. Fan, M. Numoto et al. Induction of a germline mutation at a hypervariable mouse minisatellite locus by ²⁵²Cf radiation. *J. Radiat. Res.* 37: 217-224 (1996).
- N16 Nicklas, J.A., M.T. Falta, T.C. Hunter et al. Molecular analysis of *in vivo hprt* mutations in human T lymphocytes. V. Effects of total body irradiation secondary to radio-immunoglobulin therapy. *Mutagenesis* 5: 461-468 (1990).
- N17 Nelson, S.L., K.K. Parks and A.J. Grosovsky. Ionizing radiation signature mutations in human cell mutants induced by low-dose exposures. *Mutagenesis* 11: 275-279 (1996).
- N18 Nakamura, N. and S. Sawada. Reversed dose-rate effect and RBE of ²⁵²californium radiation in the induction of 6-thioguanine resistant mutations in mouse L5178Y cells. *Mutat. Res.* 201: 65-71 (1988).
- N19 Nikjoo, H., P. O'Neill, D.T. Goodhead et al. Computational modelling of low-energy electron-induced DNA damage by early physical and chemical events. *Int. J. Radiat. Biol.* 71: 467-483 (1997).
- N20 Nevaldine, B., J.A. Longo and P.J. Hahn. The scid defect results in a much slower repair of DNA double-strand breaks but not high levels of residual breaks. *Radiat. Res.* 147: 535-540 (1997).
- N21 Nussenzweig, A., K. Sokol, P. Burgman et al. Hypersensitivity of *Ku80*-deficient cell lines and mice to DNA damage: the effects of ionizing radiation on growth, survival, and development. *Proc. Natl. Acad. Sci. U.S.A.* 94: 13588-13593 (1997).
- N22 Nikiforov, Y.E., M. Nikiforova and J.A. Fagin. Prevalence of minisatellite and microsatellite instability in radiation-induced post-Chernobyl pediatric thyroid carcinomas. *Oncogene* 17: 1983-1988 (1998).
- N23 Nelson, J.R., C.W. Lawrence and D.C. Hinckle. Thymine-thymine dimer bypass by yeast DNA polymerase zeta. *Science* 272: 1646-1649 (1996).
- N24 Nelson, J.R., C.W. Lawrence and D.C. Hinckle. Deoxycytidyl transferase activity of yeast REV1 protein. *Nature* 382: 729-731 (1996).
- N25 Nagasawa, H. and J.B. Little. Unexpected sensitivity to the induction of mutations by very low doses of alpha-particle radiation: evidence for a bystander effect. *Radiat. Res.* 152: 552-557 (1999).
- N26 Naito, T., A. Matsuura and F. Ishikawa. Circular chromosome formation in a fission yeast mutant defective in *ATM* homologues. *Nature Genet.* 20: 203-206 (1998).
- N27 Nakane, H., S. Takeuchi, S. Yuba et al. High incidence of ultraviolet-B or chemical-carcinogen-induced skin tumours in mice lacking the xeroderma pigmentosum group A gene. *Nature* 377: 165-168 (1995).

- O1 Okada, S. Radiation-induced death. p. 247-307 in: Radiation Biochemistry, Volume 1 (K.I. Altman et al., eds.). Academic Press, New York, 1970.
- O2 Olivieri, G., J. Bodycote and S. Wolff. Adaptive response of human lymphocytes to low concentrations of radioactive thymidine. *Science* 223: 594-597 (1984).
- O3 Olive, P.L. The role of DNA single and double-strand breaks in cell killing by ionizing radiation. *Radiat. Res.* 150: S42-S51 (1998).
- O4 Oliver, F.J., J. Menissier-de Murcia and G. de Murcia. Poly(ADP-ribose) polymerase in the cellular response to DNA damage, apoptosis, and disease. *Am. J. Hum. Genet.* 64: 1282-1288 (1999).
- O5 Osman, F. and S. Subramani. Double-strand break-induced recombination in eukaryotes. *Prog. Nucleic Acid Res. Mol. Biol.* 58: 263-299 (1998).
- P1 Perrino, F.W. and L.A. Loeb. Animal cell DNA polymerases in DNA repair. *Mutat. Res.* 236: 289-300 (1990).
- P2 Parshad, R., K.K. Sanford and G.M. Jones. Chromatid damage after G₂ phase X-irradiation of cells from cancer-prone individuals implicates deficiency in DNA repair. *Proc. Natl. Acad. Sci. U.S.A.* 80: 5612-5616 (1983).
- P3 Parshad, R., K.K. Sanford and G.M. Jones. Chromosomal radiosensitivity during the G₂ cell-cycle period of skin fibroblasts from individuals with familial cancer. *Proc. Natl. Acad. Sci. U.S.A.* 82: 5400-5403 (1985).
- P4 Painter, R.B. Inhibition of mammalian cell DNA synthesis by ionising radiation. *Int. J. Radiat. Biol.* 49: 771-781 (1986).
- P5 Pandita, T.K. and W.N. Hittelman. The contribution of DNA and chromosome repair deficiencies to the radiosensitivity of ataxia telangiectasia. *Radiat. Res.* 131: 214-223 (1992).
- P6 Pergola, F., M.Z. Zdzienicka and M.R. Lieber. V(D)J recombination in mammalian cell mutants defective in DNA double-strand break repair. *Mol. Cell. Biol.* 13: 3464-3471 (1993).
- P7 Plowman, P.N., B.A. Bridges, C.F. Arlett et al. An instance of clinical radiation morbidity and cellular radiosensitivity, not associated with ataxia-telangiectasia. *Br. J. Radiol.* 63: 624-628 (1990).
- P8 Pippard, E.C., A.J. Hall, D.J.P. Barker et al. Cancer in homozygotes and heterozygotes of ataxia-telangiectasia and xeroderma pigmentosum in Britain. *Cancer Res.* 48: 2929-2932 (1988).
- P9 Padovani, L., M. Appolloni, P. Anzidei et al. Do human lymphocytes exposed to the fallout of the Chernobyl accident exhibit an adaptive response? Challenge with ionising radiation. *Mutat. Res.* 332: 33-38 (1995).
- P10 Parsons, R., G-M. Li, M. Longley et al. Mismatch repair deficiency in phenotypically normal human cells. *Science* 268: 738-740 (1995).
- P11 Paterson, M.C., A.K. Anderson, B.P. Smith et al. Enhanced radiosensitivity of cultured fibroblasts from ataxia telangiectasia heterozygotes manifested by defective colony-forming ability and reduced DNA repair replication after hypoxic X-irradiation. *Cancer Res.* 39: 3725-3734 (1979).
- P12 Paterson, M.C., S.J. McFarlane, N.E. Gentner et al. Cellular hypersensitivity to chronic γ -radiation in cultured fibroblasts from ataxia-telangiectasia heterozygotes. p. 73-87 in: Ataxia-telangiectasia: Genetics, Neuropathology, and Immunology of a Degenerative Disease of Childhood (R.A. Gatti & M. Swift, eds.). Liss, New York, 1985.
- P13 Pegg, A.E. Properties of O⁶-alkylguanine-DNA transferases. *Mutat. Res.* 233: 165-175 (1990).
- P14 Pohl-Ruling, J. and P. Fischer. The dose-effect relationship of chromosome aberrations to α and γ irradiation in a population subjected an increased burden of natural radioactivity. *Radiat. Res.* 80: 61-81 (1979).
- P15 Pohl-Ruling, J., P. Fischer, O. Haas et al. Effect of low-dose acute X-irradiation on the frequencies of chromosomal aberrations in human peripheral lymphocytes *in vitro*. *Mutat. Res.* 110: 71-82 (1983).
- P16 Pohl-Ruling, J., P. Fischer, D.C. Lloyd et al. Chromosomal damage induced in human lymphocytes by low doses of D-T neutrons. *Mutat. Res.* 173: 267-272 (1986).
- P17 Pohl-Ruling, J., O. Haas, A. Brogger et al. The effect on lymphocyte chromosomes of additional radiation burden due to fallout in Salzburg (Austria) from the Chernobyl accident. *Mutat. Res.* 262: 209-217 (1991).
- P18 Potten, C.S. Extreme sensitivity of some intestinal crypt cells to X and gamma irradiation. *Nature* 269: 518-520 (1977).
- P19 Pelevina, I.I., V.A. Nikolaev, V.Ya. Gotlib et al. Adaptive response in human blood lymphocytes after low dose chronic irradiation. *Radiat. Biol. Ecol.* 34: 805-817 (1994).
- P20 Pronk, J.C., R.A. Gibson, A. Savoia et al. Localization of the Fanconi anaemia complementation group A gene to chromosome 16q24.3. *Nature Genet.* 11: 338-343 (1995).
- P21 Porter, S.E., P.W. Greenwell, K.B. Ritchie et al. The DNA-binding protein Hdf1p (a putative Ku homologue) is required for maintaining normal telomere length in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 24: 582-585 (1996).
- P22 Petrini, J.H.J., M.E. Walsh, C. Dimare et al. Isolation and characterization of the human *MRE11* homologue. *Genomics* 29: 80-86 (1995).
- P23 Plug, A.W., J. Xu, G. Reddy et al. Presynaptic association of Rad51 protein with selected sites in meiotic chromatin. *Proc. Natl. Acad. Sci. U.S.A.* 93: 5920-5924 (1996).
- P24 Ponnaiya, B., M.N. Cornforth and R.L. Ullrich. Radiation-induced chromosomal instability in BALB/c and C57BL/6 mice: the difference is as clear as black and white. *Radiat. Res.* 147: 121-125 (1997).
- P25 Pampfer, S. and C. Streffer. Increased chromosome aberration levels in cells from mouse fetuses after zygote X-irradiation. *Int. J. Radiat. Biol.* 55: 85-92 (1989).
- P26 Pennycook, J.L., Y. Chang, J. Celler et al. High frequency of normal DJH joints in B cell progenitors in severe combined immunodeficiency mice. *J. Exp. Med.* 178: 1007-1016 (1993).
- P27 Prise, K.M., S. Davies and B.D. Michael. The relationship between radiation-induced DNA double-strand breaks and cell kill in hamster V79 fibroblasts irradiated with 250 kVp X-rays, 2.3 MeV neutrons or ²³⁸Pu alpha-particles. *Int. J. Radiat. Biol.* 52: 893-902 (1987).
- P28 Peak, M.J., L. Wang, C.K. Hill et al. Comparison of repair of DNA double-strand breaks caused by neutron or gamma radiation in cultured human cells. *Int. J. Radiat. Biol.* 60: 891-898 (1991).
- P29 Pfeiffer, P. and W. Vielmetter. Joining of nonhomologous DNA double strand breaks *in vitro*. *Nucleic Acids Res.* 16: 907-924 (1988).
- P30 Pflaum, M., O. Will and B. Epe. Determination of steady-state levels of oxidative DNA base modifications in mammalian cells by means of repair endonucleases. *Carcinogenesis* 18: 2225-2231 (1997).
- P31 Pouget, J-P., J-L. Ravenat, T. Douki et al. Measurement of DNA base damage in cells exposed to low doses of gamma radiation using the comet assay associated with DNA glycosylases. *Int. J. Radiat. Biol.* 55: 51-58 (1999).

- P32 Patel, K.J., V.P.C.C. Yu, H. Lee et al. Involvement of *Brca2* in DNA repair. *Mol. Cell* 1: 347-357 (1998).
- P33 Pelevina, A.A., V.Y. Gotlieb, O.V. Kudryashova et al. The properties of irradiated cell progeny. *Tsitologiya* 40: 467-477 (1998)
- P34 Paull, T.T. and M. Gellert. Nbs1 potentiates ATP-driven DNA unwinding and endonuclease cleavage by the Mre11/Rad50 complex. *Genes Dev.* 13: 1276-1288 (1999).
- P35 Prasad, A.V., N. Mohan, B. Chandrasekar et al. Activation of nuclear factor κ B in human lymphoblastoid cells by low-dose ionizing radiation. *Radiat. Res.* 138: 367-372 (1994).
- P36 Prasad, A.V., N. Mohan, B. Chandrasekar et al. Induction of transcription of "immediate early genes" by low-dose ionizing radiation. *Radiat. Res.* 143: 263-272 (1995).
- P37 Prise, K.M., G. Ahnstrom, M. Belli et al. A review of dsb induction data for varying quality radiations. *Int. J. Radiat. Biol.* 74: 173-184 (1998).
- R1 Roth, D. and J. Wilson. Illegitimate DNA recombination in mammalian cells. p. 621-653 in: *Genetic Recombination* (R. Kucherlapati and G.R. Smith, eds.). Am. Soc. Microbiol., Washington, 1988.
- R2 Rao, B.S., N.M.S. Reddy and U. Madhvanath. Gamma radiation response and recovery studies in radiation sensitive mutants of diploid yeast. *Int. J. Radiat. Biol.* 37: 701-705 (1980).
- R3 Reddy, N.M.S., K.B. Kshiti and P. Subrahmanyam. Absence of a dose-rate effect and recovery from sub-lethal damage in *rad52* strain of diploid yeast *Saccharomyces cerevisiae* exposed to γ -rays. *Mutat. Res.* 105: 145-148 (1982).
- R4 Russo, G., M. Isobe, L. Pegoraro et al. Molecular analysis of a t(7;14)(q35;q32) chromosome translocation in a T cell leukemia of a patient with ataxia telangiectasia. *Cell* 53: 137-144 (1988).
- R5 Runger, T.M., M. Poot and K.H. Kraemer. Abnormal processing of transfected plasmid DNA in cells from patients with ataxia telangiectasia. *Mutat. Res.* 293:47-54 (1992).
- R6 Russell, W.L., L.B. Russell and E.M. Kelly. Radiation dose rate and mutation frequency. *Science* 128: 1546-1550 (1958).
- R7 Russell, W.L. Repair mechanisms in radiation mutation induction in the mouse. *Brookhaven Symp. Biol.* 20: 179-189 (1968).
- R8 Radford, I.R., T.K. Murphy, J.M. Radley et al. Radiation response of mouse lymphoid and myeloid cell lines. II. Apoptotic death is shown by all lines examined. *Int. J. Radiat. Biol.* 65: 217-227 (1994).
- R9 Rasio, D., M. Negrini, G. Manenti et al. Loss of heterozygosity at chromosome 11q in lung adenocarcinoma: identification of 3 independent regions. *Cancer Res.* 55: 3988-3991 (1995).
- R10 Reed, J.C. Bcl-2 and the regulation of programmed cell death. *J. Cell Biol.* 124: 1-6 (1994).
- R11 Reitmair, A.H., R. Schmits, A. Ewel et al. MSH2 deficient mice are viable and susceptible to lymphoid tumours. *Nature Genet.* 11: 64-70 (1995).
- R12 Rigaud, O., D. Papadopoulo and E. Moustacchi. Decreased deletion mutation in radioadapted human lymphocytes. *Radiat. Res.* 133: 94-101 (1993).
- R13 Rinchik, E.M., J.P. Stoye, W.N. Frankel et al. Molecular analysis of viable spontaneous and radiation-induced albino (*c*)-locus mutations in the mouse. *Mutat. Res.* 286: 199-207 (1993).
- R14 Russell, L.B. Information from specific-locus mutants on the nature of induced and spontaneous mutations in the mouse. p. 437-447 in: *Genetic Toxicology of Environmental Chemicals, Part B*. Liss, New York, 1986.
- R15 Russell, N.S., C.F. Arlett, H. Bartelink et al. Use of fluorescence *in situ* hybridization to determine the relationship between chromosome aberrations and cell survival in eight human fibroblast strains. *Int. J. Radiat. Biol.* 68: 185-196 (1995).
- R16 Russell, W.L. Genetic effects from internally deposited radionuclides. p. 14-17: NCRP Report No. 89 (1987).
- R17 Russell, W.L. and E.M. Kelly. Mutation frequencies in male mice and the estimation of genetic hazards of radiation in men. *Proc. Natl. Acad. Sci. U.S.A.* 79: 542-544 (1982).
- R18 Roth, D.B. and J.H. Wilson. Nonhomologous recombination in mammalian cells: role of short sequence homologies in the joining reaction. *Mol. Cell. Biol.* 6: 4295-4304 (1986).
- R19 Rampino, N., H. Yamamoto, Y. Ionov et al. Somatic frameshift mutations in the *BAX* gene in colon cancers of the microsatellite mutator phenotype. *Science* 275: 967-969 (1997).
- R20 Rosselli, F., A. Ridet, R. Soussi et al. p53-dependent pathway of radio-induced apoptosis is altered in Fanconi anemia. *Oncogene* 10: 9-17 (1995).
- R21 Reuvers, A.P., C.L. Greenstock, J.Borsa et al. Studies on the mechanism of chemical radioprotection by dimethyl sulphoxide. *Int. J. Radiat. Biol.* 24: 533-536 (1973).
- R22 Reardon, J.T., T. Bessho, H.C. Kung et al. In vitro repair of oxidative DNA damage by human nucleotide excision repair system: possible explanation for neurodegeneration in xeroderma pigmentosum patients. *Proc. Natl. Acad. Sci. U.S.A.* 94: 9463-9468 (1997).
- R23 Rajan, J.V., M. Wang, S.T. Marquis et al. *Brca2* is coordinately regulated with *Brca1* during proliferation and differentiation in mammary epithelial cells. *Proc. Natl. Acad. Sci. U.S.A.* 93: 13078-13083 (1996).
- R24 Ramsden, D.A. and M. Gellert. Ku protein stimulates DNA end joining by mammalian DNA ligases: a direct role for Ku in repair of DNA double-strand breaks. *EMBO J.* 17: 609-614 (1998).
- R25 Ritter, M.A., J.E. Cleaver and C.A. Tobias. High-LET radiations induce a large proportion of non-rejoining DNA breaks. *Nature* 266: 653-655 (1977).
- R26 Rydberg, B. Clusters of DNA damage induced by ionizing radiation: formation of short DNA fragments. II. Experimental detection. *Radiat. Res.* 145: 200-209 (1996).
- R27 Rydberg, B., M. Loblrich and P.K. Cooper. DNA double-strand breaks induced by high-energy neon and iron ions in human fibroblasts. I. Pulsed-field gel electrophoresis method. *Radiat. Res.* 139: 133-141 (1994).
- R28 Rithidech, K.N., V.P. Bond, E.P. Cronkite et al. Hypermutability of mouse chromosome 2 during development of X-ray induced murine myeloid leukaemia. *Proc. Natl. Acad. Sci. U.S.A.* 92: 1152-1156 (1995).
- R29 Rosenberg, S.M. Mutation for survival. *Curr. Opin. Genet. Dev.* 7: 829-834 (1997).
- R30 Radman, M., I. Matic, J.A. Halliday et al. Editing DNA replication and recombination by mismatch repair: from bacterial genetics to mechanisms of predisposition to cancer in humans. *Philos. Trans. R. Soc. Lond.* 347: 97-103 (1995).
- R31 Ramsay, J., G. Birrell and M. Lavin. Testing for mutations of the ataxia telangiectasia gene in radiosensitive breast cancer patients. *Radiother. Oncol.* 47: 125-128 (1998).

- R32 Rio, P.G., D. Pernin, J.O. Bay et al. Loss of heterozygosity of BRCA1, BRCA2 and ATM genes in sporadic invasive ductal breast carcinoma. *Int. J. Oncol.* 13: 849-853 (1998).
- R33 Redpath, J.L. and R.J. Antoniono. Induction of an adaptive response against spontaneous neoplastic transformation in vitro by low-dose gamma radiation. *Radiat. Res.* 149: 517-520 (1998).
- R34 Ridet, A., C. Guillouf, E. Dechaud et al. Deregulated apoptosis is a hallmark of the Fanconi anaemia syndrome. *Cancer Res.* 57: 1722-1730 (1997).
- R35 Reuven, N.B., G. Arad, A. Maor-Shoshani et al. The mutagenesis protein UmuC is a DNA polymerase activated by UmuD', RecA, and SSB and is specialized for translesion replication. *J. Biol. Chem.* 274: 31763-31766 (1999).
- R36 Roberts, S.A., A.R. Spreadborough, B. Bulman et al. Heritability of cellular radiosensitivity: a marker of low-penetrance predisposition genes in breast cancer? *Am. J. Hum. Genet.* 65: 784-794 (1999).
- R37 Robson, T., M.C. Joiner, G.D. Wilson et al. A novel human stress response-related gene with a potential role in induced radioresistance. *Radiat. Res.* 152: 451-461 (1999).
- R38 Riballo, E., S.E. Critchlow, S-H. Teo et al. Identification of a defect in DNA ligase IV in a radiosensitive leukaemia patient. *Curr. Biol.* 9: 699-702 (1999).
- R39 Radford, I.R. Review: Initiation of ionizing radiation-induced apoptosis: DNA damage-mediated or does ceramide have a role? *Int. J. Radiat. Biol.* 75: 521-528 (1999).
- R40 Riboni, R., A. Casati, T. Nardo et al. Telomeric fusions in cultured human fibroblasts as a source of genomic instability. *Cancer Genet. Cytogenet.* 95: 130-136 (1997).
- R41 Radicella, J.P., C. Dherin, C. Desmaze et al. Cloning and characterization of hOGG1, a human homolog of the OGG1 gene of *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. U.S.A.* 94: 8010-8015 (1997).
- S1 Sancar, A. and G.B. Sancar. DNA repair enzymes. *Annu. Rev. Biochem.* 57: 29-67 (1988).
- S2 Selby, C.P. and A. Sancar. Mechanisms of transcription-repair coupling and mutation frequency decline. *Microbiol. Rev.* 58: 317-329 (1994).
- S3 von Sonntag, C. *The Chemical Basis of Radiation Biology*. Taylor and Francis, London, 1987.
- S4 Sabovljevic, S.A., W.A. Cramp, P.D. Lewis et al. Use of rapid tests of cellular radiosensitivity in radiotherapeutic practice. *Lancet* ii: 787 (1985).
- S5 Sanford, K.K., R. Parshad, R. Gantt et al. Factors affecting and significance of G₂ chromatin radiosensitivity in predisposition to cancer. *Int. J. Radiat. Biol.* 55: 963-981 (1989).
- S6 Schlissel, M., A. Constantinescu, T. Morrow et al. Double-strand signal sequence breaks in V(D)J recombination are blunt, 5' phosphorylated, RAG-dependent and cell cycle regulated. *Genes Dev.* 7: 2520-2532 (1993).
- S7 Seemanova, E.E., E. Passarge, D. Beneskova et al. Familial microcephaly with normal intelligence, immunodeficiency and risk for lymphoreticular malignancies. *Am. J. Med. Genet.* 20: 639-648 (1985).
- S8 Seemanova, E. An increased risk for malignant neoplasms in heterozygotes for a syndrome of microcephaly, normal intelligence, growth retardation, remarkable facies, immunodeficiency and chromosomal instability. *Mutat. Res.* 238: 321-324 (1990).
- S9 Siede, W., A.S. Friedberg and E.C. Freidberg. *RAD9*-dependent G₁ arrest defines a second checkpoint for damaged DNA in the cell cycle of *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. U.S.A.* 90: 7985-7989 (1993).
- S10 Shinohara, A., H. Ogawa, Y. Matsuda et al. Cloning of human, mouse and fission yeast recombination genes homologous to *RAD51* and *recA*. *Nature Genet.* 4: 239-243 (1993).
- S11 Smith, P.J., T.A. Makinson and J.V. Watson. Enhanced sensitivity to camptothecin in ataxia-telangiectasia cells and its relationship with the expression of DNA topoisomerase I. *Int. J. Radiat. Biol.* 55: 217-231 (1989).
- S12 Swift, M., D. Morrell, E. Cromartie et al. The incidence and gene frequency of ataxia-telangiectasia in the United States. *Am. J. Hum. Genet.* 39: 573-583 (1986).
- S13 Swift, M., D. Morrell, R.B. Massey et al. Incidence of cancer in 161 families affected by ataxia-telangiectasia. *N. Engl. J. Med.* 325: 1831-1836 (1991).
- S14 Stamato, T.D., A. Dipatri and A. Giaccia. Cell cycle dependent repair of potentially lethal damage in the XR-1 gamma-ray sensitive CHO cell. *Radiat. Res.* 115: 325-333 (1988).
- S15 Savage, J.R.K. and P.J. Simpson. Fish painting patterns resulting from complex exchanges. *Mutat. Res.* 312: 51-60 (1994).
- S16 Seymour, C.B., C. Mothersill and T. Alper. High yields of lethal mutations in somatic mammalian cells that survive ionising radiation. *Int. J. Radiat. Biol.* 50: 167-179 (1986).
- S17 Simpson, P., T. Morris, J. Savage et al. High-resolution cytogenetic analysis of X-Ray induced mutations of the HPRT gene of primary human fibroblasts. *Cytogenet. Cell Genet.* 64: 39-45 (1993).
- S18 Singleton, B. and J. Thacker. Communication to the UNSCEAR Secretariat (1995).
- S19 Sadamoto, S., S. Suzuki, K. Kamiya et al. Radiation induction of germline mutation at a hypervariable mouse minisatellite locus. *Int. J. Radiat. Biol.* 65: 549-557 (1994).
- S20 Sanderson, B.J.S. and A.A. Morley. Exposure of human lymphocytes to ionising radiation reduces mutagenesis in subsequent ionising radiation. *Mutat. Res.* 164: 347-351 (1986).
- S21 Sands, A.T., M.B. Suraokar, A. Sanchez et al. p53 deficiency does not affect the accumulation of point mutations in a transgene target. *Proc. Natl. Acad. Sci. U.S.A.* 92: 8517-8521 (1995).
- S22 Sankaranayanan, K., A. van Duyn, M.J. Loos et al. Adaptive response of human lymphocytes to low-level radiation from radioisotopes or X-rays. *Mutat. Res.* 211: 7-12 (1989).
- S23 Sasaki, M.S. On the reaction kinetics of the radioadaptive response in cultured mouse cells. *Int. J. Radiat. Biol.* 68: 281-291 (1995).
- S24 Savitsky, K., A. Bar-Shira, S. Gilad et al. A single ataxia telangiectasia gene with a product similar to PI-3 kinase. *Science* 268: 1749-1753 (1995).
- S25 Savitsky, K., S. Sfez, D.A. Tagle et al. The complete sequence of the coding region of the ATM gene reveals similarity to cell cycle regulators in different species. *Hum. Mol. Genet.* 4: 2025-2032 (1995).
- S26 Schmid, E., M. Bauchinger and U. Nahrstedt. Adaptive response after X-irradiation of human lymphocytes? *Mutagenesis* 4: 87-89 (1989).
- S27 Scott, D., A. Spreadborough, E. Levine et al. Genetic predisposition in breast cancer. *Lancet* 344: 1444 (1994).
- S28 Scott, D., A.R. Spreadborough, L.A. Jones et al. Chromosomal radiosensitivity in G₂-phase lymphocytes as an indicator of cancer predisposition. *Radiat. Res.* 145: 3-16 (1996).

- S29 Searle, A.G. Germ-cell sensitivity in the mouse: a comparison of radiation and chemical mutagens. p. 169-177 in: Environmental Mutagens and Carcinogens (T. Sugimura et al., eds.). University of Tokyo, 1982.
- S30 Shadley, J.D. Chromosomal adaptive response in human lymphocytes. *Radiat. Res.* 138: S9-S12 (1994).
- S31 Shadley, J.D., V. Afzal and S. Wolff. Characterization of the adaptive response to ionising radiation induced by low doses of X-rays to human lymphocytes. *Radiat. Res.* 111: 511-517 (1987).
- S32 Shadley, J.D. and G. Dai. Cytogenetic and survival adaptive responses in G₁ phase human lymphocytes. *Mutat. Res.* 265: 273-281 (1992).
- S33 Shadley, J.D. and J.K. Wiencke. Induction of the adaptive response by X-rays is dependent on radiation intensity. *Int. J. Radiat. Biol.* 56: 107-118 (1989).
- S34 Shadley, J.D. and S. Wolff. Very low doses of X-rays can cause human lymphocytes to become less susceptible to ionising radiation. *Mutagenesis* 2: 95-96 (1987).
- S35 Skov, K., B. Marples, J.B. Matthews et al. A preliminary investigation into the extent of increased radiosensitivity or hyper-radiosensitivity in cells of hamster lines known to be deficient in DNA repair. *Radiat. Res.* 138: S126-S129 (1994).
- S36 Sloan, S.R., E.W. Newcomb and A. Pellicer. Neutron radiation can activate *K-ras* via a point mutation in codon 146 and induces a different spectrum of *ras* mutations than does gamma radiation. *Mol. Cell. Biol.* 10: 405-408 (1990).
- S37 Sinclair, W.K. Cyclic X-ray responses in mammalian cells in vitro. *Radiat. Res.* 33: 620-643 (1968).
- S38 Sinclair, W.K. Cell cycle dependence of the lethal radiation response in mammalian cells. *Curr. Top. Radiat. Res. Quart.* 7: 264-285 (1972).
- S39 Stacey, M., S. Thacker and A.M.R. Taylor. Cultured skin keratinocytes from both normal individuals and basal cell naevus syndrome patients are more resistant to γ -rays and UV light compared with cultured skin fibroblasts. *Int. J. Radiat. Biol.* 56: 45-58 (1989).
- S40 Savitsky, K., M. Platzer, T. Uziel et al. Ataxia-telangiectasia: structural diversity of untranslated sequences suggests complex post-transcriptional regulation of ATM gene expression. *Nucleic Acids Res.* 25: 1678-1684 (1997).
- S41 Swift, M., D. Morrell, R.B. Massey et al. Risk of breast cancer in ataxia-telangiectasia. *Lancet* 326: 1360 (1992).
- S42 Symonds, H., L. Krall, L. Remington et al. p53-dependent apoptosis suppresses tumor growth and progression in vivo. *Cell* 78: 703-711 (1994).
- S43 Sanchez, Y., B.A. Desany, W.J. Jones et al. Regulation of *RAD53* by the *ATM*-like kinases *MEC1* and *TEL1* in yeast cell cycle checkpoint pathways. *Science* 271: 357-360 (1996).
- S44 Shen, Z.Y., K. Denison, R. Lobb et al. The human and mouse homologs of the yeast *RAD52* gene: cDNA cloning, sequence analysis, assignment to human chromosome 12p12.2-p13, and mRNA expression in mouse tissues. *Genomics* 25: 199-206 (1995).
- S45 Sturzbecher, H-W., B. Donzelmann, W. Henning et al. p53 is linked directly to homologous recombination processes via *RAD51/RecA* protein interaction. *EMBO J.* 15: 1992-2002 (1996).
- S46 Strathdee, C.A., A.M.V. Duncan and M. Buchwald. Evidence for at least 4 Fanconi anaemia genes including *FACC* on chromosome 9. *Nature Genet.* 1: 196-198 (1992).
- S47 Strathdee, C.A., H. Gavish, W.R. Shannon et al. Cloning of cDNAs for Fanconi's anaemia by functional complementation. *Nature* 356: 763-767 (1992).
- S48 Saito, H., A.T. Hammond and R.E. Moses. Hyper-sensitivity to oxygen is a uniform and secondary defect in Fanconi anaemia cells. *Mutat. Res.* 294: 255 (1993).
- S49 Seeberg, E., L. Eide and M. Bjoras. The base excision repair pathway. *Trends Biochem. Sci.* 20: 391-397 (1995).
- S50 Siede, W., A.A. Friedl, I. Dianova et al. The *Saccharomyces cerevisiae* Ku autoantigen homologue affects radiosensitivity only in the absence of homologous recombination. *Genetics* 142: 91-102 (1996).
- S51 Scully, R., J. Chen, A. Plug et al. Association of *BRCA1* with *Rad51* in mitotic and meiotic cells. *Cell* 88: 265-275 (1997).
- S52 Strathern, J.N., B.K. Shafer and C.B. McGill. DNA synthesis errors associated with double-strand break repair. *Genetics* 140: 965-972 (1995).
- S53 Sabatier, L., B. Dutrillaux and M.B. Martin. Chromosomal instability. *Nature* 357: 548 (1992).
- S54 Savage, J.R.K. and D.G. Papworth. Comment on the ratio of chromosome-type dicentric interchanges to centric rings for track-clustered compared with random breaks. *Radiat. Res.* 146: 236-240 (1996).
- S55 Santana, P., L.A. Pena, A. Haimovitz-Friedman et al. Acid sphingomyelinase-deficient human lymphoblasts and mice are defective in radiation-induced apoptosis. *Cell* 86: 189-199 (1996).
- S56 Satoh, M.S., C.J. Jones, R.D. Wood et al. DNA excision-repair defect of xeroderma pigmentosum prevents removal of a class of oxygen free radical-induced base lesions. *Proc. Natl. Acad. Sci. U.S.A.* 90: 6335-6339 (1993).
- S57 Schiestl, R.H., F. Khogali and N. Carls. Reversion of the mouse *pink-eyed unstable* mutation induced by low doses of X-rays. *Science* 266: 1573-1576 (1994).
- S58 Singhal, R.K., R. Prasad and S.H. Wilson. DNA polymerase β conducts the gap-filling step in uracil-initiated base excision repair in a bovine testis nuclear extract. *J. Biol. Chem.* 270: 949-957 (1995).
- S59 Saar, K., K.H. Chrzanoska, M. Stumm et al. The gene for the ataxia-telangiectasia variant, Nijmegen breakage syndrome, maps to a 1-cm interval on chromosome 8q21. *Am. J. Hum. Genet.* 60: 605-610 (1997).
- S60 Stewart, E., C.R. Chapman, F. Al-Khodairy et al. *rhq1⁺*, a fission yeast gene related to the Bloom's and Werner's syndrome genes, is required for reversible S phase arrest. *EMBO J.* 16: 2682-2692 (1997).
- S61 Sharan, S.K., M. Morimatsu, U. Albrecht et al. Embryonic lethality and radiation hypersensitivity mediated by *Rad51* in mice lacking *Brca2*. *Nature* 386: 804-810 (1997).
- S62 Satoh, M.S., G.G. Poirier and T. Lindahl. Dual function for poly(ADP-ribose) synthesis in response to DNA strand breakage. *Biochemistry* 33: 7099-7106 (1994).
- S63 Schreiber, V., D. Hunting, C. Trucco et al. A dominant-negative mutant of human poly(ADP-ribose) polymerase affects cell recovery, apoptosis and sister-chromatid exchange following DNA damage. *Proc. Natl. Acad. Sci. U.S.A.* 92: 4753-4757 (1995).
- S64 Schar, P., G. Herrmann, G. Daly et al. A newly identified DNA ligase of *Saccharomyces cerevisiae* involved in *RAD52*-independent repair of DNA double-strand breaks. *Genes Dev.* 11: 1912-1924 (1997).
- S65 Shafman, T., K.K. Khanna, P. Kedar et al. Interaction between ATM protein and c-Abl in response to DNA damage. *Nature* 387: 520-523 (1997).
- S66 Schmid, E. and M. Bauchinger. Comments on direct biological evidence for a significant neutron dose to survivors of the Hiroshima atomic bomb. *Radiat. Res.* 146: 479-482 (1996).

- S67 Satoh, C. and M. Kodaira. Effects of radiation on children. *Nature* 383: 226 (1996).
- S68 Sala-Trepat, M., J. Boyse, P. Richard et al. Frequencies of *HPRT* mutants in T-lymphocytes and of glycophorin A variants in erythrocytes of Fanconi anemia patients and control donors. *Mutat. Res.* 289: 115-126 (1993).
- S69 Shayeghi, M., S. Seal, J. Regan et al. Heterozygosity for mutations in the ataxia telangiectasia gene is not a major cause of radiotherapy complications in breast cancer patients. *Br. J. Cancer* 78: 922-927 (1998).
- S70 Starostik, P., T. Manshouri, S. O'Brien et al. Deficiency of the ATM protein expression defines an aggressive subgroup of B-cell chronic lymphocytic leukemia. *Cancer Res.* 58: 4552-4557 (1998).
- S71 Sachs, R.K., D.J. Brenner, A.M. Chen et al. Intra-arm and inter-arm chromosome intrachanges: tools for probing geometry and dynamics of chromatin. *Radiat. Res.* 148: 330-340 (1997).
- S72 Savage, J.R.K. and D.G. Papworth. An investigation of LET 'fingerprints' in *Tradescantia*. *Mutat. Res.* 422: 313-322 (1998).
- S73 Siliciano, J.D., C.E. Canman, Y. Taya et al. DNA damage induces phosphorylation of the amino terminus of p53. *Genes Dev.* 11: 3471-3481 (1997).
- S74 Schwartz, D., N. Almog, A. Peled et al. Role of wild type p53 in the G2 phase: regulation of the gamma-irradiation-induced delay and DNA repair. *Oncogene* 15: 2597-2607 (1997).
- S75 Stewart, N., G.G. Hicks, F. Paraskevas et al. Evidence for a second cell-cycle block at G2/M by p53. *Oncogene* 10: 109-115 (1995).
- S76 Stankovic, T., P. Weber, G. Stewart et al. Inactivation of ataxia telangiectasia mutated gene in B-cell chronic lymphocytic leukaemia. *Lancet* 353: 26-29 (1999).
- S77 Suzuki, K., S. Kodama and M. Watanabe. Recruitment of ATM protein to double strand DNA irradiated with ionizing radiation. *J. Biol. Chem.* 274: 25571-25575 (1999).
- S78 Stankovic, T., A.M.J. Kidd, A. Sutcliffe et al. *ATM* mutations and phenotypes in ataxia-telangiectasia families in the British Isles: expression of mutant *ATM* and the risk of leukemia, lymphoma, and breast cancer. *Am. J. Hum. Genet.* 62: 334-345 (1998).
- S79 Sadekova, S., S. Lehnert and T.Y.-K. Chow. Induction of PBP74/mortalin/Grp75, a member of the hsp70 family, by low doses of ionizing radiation: a possible role in induced radioresistance. *Int. J. Radiat. Biol.* 72: 653-660 (1997).
- S80 Shimizu, T., T. Kato, A. Tachibana et al. Coordinated regulation of radioadaptive response by protein kinase C and p38 mitogen-activated protein kinase. *Exp. Cell Res.* 251: 424-432 (1999).
- S81 Svistuneko, D.A., G.Z. Ju, J. Wei et al. EPR study of mouse tissues in search for adaptive responses to low level whole-body X-irradiation. *Int. J. Radiat. Biol.* 62: 327-336 (1992).
- S82 Stewart, G.S., R.S. Maser, T. Stankovic et al. The DNA double-strand break repair gene *hMRE11* is mutated in individuals with an ataxia-telangiectasia-like disorder. *Cell* 99: 577-587 (1999).
- S83 Smith, B.T. and G.C. Walker. Mutagenesis and more: umuDC and the *Escherichia coli* SOS response. *Genetics* 148: 1599-1610 (1998).
- S84 Shen, M.R., I.M. Jones and H. Mohrenweiser. Non-conservative amino acid substitution variants exist at polymorphic frequency in DNA repair genes in healthy humans. *Cancer Res.* 58: 604-608 (1998).
- S85 Sargent, R.G., R.L. Rolig, A.E. Kilburn et al. Recombination-dependent deletion formation in mammalian cells deficient in the nucleotide excision repair gene ERCC1. *Proc. Natl. Acad. Sci. U.S.A.* 94: 13122-13127 (1997).
- S86 Scully, R., S. Ganesan, K. Vlasakova et al. Genetic analysis of BRCA1 function in a defined tumor cell line. *Mol. Cell* 4: 1093-1099 (1999).
- S87 Sutherland, B.M., P.V. Bennett, O. Sidorkina et al. Clustered DNA damages induced in isolated DNA and in human cells by low doses of ionizing radiation. *Proc. Natl. Acad. Sci. U.S.A.* 97: 103-108 (2000).
- S88 Stoll, U., A. Schmidt, E. Schneider et al. Killing and mutation of Chinese hamster V79 cells exposed to accelerated oxygen and neon ions. *Radiat. Res.* 142: 288-294 (1995).
- S89 Stoll, U., B. Barth, N. Scheerer et al. HPRT mutations in V79 Chinese hamster cells induced by accelerated Ni, Au and Pb ions. *Int. J. Radiat. Biol.* 70: 15-22 (1996).
- S90 Schmidt, P. and J. Kiefer. Deletion-pattern analysis of alpha-particle and X-ray induced mutations at the HPRT locus of V79 Chinese hamster cells. *Mutat. Res.* 421: 149-161 (1998).
- T1 Taylor, A.M.R. Unrepaired DNA strand breaks in irradiated ataxia telangiectasia lymphocytes suggested from cytogenetic observations. *Mutat. Res.* 50: 407-418 (1978).
- T2 Taalman, R.D.F.M., N.G.J. Jaspers, J.M. Scheres et al. Hypersensitivity to ionising radiation, in vitro, in a new chromosomal breakage disorder, the Nijmegen breakage syndrome. *Mutat. Res.* 112: 23-32 (1983).
- T3 Taylor, A.M.R., E. Flude, B. Laher et al. Variant forms of ataxia-telangiectasia. *J. Med. Genet.* 24: 669-677 (1987).
- T4 Taccioli, G.E., G. Rathbun, G. Oltz et al. Impairment of V(D)J recombination in double-strand break repair mutants. *Science* 260: 207-210 (1993).
- T5 Taccioli, G.E., T.M. Gottlieb, T. Blunt et al. Ku80: product of the XRCC5 gene and its role in DNA repair and V(D)J recombination. *Science* 265: 1442-1445 (1994).
- T6 Tatsumi-Miyajima, J., T. Yagi and H. Takebe. Analysis of mutations caused by DNA double-strand breaks produced by a restriction enzyme in shuttle vector plasmids propagated in ataxia telangiectasia cells. *Mutat. Res.* 294: 317-323 (1993).
- T7 Thacker, J. and A. Stretch. Responses of 4 X-ray-sensitive CHO cell mutants to different radiations and to irradiation conditions promoting cellular recovery. *Mutat. Res.* 146: 99-108 (1985).
- T8 Thacker, J. and A.N. Ganesh. DNA break repair, radioresistance of DNA synthesis and camptothecin sensitivity in the radiation-sensitive *irs* mutants: comparisons to ataxia-telangiectasia cells. *Mutat. Res.* 235: 49-58 (1990).
- T9 Thacker, J. and R. Wilkinson. The genetic basis of resistance to ionising radiation damage in cultured mammalian cells. *Mutat. Res.* 254: 135-142 (1991).
- T10 Thacker, J., J. Chalk, A. Ganesh et al. A mechanism for deletion formation in DNA by human cell extracts: the involvement of short sequence repeats. *Nucleic Acids Res.* 20: 6183-6188 (1992).
- T11 Thacker, J., A.N. Ganesh, A. Stretch et al. Gene mutation and V(D)J recombination in the radiosensitive *irs* lines. *Mutagenesis* 9: 163-168 (1994).
- T12 Taylor, J.A., M.A. Watson, T.R. Devereux et al. p53 mutation hotspot in radon-associated lung cancer. *Lancet* 343: 86-87 (1994).
- T13 Thacker, J. and R. Cox. Mutation induction and inactivation in mammalian cells exposed to ionizing radiation. *Nature* 258: 429-431 (1975).

- T14 Thacker, J., A. Stretch and M.A. Stephens. The induction of thioguanine-resistant mutants of Chinese hamster cells by gamma-rays. *Mutat. Res.* 42: 313-326 (1977).
- T15 Thacker, J., M.A. Stephens and A. Stretch. Mutation to ouabain-resistance in Chinese hamster cells: induction by ethyl methanesulphonate and lack of induction by ionising radiation. *Mutat. Res.* 51: 255-270 (1978).
- T16 Thacker, J., A. Stretch and M.A. Stephens. Mutation and inactivation of cultured mammalian cells exposed to beams of accelerated heavy ions. II. Chinese hamster cells. *Int. J. Radiat. Biol.* 36: 137-148 (1979).
- T17 Thacker, J. The involvement of repair processes in radiation-induced mutation of cultured mammalian cells. p. 612-620 in: *Radiation Research* (S. Okada et al. eds.). Japanese Association for Radiation Research, Tokyo, 1979.
- T18 Thacker, J., A. Stretch and D.T. Goodhead. The mutagenicity of α -particles from plutonium-238. *Radiat. Res.* 92: 343-352 (1982).
- T19 Thacker, J. and R. Cox. The relationship between specific chromosome aberrations and radiation-induced mutations in cultured mammalian cells. p. 235-275 in: *Radiation-induced Chromosome Damage in Man* (T. Ishihara and M.S. Sasaki, eds.). Liss, New York, 1983.
- T20 Thacker, J. The nature of mutants induced by ionising radiation in cultured hamster cells. III. Molecular characterization of HPRT-deficient mutants induced by gamma-rays or alpha-particles showing that the majority have deletions of all or part of the *hprt* gene. *Mutat. Res.* 160: 267-275 (1986).
- T21 Thacker, J. Molecular nature of ionising radiation-induced mutations of native and introduced genes in mammalian cells. p. 221-229 in: *Ionising Radiation Damage to DNA: Molecular Aspects* (S. Wallace and R. Painter, eds.). Wiley/Liss, New York, 1990.
- T22 Thacker, J. Radiation-induced mutation in mammalian cells at low doses and dose rates. *Adv. Radiat. Biol.* 16: 77-124 (1992).
- T23 Trakhtenbrot, L., R. Kranthgamer, P. Resnitzky et al. Deletion of chromosome 2 is an early event in the development of radiation-induced myeloid leukaemia in SJL/J mice. *Leukaemia* 2: 545-550 (1991).
- T24 Thacker, J. The study of responses the 'model' DNA breaks induced by restriction endonucleases in cells and cell-free systems: achievements and difficulties. *Int. J. Radiat. Biol.* 66: 591-596 (1994).
- T25 Takai, S., K.K. Sanford and R.E. Tarone. A procedure for carrier detection in ataxia-telangiectasia. *Cancer Genet. Cytogenet.* 46: 139-140 (1990).
- T26 Tanzarella, C., R. de Salvia, F. Degrassi et al. Effect of post-treatments with caffeine during G₂ on the frequency of chromosome type aberrations produced by X-rays in human lymphocytes during G₀ and G₁. *Mutagenesis* 1: 41-44 (1986).
- T27 Terasima, T. and L.J. Tolmach. X-ray sensitivity and DNA synthesis in synchronous populations of HeLa cells. *Science* 140: 490-492 (1963).
- T28 Thacker, J. Radiation-induced mutations in mammalian cells: what we know and what we do not know. p. 107 in: *37th Annual Meeting of the Radiation Research Society*, Seattle, 1989.
- T29 Thacker, J. Cellular radiosensitivity in ataxia-telangiectasia. *Int. J. Radiat. Biol.* 66: S87-S96 (1994).
- T30 Thacker, J. and R.E. Wilkinson. The genetic basis of cellular recovery from radiation damage: response of the radiosensitive *irs* lines to low-dose-rate irradiation. *Radiat. Res.* 144: 294-300 (1995).
- T31 Timofeeff-Ressovsky, N.W. The experimental production of mutations. *Biol. Rev. Camb. Philos. Soc.* 9: 411-457 (1934).
- T32 Tsuboi, K., T.C. Yang and D.J. Chen. Charged particle mutagenesis. I. Cytotoxic and mutagenic effects of high-LET charged iron particles on human skin fibroblasts. *Radiat. Res.* 129: 171-176 (1992).
- T33 Tuschl, H., H. Altmann, R. Kovac et al. Effects of low-dose radiation on repair processes in human lymphocytes. *Radiat. Res.* 81: 1-9 (1980).
- T34 Tsuzuki, T., Y. Fujii, K. Sakumi et al. Targeted disruption of the *Rad51* gene leads to lethality in embryonic mice. *Proc. Natl. Acad. Sci. U.S.A.* 93: 6236-6240 (1996).
- T35 Tambini, C.E., A.M. George, J.M. Rommens et al. The *XRCC2* DNA repair gene: identification of a positional candidate. *Genomics* 41: 84-92 (1997).
- T37 Tashiro, S., N. Kotomura, A. Shinohara et al. S-phase specific formation of the human Rad51 protein nuclear foci in lymphocytes. *Oncogene* 12: 2165-2170 (1996).
- T38 Tomlinson, I.P.M., M.R. Novelli and W.F. Bodmer. The mutation rate and cancer. *Proc. Natl. Acad. Sci. U.S.A.* 93: 14800-14803 (1996).
- T39 Thyagarajan, B. and C. Campbell. Elevated homologous recombination activity in Fanconi anemia fibroblasts. *J. Biol. Chem.* 272: 23328-23333 (1997).
- T40 Teo, S-H. and S.P. Jackson. Identification of *Saccharomyces cerevisiae* DNA ligase IV: involvement in DNA double-strand break rejoining. *EMBO J.* 16: 4788-4795 (1997).
- T41 Tsukamoto, Y., J. Kato and H. Ikeda. Silencing factors participate in DNA repair and recombination in *Saccharomyces cerevisiae*. *Nature* 388: 900-903 (1997).
- T42 Takata, M., M.S. Sasaki, E. Sonoda et al. Homologous recombination and non-homologous end-joining pathways of DNA double-strand break repair have overlapping roles in the maintenance of chromosomal integrity in vertebrate cells. *EMBO J.* 17: 5497-5508 (1998).
- T43 Takagi, M., D. Delia, L. Chessa et al. Defective control of apoptosis, radiosensitivity, and spindle checkpoint in ataxia telangiectasia. *Cancer Res.* 58: 4923-4929 (1998).
- T44 Thacker, J. Repair of ionizing radiation damage in mammalian cells: alternative pathways and their fidelity. *C.R. Acad. Sci. Paris* 322: (1999, in press).
- T45 Tang, M., X. Shen, E.G. Frank et al. UmuD'(2)C is an error-prone DNA polymerase, *Escherichia coli* pol V. *Proc. Natl. Acad. Sci. U.S.A.* 96: 8919-8924 (1999).
- T46 Tutt, A., A. Gabriel, D. Bertwistle et al. Absence of *Brca2* causes genome instability by chromosome breakage and loss associated with centrosome amplification. *Curr. Biol.* 9: 1107-1110 (1999).
- T47 Thacker, J. A surfeit of RAD51-like genes? *Trends Genet.* 15: 166-168 (1999).
- T48 Tchou, J., H. Kasai, S. Shibutani et al. 8-oxoguanine (8-hydroxyguanine) DNA glycosylase and its substrate specificity. *Proc. Natl. Acad. Sci. U.S.A.* 88: 4690-4694 (1991).
- T49 Thomas, C.B., D.O. Nelson, P. Pleshanov et al. Elevated frequencies of hypoxanthine phosphoribosyltransferase lymphocyte mutants are detected in Russian liquidators 6 to 10 years after exposure to radiation from the Chernobyl nuclear power plant accident. *Mutat. Res.* 439: 105-119 (1999).
- U2 United Nations. Sources and Effects of Ionizing Radiation. United Nations Scientific Committee on the Effects of Atomic Radiation, 1994 Report to the General Assembly, with scientific annexes. United Nations sales publication E.94.IX.11. United Nations, New York, 1994.

- U3 United Nations. Sources and Effects of Ionizing Radiation. United Nations Scientific Committee on the Effects of Atomic Radiation, 1993 Report to the General Assembly, with scientific annexes. United Nations sales publication E.94.IX.2. United Nations, New York, 1993.
- U14 Umansky, S.R., B.A. Korol and P.A. Nelipovitch. *In vivo* DNA degradation in thymocytes of gamma-irradiated or hydrocortisone-treated rats. *Biochim. Biophys. Acta* 665: 9-17 (1981).
- U15 Utsumi, H. and M.S. Sasaki. Deficient repair of potentially lethal damage in actively growing ataxia telangiectasia cells. *Radiat. Res.* 97: 407-413 (1984).
- U16 Unzhakov, S.V., I.M. Vasilieva, I.A. Meliksetova et al. Adaptive response formation in lymphocytes from the children exposed to small doses of radiation as a result of the Chernobyl accident. *Radiat. Biol. Ecol.* 34: 827-831 (1994).
- U17 Ueno, A.M., D.B. Vannais, D.L. Gustafson et al. A low adaptive dose of gamma rays reduced the number and altered the spectrum of S1-mutants in human-hamster hybrid A₁ cells. *Mutat. Res.* 358: 161-169 (1996).
- U18 Usui, T., T. Ohta, H. Oshiumi et al. Complex formation and functional versatility of Mre11 of budding yeast in recombination. *Cell* 95: 705-716 (1998).
- V1 Vorechovsky, I., D. Scott, M.R. Haeney et al. Chromosomal radiosensitivity in common variable immune deficiency. *Mutat. Res.* 290: 255-264 (1993).
- V2 Vahakangas, K.H., J.M. Samet, R.A. Metcalf et al. Mutations of p53 and ras genes associated lung cancer from uranium miners. *Lancet* 339: 576-580 (1992).
- V3 Venitt, S. and P.J. Biggs. Radon, mycotoxins, p53 and uranium mining. *Lancet* 343: 795 (1994).
- V4 Vijayalaxmi and W. Burkart. Resistance and cross-resistance to chromosome damage in human blood lymphocytes adapted to bleomycin. *Mutat. Res.* 211: 1-5 (1989).
- V5 Vorechovsky, I., L. Luo, M.J.S. Dyer et al. Clustering of missense mutations in the ataxia-telangiectasia gene in a sporadic T-cell leukaemia. *Nature Genet.* 17: 96-99 (1997).
- V6 Varon, R., C. Vissinga, M. Platzer et al. Nibrin, a novel DNA double-strand break repair protein, is mutated in Nijmegen breakage syndrome. *Cell* 93: 467-476 (1998).
- V7 van der Burgt, I., K.H. Chrzanowska, D. Smeets et al. Nijmegen breakage syndrome. *J. Med. Genet.* 33: 153-156 (1996).
- V8 Vorechovsky, I., D. Rasio, L. Luo et al. The *ATM* gene and susceptibility to breast cancer: analysis of 38 breast tumours reveals no evidence for mutation. *Cancer Res.* 56: 2726-2732 (1996).
- V9 Vorechovsky, I., L. Luo, A. Lindblom et al. *ATM* mutations in cancer families. *Cancer Res.* 56: 4130-4133 (1996).
- W1 Wallace, S.S. AP endonucleases and DNA glycosylases that recognize oxidative DNA damage. *Environ. Mol. Mutagen.* 12: 431-477 (1988).
- W2 Weichselbaum, R.R., J. Nove and J.B. Little. Deficient recovery from potentially lethal radiation damage in ataxia telangiectasia and xeroderma pigmentosum. *Nature* 271: 261-262 (1978).
- W3 Weichselbaum, R.R., J. Nove and J.B. Little. X-ray sensitivity of 53 human diploid fibroblast cell strains from patients with characterized genetic disorders. *Cancer Res.* 40: 920-925 (1980).
- W4 Weemaes, C.M.R., T.W.J. Hustinx, J.M.J.C. Scheres et al. New chromosome instability disorder: the Nijmegen breakage syndrome. *Acta Paediatr. Scand.* 70: 557-562 (1981).
- W5 Wegner, R.D., M. Metzger, F. Hanefeld et al. A new chromosome instability disorder confirmed by complementation studies. *Clin. Genet.* 33: 20-32 (1988).
- W6 Weinert, T.A. and L.H. Hartwell. The *rad9* gene controls the cell cycle response to DNA damage in *Saccharomyces cerevisiae*. *Science* 241: 317-322 (1988).
- W7 Weinert, T.A. and L.H. Hartwell. Characterization of *RAD9* of *Saccharomyces cerevisiae* and evidence that its function acts posttranslationally in cell cycle arrest after DNA damage. *Mol. Cell. Biol.* 10: 6554-6564 (1990).
- W8 Williams, G.T. Programmed cell death - apoptosis and oncogenesis. *Cell* 65: 1097-1098 (1991).
- W9 Woods, W.G., T.D. Byrne and T.H. Kim. Sensitivity of cultured cells to gamma radiation in a patient exhibiting marked *in vivo* radiation sensitivity. *Cancer* 62: 2341-2345 (1988).
- W10 Woods, C.G., S.E. Bunday and A.M.R. Taylor. Unusual features in the inheritance of ataxia telangiectasia. *Hum. Genet.* 84: 555-562 (1990).
- W11 Walker, G.C. Mutagenesis and inducible responses to deoxyribonucleic acid damage in *Escherichia coli*. *Microbiol. Rev.* 48: 60-93 (1984).
- W12 Waldren, C., L. Correll, M.A. Sognier et al. Measurement of low levels of X-ray mutagenesis in relation to human disease. *Proc. Natl. Acad. Sci. U.S.A.* 83: 4839-4843 (1986).
- W13 Waghray, M., S. Al-Sedairy, P.T. Ozand et al. Cytogenetic characterization of ataxia telangiectasia heterozygotes using lymphoblastoid cell lines and chronic γ -irradiation. *Hum. Genet.* 84: 532-534 (1990).
- W14 Wagner, L.K. Risk of breast cancer in ataxia-telangiectasia. *Lancet* 326: 1358 (1992).
- W15 Wang, Z.-Q., S. Saigusa and M.S. Sasaki. Adaptive response to chromosome damage in cultured human lymphocytes primed with low doses of X-rays. *Mutat. Res.* 246: 179-186 (1991).
- W16 Warenius, H.M. and J.D. Down. RBE of fast neutrons for apoptosis in mouse thymocytes. *Int. J. Radiat. Biol.* 68: 625-629 (1995).
- W17 Watson, P. and H.T. Lynch. Extracolonic cancer in hereditary nonpolyposis colorectal cancer. *Cancer* 71: 677-685 (1993).
- W18 Weeks, D.E., M.C. Paterson, K. Lange et al. Assessment of chronic γ radiosensitivity as an *in vitro* assay for heterozygote identification of ataxia-telangiectasia. *Radiat. Res.* 128: 90-99 (1991).
- W19 Wei, Y.F., P. Robins, K. Carter et al. Molecular cloning and expression of human cDNAs encoding a novel DNA ligase IV and DNA ligase III, an enzyme active in DNA repair and recombination. *Mol. Cell. Biol.* 15: 3206-3216 (1995).
- W20 Weichselbaum, R.R., D.E. Hallahan, V. Sukhatme et al. Biological consequences of gene regulation after ionising radiation exposure. *J. Natl. Cancer Inst.* 83: 480-484 (1991).
- W21 West, C.M.L., S.A.G. Elyan, P. Berry et al. A comparison of the radiosensitivity of lymphocytes from normal donors, cancer patients, individuals with ataxia-telangiectasia (A-T) and A-T heterozygotes. *Int. J. Radiat. Biol.* 68: 197-203 (1995).
- W22 Whaley, J.M. and J.B. Little. Molecular characterization of hprt mutants induced by low- and high-LET radiations in human cells. *Mutat. Res.* 243: 35-45 (1990).
- W23 Wiencke, J.K., V. Afzal, G. Olivieri et al. Evidence that the [³H] thymidine-induced adaptive response of human lymphocytes to subsequent doses of X-rays involves the induction of a chromosomal repair mechanism. *Mutagenesis* 1: 375-380 (1986).

- W24 Wiencke, J.K., J.D. Shadley, K.T. Kelsey et al. Failure of high-intensity x-ray treatments or densely-ionising fast neutrons to induce the adaptive response in human lymphocytes. p. 212 in: Proceedings of the 8th International Congress on Radiation Research, Volume 1 (E.M. Fielden et al., eds.). Edinburgh, 1987.
- W25 Winans, L.F., W.C. Dewey and C.M. Dettor. Repair of sublethal and potentially lethal X-ray damage in synchronous Chinese hamster cells. *Radiat. Res.* 52: 333-351 (1972).
- W26 Wojcik, A. and H. Tuschl. Indications of an adaptive response in C57BL mice pre-exposed *in vivo* to low doses of ionising radiation. *Mutat. Res.* 243: 67-73 (1990).
- W27 Wojcik, A. and C. Streffer. Application of a multiple fixation regimen to study the adaptive response to ionising radiation in lymphocytes of two human donors. *Mutat. Res.* 326: 109-116 (1995).
- W28 Wolff, S., R. Jostes, F.T. Cross et al. Adaptive response of human lymphocytes for the repair of radon-induced chromosomal damage. *Mutat. Res.* 250: 299-306 (1991).
- W29 Wolff, S., J.K. Wiencke, V. Afzal et al. The adaptive response of human lymphocytes to very low doses of ionising radiation: a case of induced chromosomal repair with the induction of specific proteins. p. 446-454 in: *Low Dose Radiation* (K.F. Baverstock and J.W. Stather, eds.). Taylor and Francis, London, 1989.
- W30 Wolff, R., R. Plaetke, A.J. Jeffreys et al. Unequal crossing over between homologous chromosomes is not the major mechanism involved in the generation of new alleles at VNTR loci. *Genomics* 5: 382-384 (1989).
- W31 Wooster, R., D. Ford, J. Mangion et al. Absence of linkage to the ataxia-telangiectasia locus in familial breast cancer. *Hum. Genet.* 92: 91-94 (1993).
- W32 Wouters, B. and L.D. Skarsgard. The response of a human tumor cell line to low radiation doses: evidence of enhanced sensitivity. *Radiat. Res.* 138: S76-S80 (1994).
- W33 Wyllie, A.H., J.F.R. Kerr and A.R. Currie. Cell death: the significance of apoptosis. *Int. Rev. Cytol.* 68: 251-306 (1980).
- W34 Wyllie, A.H. The genetic regulation of apoptosis. *Curr. Opin. Genet. Dev.* 5: 97-104 (1995).
- W35 Watt, P.M., E.J. Louis, R.H. Borts et al. Sgs1: a eukaryotic homolog of *E. coli* RecQ that interacts with topoisomerase II *in vivo* and is required for faithful chromosome segregation. *Cell* 81: 253-260 (1995).
- W36 Weksberg, R., C. Smith, L. Anson-Cartwright et al. Bloom syndrome: a single complementation group defines patients of diverse ethnic origin. *Am. J. Hum. Genet.* 42: 816-824 (1988).
- W37 Whitney, M., M. Thayer, C. Reifsteck et al. Microcell-mediated chromosome transfer maps the Fanconi anaemia group D gene to chromosome 3p. *Nature Genet.* 11: 341-343 (1995).
- W38 Watson, G.E., S.A. Lorimore and E.G. Wright. Long-term *in vivo* transmission of α -particle-induced chromosomal instability in murine haemopoietic cells. *Int. J. Radiat. Biol.* 69: 175-182 (1996).
- W39 Ward, J.F. DNA damage produced by ionizing radiation in mammalian cells: identities, mechanisms of formation and reparability. *Prog. Nucleic Acid Res. Mol. Biol.* 35: 95-125 (1988).
- W40 Weissenborn, U. and C. Streffer. Analysis of structural and numerical chromosomal anomalies at the first, second and third mitosis after irradiation of one-cell mouse embryos with X-rays or neutrons. *Int. J. Radiat. Biol.* 54: 381-394 (1988).
- W41 Weissenborn, U. and C. Streffer. The one-cell mouse embryo: cell cycle-dependent radiosensitivity and development of chromosomal anomalies in post-irradiation cell cycles. *Int. J. Radiat. Biol.* 54: 659-674 (1988).
- W42 Weissenborn, U. and C. Streffer. Analysis of structural and numerical chromosomal aberrations at the first and second mitosis after X irradiation of two-cell mouse embryos. *Radiat. Res.* 117: 214-220 (1989).
- W43 Ward, J.F. Molecular mechanisms of radiation-induced damage to nucleic acids. *Adv. Radiat. Biol.* 5: 181-239 (1975).
- W44 Ward, J.F. The complexity of DNA damage: relevance to biological consequences. *Int. J. Radiat. Biol.* 66: 427-432 (1994).
- W45 Wilson, D.M. and L.H. Thompson. Life without DNA repair. *Proc. Natl. Acad. Sci. U.S.A.* 94: 12754-12757 (1997).
- W46 Wilson, T.E., U. Grawunder and M.R. Lieber. Yeast DNA ligase IV mediates non-homologous DNA end joining. *Nature* 388: 495-498 (1997).
- W47 Westphal, C.H., C. Schmaltz, S. Rowan et al. Genetic interactions between *atm* and *p53* influence cellular proliferation and irradiation-induced cell cycle checkpoints. *Cancer Res.* 57: 1664-1667 (1997).
- W48 Wojcik, A., K. Bonk, W-U. Muller et al. Do DNA double-strand breaks induced by AluI lead to development of novel aberrations in the second and third post-treatment mitosis? *Radiat. Res.* 145: 119-127 (1996).
- W49 Wang, Y.A., A. Elson and P. Leder. Loss of p21 increases sensitivity to ionizing radiation and delays the onset of lymphoma in *atm*-deficient mice. *Proc. Natl. Acad. Sci. U.S.A.* 94: 14590-14595 (1997).
- W50 Wright, J.A., K.S. Keegan, D.R. Herendeen et al. Protein kinase mutants of human ATR increase sensitivity to UV and ionizing radiation and abrogate cell cycle checkpoint control. *Proc. Natl. Acad. Sci. U.S.A.* 95: 7445-7450 (1998).
- W51 Wu, L-J., G. Randers-Pehrson, A. Xu et al. Targeted cytoplasmic irradiation with alpha particles induces mutations in mammalian cells. *Proc. Natl. Acad. Sci. U.S.A.* 96: 4959-4964 (1999).
- W52 Wood, R.D. Nucleotide excision repair in mammalian cells. *J. Biol. Chem.* 272: 23465-23468 (1997).
- W53 Wishkerman, V.Y., M.R. Quastel, A. Douvdevani et al. Somatic mutations at the glycoprotein A (GPA) locus measured in red cells of Chernobyl liquidators who immigrated to Israel. *Environ. Health Perspect.* 105: 1451-1454 (1997).
- X1 Xia, F., S.A. Amundson, J.A. Nickoloff et al. Different capacities for recombination in closely related human lymphoblastoid cell lines with different mutational responses. *Mol. Cell. Biol.* 14: 5850-5857 (1994).
- X2 Xia, F., X. Wang, Y-H. Wang et al. Altered p53 status correlates with differences in sensitivity to radiation-induced mutation and apoptosis in two closely related human lymphoblast lines. *Cancer Res.* 55: 12-15 (1995).
- X3 Xu, Y., T. Ashley, E.B. Brainerd et al. Targeted disruption of *ATM* leads to growth retardation, chromosomal fragmentation during meiosis, immune defects, and thymic lymphoma. *Genes Dev.* 10: 2411-2422 (1996).
- X4 Xu, Y. and D. Baltimore. Dual roles of ATM in the cellular response to radiation and in cell growth control. *Genes Dev.* 10: 2401-2410 (1996).
- X5 Xanthoudakis, S., G. Miao, F. Wang et al. Redox activation of Fos-Jun DNA binding activity is mediated by a DNA repair enzyme. *EMBO J.* 11: 3323-3335 (1992).

- X6 Xanthoudakis, S., R.J. Smeyne, J.D. Wallace et al. The redox/DNA repair protein, Ref-1, is essential for early embryonic development in mice. *Proc. Natl. Acad. Sci. U.S.A.* 93: 8919-8923 (1996).
- X7 Xu, X., K-U. Wagner, D. Larson et al. Conditional mutation of *Brcal* in mammary epithelial cells results in blunted ductal morphogenesis and tumour formation. *Nature Genet.* 22: 37-43 (1999).
- X8 Xu, X., Z. Weaver, S.P. Linke et al. Centrosome amplification and a defective G2-M cell cycle checkpoint induce genetic instability in BRCA1 exon 11 isoform-deficient cells. *Mol. Cell* 3: 389-395 (1999).
- Y1 Ying, K.L. and W.E. Decoteau. Cytogenetic anomalies in a patient with ataxia, immune deficiency and high α -fetoprotein in the absence of telangiectasia. *Cancer Genet. Cytogenet.* 4: 311-317 (1981).
- Y2 Yandell, D.W., T.P. Dryja and J.B. Little. Somatic mutations at a heterozygous autosomal locus in human cells occur more frequently by allele loss than by intragenic structural alterations. *Somat. Cell Mol. Genet.* 12: 255-263 (1986).
- Y3 Yamada, T. and H. Ohyama. Radiation-induced interphase death of rat thymocytes is internal programmed (apoptosis). *Int. J. Radiat. Biol.* 53: 65-75 (1988).
- Y4 Yamaguchi, T. Relationship between survival period and dose of irradiation in rat thymocytes *in vitro*. *Int. J. Radiat. Biol.* 12: 235-242 (1967).
- Y5 Yanagihara, K., M. Nii, M. Numoto et al. Radiation-induced apoptotic cell death in human gastric epithelial tumour cells; correlation between mitotic death and apoptosis. *Int. J. Radiat. Biol.* 67: 677-685 (1995).
- Y6 Yoshimura, Y., T. Morita, A. Yamamoto et al. Cloning and sequence of the human *RecA*-like gene cDNA. *Nucleic Acids Res.* 21: 1665 (1993).
- Y7 Yamashita, T., D.L. Barber, Y. Zhu et al. The Fanconi anaemia polypeptide FACC is localized to the cytoplasm. *Proc. Natl. Acad. Sci. U.S.A.* 91: 6712-6716 (1994).
- Y8 Yuille, M.A., L.J. Coignet, S.M. Abraham et al. ATM is usually rearranged in T-cell prolymphocytic leukaemia. *Oncogene* 16: 789-796 (1998).
- Z1 Ziv, Y., A. Amiel, N.G.J. Jaspers et al. Ataxia-telangiectasia: a variant with altered *in vitro* phenotype of fibroblast cells. *Mutat. Res.* 210: 211-219 (1989).
- Z2 Ziv, Y., G. Rotman, M. Frydman et al. The ATC (ataxia-telangiectasia complementation group C) locus localizes to 11q22-q23. *Genomics* 9: 373-375 (1991).
- Z3 Ziv, Y., M. Frydman, E. Lange et al. Ataxia-telangiectasia -linkage analysis in highly inbred arab and Druze families and differentiation from an ataxia-microcephaly-cataract syndrome. *Hum. Genet.* 88: 619-626 (1992).
- Z4 Ziegler, A., A.S. Jonason, D.J. Leffell et al. Sunburn and p53 in the onset of skin cancer. *Nature* 372: 773-776 (1994).
- Z5 Zdzienicka, M.Z. and J.W.I.M. Simons. Mutagen-sensitive cell lines are obtained with a high frequency in V79 Chinese hamster cells. *Mutat. Res.* 178: 235-244 (1987).
- Z6 Zdzienicka, M.Z., N.G.J. Jaspers, G.P. van der Schans et al. Ataxia-telangiectasia-like Chinese hamster V79 cell mutants with radioresistant DNA synthesis, chromosomal instability and normal DNA strand break repair. *Cancer Res.* 49: 1481-1485 (1989).
- Z7 Zhen, W., C.M. Denault, K. Loviscek et al. The relative radiosensitivity of TK6 and WI-L2-NS lymphoblastoid cells derived from a common source is primarily determined by their p53 mutational status. *Mutat. Res.* 346: 85-92 (1995).
- Z8 Zhou, P.-K., W-Z. Sun, X-Y. Liu et al. Adaptive response of mutagenesis and DNA double-strand break repair in mouse cells induced by low dose of γ -ray. p. 271-274 in: *Low Dose Irradiation and Biological Defense Mechanisms* (T. Sugahara et al., eds.). Elsevier, 1992.
- Z9 Zhu, C., M.A. Bogue, D-S. Lim et al. Ku-86 deficient mice exhibit severe combined immunodeficiency and defective processing of V(D)J recombination intermediates. *Cell* 86: 379-389 (1996).
- Z10 Zhang, Q-M., S. Yonei and M. Kato. Multiple pathways for repair of oxidative damages caused by X rays and hydrogen peroxide in *Escherichia coli*. *Radiat. Res.* 132: 334-338 (1992).
- Z11 Zasukhina, G.D., T.A. Sinel'shchikova, I.M. Vasil'eva et al. Radioadaptive response in repair-defective cells from patients with homocystinuria. *Genetika* 32: 1592-1595 (1996).
- Z12 Zhong, Q., C-F. Chen, S. Li et al. Association of BRCA1 with the hRad50-hMre11-p95 complex and the DNA damage response. *Science* 285: 747-750 (1999).
- Z13 Zdzienicka, M.Z. Mammalian X ray sensitive mutants: a tool for the elucidation of the cellular response to ionizing radiation. *Cancer Surv.* 28: 281-293 (1996).
- Z14 Zhang, H., R.L. Zheng, Z.Q. Wei et al. Effects of pre-exposure of mouse testis with low dose $^{16}\text{O}8^+$ ions or ^{60}Co gamma-rays on sperm shape abnormalities, lipid peroxidation and superoxide dismutase activity induced by subsequent high dose irradiation. *Int. J. Radiat. Biol.* 73: 163-167 (1998).